

### STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If Inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If On Skin: Remove contaminated clothing and wash affected areas with soap and water.

If In Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Physician: This product may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and pot-holes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### REENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written or oral warnings must include the following information:

Do not contaminate water, food, or feed by storage or disposal.  
**STORAGE:** Store in cool, dry, well-ventilated area. Store separately from strong alkalis and strong oxidizers. Keep container tightly closed when not in use.  
**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticides, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.  
**CONTAINER DISPOSAL:** Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved of by state and local authorities.

### DIRECTIONS FOR DILUTION

**NOTE:** The rates of water necessary to add to the tank will vary with the application method. The normal requirements for the following application methods are as follows:  
**AIR APPLICATION—**5 to 20 gals. per acre. **GROUND APPLICATION—**15 to 200 gals. per acre. **ORCHARD APPLICATION—**70 to 800 gals. per acre. (Citrus may require up to 1,500 gals.) (70 to 100 gals. per acre rates are considered concentrate rates and require special concentrate equipment.)

**CHEMGATION—**Refer to supplemental labeling entitled "APPLICATION THROUGH IRRIGATION SYSTEMS—CHEMGATION" for use directions for chemigation. Do not apply this product through any irrigation systems unless the supplemental labeling on chemigation is followed.

### APPLICATIONS

**BEANS:** Lygus Bug, Aphid. 1 to 1 1/4 pts./acre. Do not apply within 1 day of harvest.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** Aphid, Cabbage Looper, Imported Cabbage Worm. 1 to 2 1/2 pts./acre. Do not apply within 3 days of harvest on Broccoli, or within 7 days on Cabbage, Brussels Sprouts or Cauliflower.

**LETTUCE, SPINACH:** Aphid, Leafhopper. 1 to 2 pts./acre. Do not apply within 14 days of harvest on Leaf Lettuce, or within 7 days of harvest on Head Lettuce or Spinach.

**ONIONS (including Green Onions):** Onion Thrip. 1 to 2 pts./acre. Do not apply within 3 days of harvest on Green Onions.

Size 17 1/4 x 8 3/4

# MALASPRAY 8EC

### ACTIVE INGREDIENT:

Malathion (O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate)	80%
INERT INGREDIENTS*	20%
<b>TOTAL</b>	<b>100%</b>

\*Contains xylene range aromatics

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Side Panel For Additional Precautionary Statements

NET CONTENTS \_\_\_\_\_ GALLONS

EPA REG. NO. 34704-458

EPA EST. NO. \_\_\_\_\_



25888

05788 A

**WALNUTS:** Walnut Husk Weevils.  
**EGGPLANT:** Aphid, Leafminer. Do not apply within 14 days of harvest.  
**PEAS:** Pea Aphid, Lygus Weevil. 1 to 2 1/2 pts./acre harvest.  
**CORN:** Aphid, Corn Earworm, Young Grasshopper within 5 days of harvest may occur in the whorls.  
**HOPS:** Aphids. 1/2 to 1 1/2 pts./acre.  
**CUCUMBERS, SQUASH:** Spider Mite, Pickleworm, Dotted Ground Squirrel. Do not apply within 1 day of harvest.  
**POTATOES:** Potato Aphid, Bug, Leafhopper, Mealworm.  
**TOMATOES:** Aphid, Spider Mite. Apply within 1 day of harvest.  
**ALFALFA, CLOVER:** Alfalfa Weevil Larvae, Grasshopper, Lygus Bug to 1 1/2 pts./acre. No time in bloom.  
**APPLES:** Red-banded Leafminer, Bud Mot. 1 1/2 pts./acre per 100 gals. water. May cause injury if applied within 4 weeks of bloom.  
**CHERRIES:** Black Cherry Aphid. Do not apply within 14 days of harvest. May injure foliage on some varieties.  
**PEARS:** Red-banded Leafminer, Woolly Aphid, Mealybug. 1 1/2 pts./acre per 100 gals. water. Do not apply within 7 days of harvest.  
**PEACHES:** Spider Mite. 1 1/2 pts./100 gals. water. Do not apply within 7 days of harvest.  
**PLUMS, PRUNES:** Mealworm. 1/2 pt./100 gals. water. Do not apply within 3 days of harvest.  
**GRAPES:** Grape Leafhopper, Spider Mite, Mealybug. Do not apply within 3 days of harvest.  
**STRAWBERRIES:** Aphid, Leafroller, Strawberry Weevil.

**BEST AVAILABLE COPY**