



SUPER 94 440 SPRAY OIL

ACTIVE INGREDIENT:	BY WT.
Petroleum Oil	99%
INERT INGREDIENTS:	1%
TOTAL	100%

Petroleum Oil Minimum Unsulfonated Residue—99%
Petroleum Oil Classification: Unclassified

neoprene gloves and overboes, goggles and face protection.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in a cool, dry area, away from heat and open flame.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

KEEP OUT OF REACH OF CHILDREN CAUTION

See Below For Additional Precautionary Statements.

EPA REG. NO. 34704-464

EPA EST. NO. _____

NET CONTENTS _____ GALLONS

07/93

ACCEPTED

SEP - 8 1993

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION

HARMFUL IF SWALLOWED.

DO NOT INDUCE VOMITING.

CALL A PHYSICIAN IMMEDIATELY.

AVOID BREATHING OF SPRAY MIST.

READ ENTIRE LABEL. Use strictly in accordance with label cautions, warnings and directions; and conformity with Federal and State Regulations.

STATEMENT OF PRACTICAL TREATMENT

- If swallowed: Do not induce vomiting. Call a physician immediately.
- If in eyes: Wash thoroughly with water and if irritation persists, consult a physician.
- If inhaled: Remove victim to fresh air. Apply respirator if indicated.
- If on skin: Remove contaminated clothing and wash affected area with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Apply this product only as specified on the label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until the spray has dried. Certain states may require more restrictive reentry intervals for various crops treated with this product. Consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written and oral warnings must include the following information:

CAUTION

Area treated with Petroleum Oil on (date of application). Do not enter without appropriate protective clothing until spray has dried. Required Protective Clothing: long-sleeved shirt, long-legged pants,

APPLICATIONS

This spray oil is a straight-cut narrow boiling range petroleum-oil designed specifically for use on deciduous fruit trees.

Apply so as to give total coverage to all parts of tree. Thorough drenching of trunks, limbs, twigs and leaves is essential for maximum insect control.

This product may be used as a carrier for appropriate insecticides and fungicides which are registered for use on deciduous fruits and nuts and which can be applied to give a uniform spray. Consult your local Agricultural Extension Service.

DO NOT COMBINE WITH SULFUR SPRAYS AND DO NOT APPLY spray oil within 30 days before or after an application of sulfur.

DO NOT MIX with insecticides or fungicides incompatible with oil. Consult your State Agricultural Extension Service for specific compatibilities. DO NOT APPLY to trees weakened by disease, drought, drying winds, or high nitrogen applications.

DO NOT USE this product when temperatures are above 90°F.

Use only in spray machines with powerful positive agitation. Keep agitator running at all times during spray operations. This product is formulated as a "fast break" spray oil and will not stay in solution unless agitated constantly.

Do not apply this product through any type of irrigation system.

Unless otherwise noted, amounts given are for 100 gallons of diluted spray. See specific instructions for concentrate and aerial application.

Fill spray tank with 1/2 the required amount of water. Start agitation. Add total amount of spray oil. Continue operation of agitator while adding balance of water and while spraying.

ALMONDS, APRICOTS, APPLES, CHERRIES, HECTARINES, OLIVES, PEACHES, PEARs, PLUMS, PRUNES: Dormant Treatment: Brown Apricot Scale, Black Scale, San Jose Scale, Parasitic Scale, Oleander Scale, Italian Pear Scale, Lecanium Scale, Aphid Eggs, Spider Mite. Apply 1 1/2 to 2 gallons per 100 gallons of water. Delayed Dormant Treatment: Pear Psylla. Apply 2 gallons per 100 gallons of water. Summer Treatment: Spider Mite. Apply 3 to 4 quarts per 100 gallons of water. San Jose Scale, Italian Pear Scale, Pear Psylla. Apply 1 to 1 1/2 gallons per 100 gallons of water. Aerial Application: Dormant and Delayed Dormant Application Only: Apply 3-5 gallons in a minimum of 15 gallons of water per acre. Apply by air only when soil conditions do not permit regular ground application.

WALNUTS (summer treatment only): Spider Mite. Apply 3 to 4 quarts per 100 gallons of water. Summer stages of Frost Scale, European Fruit Scale, San Jose and other soft-shelled Scale: Apply 1 to 1 1/2 gallons per 100 gallons of water. Do not apply after harvest.

ORCHARD CONCENTRATE APPLICATION: Apply 3 to 5 gallons in a minimum of 50 gallons of water per acre. Use only on crops listed on this label. Special concentrate spray equipment is necessary for this application.

CITRUS: California Only (Lemons, Grapefruit and Oranges (Navel and Valencia)): Black Scale, Brown Soft Scale, California Red Scale, Citricola Scale, Citrus Bud Mite, Citrus Red Mite, Purple Scale, Yellow Scale, Citrus Flat Mite. Central California: Apply 0.5 to 1.0 gallons per

Post Harvest and

BEST AVAILABLE COPY

39
2

SUPER 94 440 SPRAY OIL

EPA REG. NO. 34704-464

100 gallons of water as a thorough coverage spray (TC). For Lemons apply in August and September only; for Grapefruit apply July through September; and for Oranges (Navels and Valencia) apply July through September. Southern California: For coastal Lemons, apply 0.5 to 1.4 gallons per 100 gallons of water in May and June and/or September through December. Apply as a thorough coverage spray (TC). For interior Lemons, apply 0.5 to 1.4 gallons per 100 gallons of water in April and May and/or September through November. Apply as a thorough coverage spray (TC). For Lemons, Grapefruit and Oranges (Navels and Valencia), apply to 10 gallons per acre on bearing trees up to 10 feet in height. Add 1 gallon of oil per acre for each foot of tree height over 10 feet. Apply as a low volume spray (LV). For Lemons, make applications in all months except December through February; for Grapefruit make applications July through October; for Oranges make applications on Navels July through September and on Valencia July through October. Grand Plains Area: For Lemons, Grapefruit and Oranges (Navels and Valencia), follow the application rates stated for Southern California but delete timing restrictions. Make applications as needed throughout the year.

GRAPES: Delayed Dormant or Summer Applications: Mealybug. Apply 1 gallon plus the appropriate label rate of an insecticide registered for use on grapes per 100 gallons of water. Do not apply over 250 gallons of water.


NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNWANTED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.

BEST AVAILABLE COPY

FORMULATED FOR
PLATTE CHEMICAL CO.
156 SO. MAIN STREET FREMONT, NEBRASKA 68025-6467



Super 94 Spray Oil (aka Super 94 400 Spray Oil)
EPA Reg. No. 34704-464

The following statement is the proposed wording for the replacement of the present sentence highlighted on the current label.

DO NOT COMBINE WITH SULFUR SPRAYS AND DO NOT APPLY spray oil within 30 days before or after an application of sulfur except in the Northwest (Oregon and Washington) where it may be used in combination on Pears as a Post Harvest (after all fruit is off the tree), Dormant and Delayed Dormant (before the scales slip, or at or prior to bud swell) treatment and on Apples prior to or up to the Delayed Dormant (1/2" green) period.