

PH. 12 2 2000-93

15

493225

AUG 29 1995

Ms. Glenda Haage
Platte Chemical Company
P.O. Box 667
Greeley, CO 80632-0667

Dear Ms. Haage:

Subject: Revised Confidential Statements of Formula, and
Finished labeling per Agency letter dated July 19, 1995
Diazinon 5 Granules
EPA Registration Number 34704-493
Your submissions dated August 1, 1995

As noted in our telephone conversation of August 28, 1995, the basic and alternate Confidential Statements of Formula (CSFs) dated July 31, 1995 show certified limits for the impurities of toxicological concern which are not in agreement with your suppliers' figures for these impurities. The Agency understand that you will soon be submitting revised basic and alternate CSFs bearing corrected certified limits.

The labeling referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable and a stamped copy is enclosed for your records. At the next label printing, please add the missing comma after "(ducks or geese)" in the Environmental Hazards.

Sincerely,

George T. LaRocca
Product Manager (13)
Insec.icide-Rodenticide Branch
Registration Division (7505C)

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS.

ACCEPTED

AUG 29 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 34704-493

DIAZINON 5 GRANULES

FOR OUTDOOR USE ON THE HOME LAWN AND IN THE HOME GARDEN ONLY

ACTIVE INGREDIENT:	BY WT.
Diazinon: O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	5.0%
INERT INGREDIENTS:	95.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

EPA REG. NO. 34704-493

EPA EST. NO. _____

NET WEIGHT 50 LBS. (22.68 KG)

4777

07Y95

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl (ducks or geese) feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds. Avoid overlapping granules. Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil. For lawns, follow all watering instructions in the directions for use, however, stop watering before puddling occurs. For lawns, do not apply this product to sites occupied by waterfowl, or within 75 ft. of any body of water which may attract waterfowl, such as ponds or lakes. Do not apply directly to water. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Runoff may be hazardous to fish and other aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration, if indicated.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physicians: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a cool, dry, well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-303-358-4400 or 1-800-424-9300 day or night.

In Lawns this product controls:

Ants	Crickets	Sod webworms
Armyworms	Cutworms	Sowbugs
Bermudagrass	Digger wasps	(Pillbugs)
mites	Earwigs	Springtails
Billbugs	Fire Ants	White grubs of:
Brown dog ticks	Fleas	* European chaffer
Chiggers	Lawn moths	* Japanese beetle
Chinch bugs	Leathoppers	* Southern chaffer
Clover mites	Millipedes	* Aids in control

In Gardens this product controls:

Cutworms	Onion maggots	Wireworms
Mole crickets	Root maggots	

OVERALL LAWN APPLICATION

Special Precautions To Reduce Risk To Birds

Birds, especially waterfowl (ducks or geese) feeding or drinking on treated areas may be killed. To reduce risks to birds:

1. Do not treat areas larger than 15,000 sq. ft.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20.
3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
4. Avoid overlapping treated areas.
5. Follow all watering instructions in the directions for use, however, stop watering before puddling occurs.
6. Do not apply this product to sites occupied by waterfowl or to sites within 75 ft. of any body of water that may attract waterfowl, such as lakes or ponds.
7. If it is necessary to repeat application, wait a minimum of 3 weeks between applications. Make no more than 4 applications per year.

This product controls certain lawn insects on both grass and dichondra lawns. Apply this product with fertilizer spreader. Refer to the tables that follow for use rates and spreader settings.

DIAZINON 5 GRANULES

EPA REG. NO. 34704-493

Pest	Rate*	Timing/Special Directions
Ants, Armyworms, Bermudagrass mites, Brown dog ticks, Chiggers, Chirch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Leafhoppers, Millipedes, Sowbugs (Pillbugs), Springtails (Collembola)	2 lbs. per 1,000 sq. ft.	Apply when grass is dry. Water lawn lightly after application. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year.
Billbugs, White grubs of Japanese beetle, European chafer, Southern chafer, <i>Atractodes pretulius</i> (dung beetle)	2 lbs. per 1,000 sq. ft.	Water grass with at least 1/4 inch of water after application, however, stop watering if puddling occurs. For billbug control, apply when activity is first observed or when chewed grass indicates damage. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or Summer when damage is first noticed. Water grass with at least 1/4 inch of water application, however, stop watering if puddling occurs. Treat again in late Summer or early Fall, if a second generation occurs.
Hyperodes weevils	2 lbs. per 1,000 sq. ft.	Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. Treat problem areas in mid-April and again in mid-May.

*1,000 sq. ft. = an area 20 ft. x 50 ft.

100 sq. ft. = an area 10 ft. x 10 ft.

2 lbs. per 1,000 sq. ft. = 3.2 oz. per 100 sq. ft.

Note: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting, if necessary in order to assure correct rate of application.

LAWN SPREADER SETTING NO. ** (To deliver 2 lbs. per 1,000 sq. ft.)

Spreader Make and Model	Setting
Central Imperial Mark No. 1662-G	6 1/2
Central Suburbia Lawn Spreader No. 1622-0	6 1/2
Cyclone Model B1 Rotary	4 1/2
Greenfield Model 105	6 1/2
ORTHO Drop Spreader Model 3000	4 1/2
ORTHO Broadcast Spreader Model 4000	4 1/2
J.C. Penney Model 6005	7 1/4
J.C. Penney Rotary Spreader	5
Scott Precision Flow Spreader Model PF-4	4
Sears Craftsman Model 871.19198	6

** For application with other types of equipment, apply at the rate of 2 lbs. per 1,000 sq. ft.

LAWN SPOT TREATMENT

Ants: Apply 1 1/2 teaspoons of this product over and around each ant hill.

Fire ants: To aid in the control of fire ants in lawns and other outdoor home recreation areas, apply 1/2 cup over, and 2 ft. around each fire ant mound. Immediately after application, water in thoroughly. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. A minimum of 1 gal. of water should be applied to each mound area. Reapply as necessary.

Note: For best results, apply in cool weather, 65-80° F. or in early morning or late evening hours. Works within 24 hours. No walking for days or weeks. Treat new mounds as they appear.

Digger wasps: Apply 1 1/2 teaspoons of this product in and around each nest opening. Apply at dusk when wasps are not active.

BAND TREATMENT AROUND HOUSE FOUNDATION

The following pests are commonly found around and near house foundations, and may enter the house:

Ants	Crickets	Millipedes
Brown dog ticks	Earwigs	Sowbugs (Pillbugs)
Clover mites	Fleas	Springtails (Collembola)

Apply a 2-foot band of this product next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (2' x 50') treated. If necessary, repeat application at 3-week intervals.

VEGETABLES — INSECT CONTROL

Special Precautions To Reduce Risk To Birds

- Birds feeding on treated areas may be killed. To reduce risk to birds:
1. Thoroughly mix this product into the soil during or immediately after application. Refer to the Soil Incorporation table that follows.
 2. Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil.

This product controls certain garden insects on crops listed in the following tables. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate only for heavier infestations.

Do not apply this product to food crops grown in greenhouses.

Applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.

SOIL INCORPORATION

Insect	Depth of Mixing or Incorporation into Soil
Surface cutworms*	2-3 inches
Subterranean cutworms*	3-6 inches
Mole crickets	1-2 inches
Root maggots	3-4 inches
Wireworms	4-8 inches

*If type of cutworm cannot be identified, mix into soil 4-6 inches deep.

Crop	Pest	Rate	Timing/Special Directions
Beans (Lima, Pole and Snap) (succulent only)	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Beets, Red (Table)	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
	Mole Crickets	7 oz. per 1,000 sq. ft.	
Broccoli	Root Maggots	14-21 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	14-28 oz. per 1,000 sq. ft.	
	Wireworms	21-28 oz. per 1,000 sq. ft.	
	Mole Crickets	7 oz. per 1,000 sq. ft.	
Brussels Sprouts	Root Maggots	14-21 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	14-28 oz. per 1,000 sq. ft.	
	Wireworms	21-28 oz. per 1,000 sq. ft.	

DIAZINON 5 GRANULES
EPA REG. NO. 34704-493

Crop	Pest	Rate	Timing/Special Directions
Cabbage	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Root Maggots	14-21 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Carrots	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Cauliflower	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Root Maggots	14-21 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Celery	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Collards	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Cucumbers	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Endive (Escarole)	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Kale	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Lettuce	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	

Crop	Pest	Rate	Timing/Special Directions
Melons (Cantaloupes, Casabas, Cranshaws, Honeydew, Musk-melons, Persians and Hybrids of these) and Watermelons	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Mustard	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Onions (Bulb & Green)	Onion Maggots	14-28 oz. per 1,000 sq. ft.	Broadcast just before planting and mix into the top 3-4 inches of soil. NOTE: Resistant onion maggots will not be controlled by diazinon or similar type insecticides.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Parsley	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Peas (succulent only)	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Note: Do not use on dried pea varieties such as dried blackeyed peas.			
Peppers	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
	Mole Crickets	7 oz. per 1,000 sq. ft.	
Potatoes	Cutworms (Surface & Subterranean)	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Radishes	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Spinach	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Squash (Summer and Winter)	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Sweet Corn	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
	Seed Corn Maggots	14-28 oz. per 1,000 sq. ft.	
Sweet Potatoes	Wireworms, Flea Beetles	21 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
Swiss Chard	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
Tomatoes	Cutworms	14-28 oz. per 1,000 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	
	Mole Crickets	7 oz. per 1,000 sq. ft.	
Turnips	Mole Crickets	7 oz. per 1,000 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	21-28 oz. per 1,000 sq. ft.	

DIAZINON 5 GRANULES

EPA REG. NO. 34704-493

NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

FORMULATED FOR
PLATTE CHEMICAL CO.

150 SO. MAIN STREET FREMONT, NEBRASKA 68025-5697