

PM 12 34704-494

RESTRICTED USE PESTICIDE

Due to very high acute toxicity to humans and birds.

For use only by persons who have been trained and certified in the use of restricted use pesticides. For more information on restricted use pesticides, contact your state pesticide applicator or the National Pesticide Information Center, 1515 Greenburg Road, Corvallis, Oregon 97331, (503) 852-8900.

ETHYL METHYL PARATHION 6-3 EC

ACTIVE INGREDIENTS	BY WT
ETHYL METHYL PARATHION	6.3
INERT INGREDIENTS	93.7
TOTAL	100.0

EPA Reg. No. 106-11-01
EPA Est. No. 106-11-01



POISON

KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

PRECAUTIONARY STATEMENTS

POISONOUS Symptoms: Nausea, vomiting, diarrhea, dizziness, headache, blurred vision, difficulty breathing, weakness, and loss of consciousness. If symptoms occur, seek medical attention immediately.

STATEMENT OF PRACTICAL TREATMENT

If breathing has stopped, artificial respiration should be given. Do not use mouth-to-mouth respiration. If necessary, use a resuscitator. Do not induce vomiting.

If swallowed: Do not induce vomiting. Give water or milk to drink. Do not give anything to eat or drink until a doctor has been consulted.

In case of contact: Wash thoroughly with soap and water. Remove contaminated clothing and shoes. Do not use solvents to clean skin.

NOTE TO PHYSICIAN

For information on the toxicity of this pesticide, contact the National Pesticide Information Center, 1515 Greenburg Road, Corvallis, Oregon 97331, (503) 852-8900. For information on the treatment of poisoning, contact your local poison control center.

Wear long pants and coat, heavy-duty chemical-resistant gloves, a hood or wide-brimmed hat, safety goggles or face shield, and a chemical-resistant respirator. In addition, applicators must wear a chemical-resistant apron when using the restricted product. During aerial application, an enclosed hood or a helmet with a visor may be substituted for the hood or wide-brimmed hat and safety goggles or face shield requirements. IF MIXING AND LOADING IS PERFORMED USING A CLOSED SYSTEM, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE:

Heavy-duty chemical-resistant gloves, chemical-resistant apron, long-sleeved shirt or pants and short-sleeved shirt and long-legged pants, shoes and socks.

Safety goggles or a face shield must be worn when the system is under pressure. A other protective clothing and equipment required for use with such systems must be available nearby.

IF APPLICATION IS PERFORMED USING AN ENCLOSED CAB OR EQUIPMENT, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATIVE:

Clean long-sleeved shirt and long-legged pants. An other protective clothing and equipment may be used during application if it is available in the cab and must be worn when exiting the cab to treat areas not used for this purpose. Contaminated clothing may not be worn back into the cab unless it is enclosed, such as a plastic bag. REMEMBER: THE FOLLOWING IS NOT INTENDED TO PROTECT OPERATORS FROM REPAIR AND CLEANING OF APPLICATION EQUIPMENT. REPAIR AND EARLY RE-ENTRY REFER TO THE INSTRUCTIONS ON THE LABEL.

IF WASHING CLOTHES ARE STRICTLY PROHIBITED DURING APPLICATION.

IMPORTANT: Do not come in contact with skin, wash thoroughly with soap and water immediately. Always wash hands, face and arms with soap and water before smoking, eating, or drinking.

AFTER WORK: Wash gloves with soap and water before entering the house. Wash clothes and shoes. Store protective clothing separately from other clothing. Wash protective clothing separately from other clothing with soap and water. Aerator's and sprayer's parts after use should be washed with soap and water. Do not use for other purposes. Do not use for other purposes. Do not use for other purposes. Do not use for other purposes.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to birds and fish. Birds in treated areas should be kept out and other aquatic organisms should be protected. Do not apply to water bodies. Do not apply to water bodies. Do not apply to water bodies.

RE-ENTRY STATEMENT

Do not re-enter treated areas until the pesticide has dried. Do not re-enter treated areas until the pesticide has dried. Do not re-enter treated areas until the pesticide has dried.

STORAGE AND DISPOSAL

Store in a cool, dry place. Do not store in a hot or cold place. Do not store in a hot or cold place. Do not store in a hot or cold place.

BEST AVAILABLE COPY

Spinach Aphids, Leafminers, Apply 1.4 to 2.0 pint per acre. Armyworm, Apply 1.0 to 1.5 pint per acre. Flea Beetle, Marengo Bug, Leafhoppers, Cabbage Looper, Mites, Seed Corn Maggot, in corn. Mites, Vegetable Weevils, Wireworm, Apply 2.0 pint per acre. Rates up to 1.0 pint, do not apply within 14 days of harvest. Rate 1.0 pint through 2.0 pint, do not apply within 21 days of harvest.

Strawberries Aphids, Cicadas, Flea Beetles, Leafhoppers, Leafminers, Lygus Bugs, Spider Mites, Pomegranate Thrips, Apply 1.0 to 2.0 pint per acre. Do not apply within 14 days of harvest.

Sugar Beets Leafhoppers, Leafminers, Lygus Bugs, Apply 1.0 to 2.0 pint per acre. Aphids, Armyworm, Baster Beetle, Cucumber Cutworm, Flea Beetle, Grasshoppers, Mite, Looper, Mites, Stink Bugs, weevils, Apply 2.0 pint per acre. Do not apply within 20 days of harvest. Thrips are not be fed. If thrips are going to be fed, do not apply within 50 days of harvest.

Sweet Potatoes Aphids, Leafhoppers, Leafminers, Spider Mites, Stink Bugs, Apply 2.0 to 3.4 pint per acre. Do not apply within 14 days of harvest.

Tobacco Aphids, Flea Beetles, Sucking, Apply 1.2 pint per acre. Do not apply within 5 days of curing. Avoid contact with plant surfaces when priming or curing tobacco. When necessary to enter treated fields within 24 hours after application, protective clothing should be worn.

Tomatoes Aphids, Baster Beetle, Colorado Potato Beetle, Flea Beetle, Grasshoppers, Leaf Footed Bugs, Leafhoppers, Leafminers, Spider Mites, Stink Bugs, Thrips, Wire Weevils, Apply 1.0 to 2.0 pint per acre. Armyworm, Banded Cucumber Beetle, Looper, Mites, Plant Borer, Tomato Pinworm, Vegetable Weevils, Apply 2.0 to 3.4 pint per acre. Do not apply within 10 days of harvest. Rates over 3.4 pint per acre, 1.4 pints, do not apply within 14 days of harvest.

Turnips Aphids, Armyworm, Cucumber Flea Beetle, Mite, Flea Beetle, Spider Bug, Flea Beetle, Marengo Bug, Important Cabbage Looper, Mites, Vegetable Weevils, Cabbage Weevils, Apply 1.0 to 2.0 pint per acre. Leafminers, Cabbage Looper, Apply 2.0 pint per acre. Do not apply within 21 days of harvest. Thrips are used by the grower, apply within 14 days of harvest. Thrips are not used by the grower.

Simplot

J. R. Simplot Company

P.O. Box 198, Lathrop, California 95330

... ..

APR 1964

BEST AVAILABLE COPY

11/14