

DICOFOL 4 EC

ACTIVE INGREDIENT:

1,1-Dichloro-2,2-bis(4-chlorophenyl)ethane

WT. %

41.1%

INERT INGREDIENTS:

58.9%

TOTAL

100.0%

Contains 4 lbs. Dicofol per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See side panels for additional Precautionary Statements
and Directions for Use.

EPA REG. NO. 34704-513

EPA EST. NO. _____

NET CONTENTS 2 1/2 GALLONS



FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
150 SO. MAIN STREET FREMONT, NEBRASKA 68725

23115

11128 A

103

BEST AVAILABLE COPY

2007

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Avoid skin contact. Wash thoroughly after using. Avoid inhalation of mist. Harmful if swallowed. If taken internally get medical attention. Do not store near food or foodstuffs.

STATEMENT OF PRACTICAL TREATMENT:

IF SWALLOWED call a physician immediately. If no other treatment is available induce vomiting by giving victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED remove victim to fresh air. Apply respiration if breathing is irregular or has stopped. Call a physician immediately.

IF ON SKIN wash immediately with soap and water.

IF IN EYES flush eyes for at least 15 minutes with water. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water.



CHEMICAL HAZARDS

NOTE—This product must be stored at temperatures above 40°F. Storage below may affect the quality of the product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Skin contact with this pesticide may be hazardous: wear chemical resistant gloves when mixing, loading, or applying this product.

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written and oral warnings must include the following information.

CAUTION

Area treated with DIOFOL 4EC on (date of application) Do not enter without appropriate protective clothing until sprays have dried.

REQUIRED PROTECTIVE CLOTHING: Long sleeved shirt, long legged pants, rubber gloves and a respirator approved by the Mining, Environment and Safety Administration and by the National Institute for Occupational Safety and Health.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

STORAGE: Store at temperatures above 40°F. Storage below may affect the quality of the product.

CONTAINER DISPOSAL: Metal: triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Do not apply this product through any type of irrigation system.

DIRECTIONS FOR DILUTION

DILUTE APPLICATION: Ground Application: Apply specified rate in 20 to 60 gallons of water per acre. Orchard Application: Apply specified rate in 100 to 800 gallons of water per acre.

CONCENTRATE APPLICATION: Ground Application: Apply specified rate in not less than 5 gallons of water per acre. Orchard Application: Apply specified rate in 20 to 100 gallons of water per acre. Special concentrate equipment is necessary for these uses.

AIR APPLICATION: Apply specified rate in 5 to 20 gallons of water per acre. Orchard rates should not be applied in less than 10 gallons of water per acre.

Apples, Pears (Western states only): Apple Rust, Brown, European Red, McDaniels Spider, Pacific, Schoene Spider, Willamette and Yellow Spider Mites. Apply 1 1/2 to 4 qts. per acre. Do not apply within 7 days of harvest. Dicofol may cause fruit spotting under certain conditions and should not be used in late cover spray. Claims for Brown Mite and European Red Mite are for cleanup of late spring populations. Repeat as needed but allow 30 days between treatments.

Beans: Carmine Spider, Strawberry Spider, Pacific Spider, Two-Spotted Mite. Apply 1/2 to 1 1/2 qts. per acre other than California. Apply 7 days before harvest through 1/2 qt. per acre. Apply 1/2 to 1 1/2 qts. 21 days before harvest. California Only: Apply 1/2 to 1 1/2 qts. 45 days before harvest. Do not feed treated fresh hay, spent hay to dairy or meat animals.

Citrus (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangerines, Tangelos, Tangelos): Citrus Flat, Citrus Red, Citrus Rust, Six-Spotted, Texas Citrus, Yuma Spider Mite, Pacific Spider, Two-Spotted Spider Mite. Apply 1 1/2 to 4 qts. per acre. Do not apply within 7 days of harvest.

Cotton: Desert Spider, Pacific Spider, Two-Spotted Spider, Strawberry Spider Mite. Apply 1/2 to 1 1/2 qts. per acre. Do not apply within 14 days of harvest. Do not graze dairy or meat animals in treated fields or feed grain to them. Crabapples: Apple Rust Mite, Brown Mite, European Red Mite, McDaniels Spider Mite, Pacific Spider Mite, Schoene Spider Mite, Yellow Spider Mite, Two-Spotted Spider Mite, Willamette Mite. Apply 1 1/2 to 4 qts. per acre. Claims for Brown Mite and European Red Mite are for cleanup of late spring populations. Repeat

BEST AVAILABLE COPY

needed, but allow 30 days between treatments. As a preventive schedule apply 1/2 pt. per acre. Make two successive applications at 10 to 14 day intervals.
Cucumbers, Melons (including Watermelons), Pumpkins, Squash: Pacific Spider Mite, Two-Spotted Spider Mite. Apply 1/2 to 1/4 pt. per acre. Do not apply within 2 days of harvest.

Pilberts, Walnuts, Hickory Nuts, Peasans, Chestnuts: Brown Mite, European Red, Two-Spotted Spider, Pacific Spider Mites, Yellow Spider Mite. 1 1/2 to 4 qts. per acre. Do not apply within 14 days of harvest. Do not feed treated material to livestock.

Grapes: European Red, McDaniel Spider, Pacific Spider, Two-Spotted Spider, Williams Mite. Apply 1/2 to 1 qt. per acre. Do not apply within 7 days of harvest.

Hops: Pacific Spider, Two-Spotted Spider Mite. Apply 1/2 to 1 qt. per acre. Do not apply within 7 days of harvest.

Mint (peppermint and spearmint for oil production): Two-spotted Spider Mite. Apply 1/2 to 1 qt. per acre. Do not apply within 30 days of harvest. Do not make more than one application per crop season. Do not feed treated fresh hay or spent hay to livestock.

Pears: Apple Rust Mite, Brown Mite, European Red Mite, McDaniel Spider, Pacific Spider, Schoone Spider, Two-Spotted Spider, Williams, Yellow Spider Mite. Apply 1 1/2 to 4 qts. per acre. Claims for Brown Mite and European Red Mite are for clean-up of late spring populations. Repeat as needed but allow 30 days between treatments. Dical should not be used in late cover sprays as plant injury may result.

Peppers, Tomatoes: Pacific Spider, Tomato Rust, Tropical Spider, Two-Spotted Spider Mites. Apply 1/2 to 1/4 qt. per acre. Do not apply within 2 days of harvest.

Quince: Apple Rust, Brown, European Red, McDaniel Spider, Pacific Spider, Schoone Spider, Two-Spotted Spider, Williams, Yellow Spider Mite. Apply 1 1/2 to 4 qts. per acre. Do not apply within 7 days of harvest.

Strawberries: Strawberry Spider, Two-Spotted Spider Mite. Apply 1/2 to 2 qts. per acre. Cyclamen Mites. Apply 2 qts. per acre. Do not apply within 2 days of harvest. To control Cyclamen Mites, make 2 or 3 applications at 10 to 20 day intervals starting when new spring growth begins. Small areas of infestation may also be treated during the production season.

ORNAMENTALS


Lawns, Turf: Bermudagrass—Clover Mites. Apply 1 oz. in 2 1/2 gallons of spray per 1,000 sq ft. Repeat in 10 to 14 days if needed. For Clover Mite be sure to treat areas at base of buildings.

Flowering Plants (including African Violets, Chrysanthemums, Sweet Peas), Orchids, Shrubs, Trees, Vines, Nursery Stock: Mites. Apply 1 oz. in 2 1/2 gallons of water. Thoroughly cover upper and lower leaf surfaces. Wet to the dripping point.

DISCLAIMER OF WARRANTIES

Sellec warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS, EXPRESSED, IMPLIED OR STATUTORY INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond

the control of Sellec. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of this product or a refund of the purchase price of the product, at Buyer's option, as full discharge of Seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product and no such warranties, guarantees or directions shall be valid or binding upon Sellec.

ACCEPTED 
JUN 29 1983
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 34704-513

NOTE
THIS PRODUCT MUST BE STORED AT TEMPERATURES ABOVE 40°F.
STORAGE BELOW MAY AFFECT THE QUALITY OF THE PRODUCT.

INSECTICIDE, LIQUID N.O.S.
NA1993 RQ (KELTHANE)
(COMBUSTIBLE LIQUID)

BEST AVAILABLE COPY

3rd 12

20