

To kill roaches, waterbugs, silverfish, crickets, ants and certain other crawling insects: Spray baseboards, cracks, crevices, moist areas and openings around sinks and pipes and other hiding places, hitting the insects directly. Spray ant hills and ant trails over window and door sills and other locations.

To kill clothes moths, carpet beetles: Spray all articles to be protected giving particular attention to folds and seams. Spray interior of storage containers such as closets, chests and trunks regularly. Treatment should be repeated every thirty days unless the articles are stored in a mothtight container.

To kill bedbugs: Spray mattresses thoroughly, especially around seams and tufts. Spray bedstead, baseboards, floor boards. Repeat frequently.

TO KILL GARDEN PESTS (Outdoors)

For bugs on such plants as Asters, Azaleas, Begonias, Chrysanthemums, Dahlias, Delphiniums, Gladiolus, Irises, Marigolds, Pctunias, Roses and Snapdragons. Also for bugs on evergreen and small ornamental deciduous trees, shrubs and vines.

When sprayed directly on insects this spray kills sucking insects such as aphids, leafhoppers, spotted mites, thrips (exposed) and lace bug. Kills chewing insects such as Salt Marsh caterpillar, Mexican bean beetles, climbing cutworms and Japanese beetles.

For best results spray in evening or on a cloudy day when air is still. Use slow sweeping motion. Keep can 18 inches from plants. Avoid wetting foliage with the spray. Spray rows at the rate of 1 ft. per second. For heavy growth, spray row on both sides. Repeat when needed. Spray with caution on young plants and new growth. Do not spray garden when temperature exceeds 90°F.

NOTICE OF WARRANTY

Seller and Manufacturer make no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN ST. FREMONT, NE 68025



PEST CONTROL FOR HOUSE & GARDEN

For Indoor and Outdoor Use

Kills Ants, Flies, Moths
and Mosquitoes

ACTIVE INGREDIENTS:

Pyrethrins	0.250%
Technical Piperonyl Butoxide*	0.500%
N-Octyl Bicycloheptene Dicarb- oximide	0.500%
INERT INGREDIENTS**	99.750%
TOTAL	100.000%

*Consists of 0.400% (Butylcarbityl) (6-Propylpiperonyl) ether and 0.100% related compounds.

**This product contains
Aliphatic petroleum compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

Read cautions and directions on side panels.


EPA REG. NO. 34704-526

EPA EST. NO. _____

**NET WEIGHT 14 OUNCES
CONTENTS UNDER PRESSURE**

8201

EXP-1Y



CAUTION

Cover or remove fish bowls for 1 hour. Cover exposed food and utensils; this product can be safely used around foodstuffs and utensils if they are not protected from direct contact during spraying. Do not spray on plastics or asphalt tile.

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 120°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store upright in a cool place (not over 120°F). Do not place on hot surface. Do not use or store near heat or open flame. Exposure to temperature above 120°F may cause bursting. Store in original container only, preferably in a locked storage area.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Replace cap and discard container in trash. Do not incinerate or puncture.

Remove protective cap. Hold upright. Point spray opening away from you and press valve downward. Apply spray in short bursts.

TO KILL HOUSEHOLD INSECTS (Indoors)

To kill flies, mosquitoes, gnats, wasps, flying moths and certain other flying insects. Close all doors and windows and direct spray upward into center of room with a slow sweeping motion. Spray 5 to 10 seconds for average room. Keep nozzle 3 feet from surface. Keep room closed for 15 minutes after spraying.

**FOR CONTROL OF SODWORM (LAWN MOTH),
CUTWORMS, EARWIGS, ANTS, CHINCH BUGS,
GRASSHOPPERS, SOWBUGS AND WHITE GRUBS
INFESTING HOME LAWNS AND OTHER
ORNAMENTAL TURF AREAS.**

Contains Dursban®

ACTIVE INGREDIENTS:	BY WT.
Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate]	6.7%
INERT INGREDIENTS:*	93.3%
TOTAL	100.0%

Contains 1/2 lb. of chlorpyrifos per gallon

*This product contains aliphatic petroleum hydrocarbons.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Below For Additional Precautionary Statements

EPA REG. NO. 34704-523

EPA EST. NO. _____

NET CONTENTS _____

• DURSBBAN is a Registered T.M. of the Dow Chemical Company.
U.S. Patent No. 3,244,566

25335

12V69

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Do not smoke while using. Wash thoroughly after handling. Wash contaminated clothing before reuse. In case of contact, immediately flush exposed eyes or skin with plenty of water. Get medical attention if irritation persists. Avoid breathing spray mist or vapors. Keep children and pets off treated areas until spray has dried.

NOTE TO PHYSICIAN: Active ingredient is a Cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in original container only, preferably in locked areas. Keep container tightly closed when not in use. Store in a cool, dry place. Do not use food or drink containers for mixing or storage.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

Read entire label. Use strictly in accord with label cautions, warnings

and directions. This product contains Dursban® insecticide. This phosphate insecticide provides control of insects attacking home lawns and other ornamental and recreational turf grass areas. Insects controlled: Lawn Moth (Sod Webworm), Chinch Bugs, Cutworms, Earwigs, Grasshoppers, Armyworms (*Prodenia* spp.). Spray when pests first appear; repeat when needed. One application usually will control Chinch Bugs for up to 8 weeks. Mix this product with water according to the directions in the following tables. Apply as a coarse (large droplet) spray using a low pressure sprayer such as a hose-attached applicator, knapsack-type, tank-type, or similar equipment. For best results, the lawn should be moist at time of application and care should be taken to obtain complete coverage of pest infested areas.

DOSAGE AND DILUTIONS FOR USE ON TURF

Area To Be Treated	Amount To Use*	Dilute With Water To Give
500 sq. ft.	3 fl. oz.	15 gals.
1,000 sq. ft.	6 fl. oz.	30 gals.
10,700 sq. ft.	2 qts.	325 gals.
One Acre	2 gals.	1,300 gals.

*1 fluid ounce 2 tablespoonful 6 teaspoonful

TO CONTROL CERTAIN WHITE GRUBS (larvae of European chafer and Japanese Beetle).

Use two to four times the amounts of DURSBBAN INSECT SPRAY shown in the TURF table (i.e. 12-24 fl. oz. per 1000 sq. ft.). Apply in enough water to obtain uniform coverage of the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

HORTICULTURAL NOTE: Damaged lawns and ornamental turf should be fertilized to stimulate re-growth and recovery from insect injury.

PESTS: Out of Doors:

To help prevent infestation of buildings by ants, crickets, earwigs and sowbugs (pillbugs), treat a band of soil 2 to 3 ft. wide around and adjacent to the building, also the building foundation to a height of 2 to 3 ft., where pests are active and may find entrance. Use the same dosage and dilution recommended in the table for use on turf. Apply as a coarse spray to thoroughly and uniformly wet the band area.

PESTS OF ORNAMENTALS: To control bagworms, cutworms, grasshoppers, leafhoppers, mealybugs, mites, spittlebugs, thrips (exposed) and whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute with water according to directions given in the following table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 days intervals, if needed. NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants.

**DOSAGE AND DILUTIONS
FOR USE ON ORNAMENTAL PLANTS**

PESTS CONTROLLED	AMOUNT IN WATER TO MAKE		
Bagworms, Grasshoppers,	3 Gals.	9 Gals.	100 Gals.
Mites Spittlebugs and	2 fl. oz.	6 fl. oz.	2 qts.
Whiteflies			
Cutworms, Mealybugs, Leaf-	4 fl. oz.	12 fl. oz.	1 gal.
hoppers and Thrips (ex-			
posed)			

* Equivalent to 4 ozs. of chlorpyrifos per 100 gals.

** Equivalent to 8 ozs. of chlorpyrifos per 100 gals.

(continued on page 2)

