

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: If container is damaged. Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required as to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material such as clay, diatomaceous earth, sand, or sawdust. Sweep the contaminated absorbent on to a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Do not store, use, pour or spill near heat or open flame. Do not store in car trunk where temperatures may exceed 120° F.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or incineration or if allowed by State and local authorities, by burning. If burned, stay out of the smoke.

CONSUMER STORAGE AND DISPOSAL STATEMENTS

STORAGE: Store in original container only. Place in a locked storage area.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

Directions For Applying

Problem Area	Pests—Rate and Area Covered
Lawns	Ants, Armyworms, Bermudagrass Thrips, Brown Dog Ticks*, Chiggers, Crickets, Cutworms, Earwigs, Fleas, Frit Flies, Lawn Billbugs**, Chinch Bug†, Lawn Moths (Sod Webworms), Leafhoppers, Mites (Bermuda Clover), Sowbugs (Pillbugs), Springtails (Collembola)
Bent, Bermuda, Bluegrass, Dichondra, Fescue, Marion, St. Augustine	4 fl. oz. in 3 gals. of water covers 125 sq. ft. 32 fl. oz. in 24 gals. of water covers 1,000 sq. ft. Thoroughly wet down grass a few hours before applying. *Brown Dog Ticks: Do not apply to animals. In particular, spray under shrubbery and the grass next to house. **Lawn Billbugs: Water the lawn thoroughly immediately after treatment. †In the North, if Chinch Bugs are the only lawn insect problem, a lower rate of application can be used: 2 fl. oz. in 3 gals. of water covers 125 sq. ft. Milkweed: 8 fl. oz. in 3 gals. of water covers 125 sq. ft. Thoroughly wet grass a few hours before applying. Rodegrass Scales 4 fl. oz. in 3 gals. of water covers 100 sq. ft. Apply when Scale Crawlers first emerge. Ants, Digger Wasps* 4 fl. oz. in 3 gals. of water covers 125 sq. ft. Spot spray ant hills and wasp ground nest openings. *Make application at dusk when the wasps are not active. Billbugs*, White Grubs** of Japanese Beetle, European Chaffer, Southern Chaffer 4 fl. oz. in 3 gals. of water covers 125 sq. ft. Water grass thoroughly after application. *Billbug control: Apply when activity is first observed or when chewed or brown grass is present, indicating damage from the insect. **Grub control: Make application any time between late July and early October. Hyperodes Weevil (New York only) 3 fl. oz. in 3 gals. of water covers 125 sq. ft. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

Directions For Applying (Cont'd)

Problem Area	Pests—Rate and Area Covered
Trees & Shrubs	Wet leaves and branches to the dripping point. Try to penetrate dense foliage.
Arborvitae, Azalea, Birch, Boxwood, Camellia, Douglas Fir, Elm, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Plum, Poplar, Rhododendron, Spruce, Willow	Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dpterous Leaf Miners (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak), Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Pinet, Two-spotted Spider), Obecure Root Weevils, Omnivorous Leaf Tiers, Scale Crawlers (Cottony-cushion, Euonymus, Lecanum, Pine Needle, Pitt Scale, San Jose, Soft) 4 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray. Webworms, (Juniper, Mimosa*), Whiteflies 4 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray. *Mimosa Webworms: Apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals. Apple-and-Thorn Skeletonizers, Hemlock Chermes, Oak Looper, Oblique-banded Leaf Roller, Pear Slugs, Tent Caterpillars, Webworms (Cotoneaster, Fall) 12 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray.
Roses & Flowers	Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage.
Azaleas, Camellias, Chrysanthemums, Carnations, Gladiol, Lilac, Rhododendron, Roses	Aphids, Euonymus Scale Crawlers, Flea Beetles, Dpterous Leaf Miners, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies 4 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray. Do not use on certain ferns, hibiscus, poinsettia, papaya, pine, gardenia, and orchids since injury may occur.
Fruits* & Nuts	Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted. If infestations warrant, repeat applications at 7 to 10 day intervals.
Almonds, Filberts, Pecans, Walnuts	Mites 1 fl. oz. in 3 gals. of water. Apply as a full cover spray. Do not apply after husks open or split.
Apples, Pears	Aphids, Codling Moths, Eye-spotted Bud Moths, Fruit Tree Leaf Rollers, Leafhoppers, Tentiform Leaf Miners (pears), Maggots (apples), Mealybugs, Mites (European Red, Clover [brown], Pacific, Two-spotted Spider), Pearls (pears), Scale Crawlers (Cottony-cushion, San Jose, Forbes, Soft) 4 fl. oz. in 3 gals. of water. Apply as a full cover spray. On apples, slight russetting may occur on some varieties such as Golden Delicious.
Apricots, Peaches, Nectarines	Aphids, Apricot Mealybug (apricot), Mites (Clover [brown], Two-spotted), Scale Crawlers (Olive, San Jose) 4 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Cherries	Black Cherry Aphid, Cherry Fruit Fly, Cherry Fruitworm (Maggot), Fruit Tree Leaf Roller, Mites (Cherry Rust), Scale Crawlers (San Jose) 4 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Figs	Dried Fruit Beetle, Vinegar Flies (Drosophila spp.) 4 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Plums, Prunes	Aphids (Leaf Curl, Plum, Mealy Plum, Thistle), Mites (Clover [brown], European Red) 4 fl. oz. in 3 gals. of water. Apply as a full cover spray.
Caneberries (blackberries, boysenberries, dewberries, loganberries, raspberries)	Aphids, Leafhoppers, Raspberry Sawfly, Raspberry Fruitworm, Two-spotted Mites, Thrips 4 fl. oz. in 3 gals. of water. Apply as a full cover spray.

