

25% DIAZINON GARDEN INSECT SPRAY provides positive control of many hard-to-kill insects. It works quickly and remains effective for an extended period of time.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### WARNING

KEEP OUT OF REACH OF CHILDREN. May be fatal if swallowed. May be absorbed through the skin. Avoid contact with skin, eyes, and clothing. Do not breathe spray mist. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto sprayed grass until spray has completely dried.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. This product is highly toxic to bees exposed to direct treatment or to residues remaining on treated areas. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not get into any body of water. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated.

Do not use on ferns, poinsettia, hibiscus, papaya, pine gardenia, and orchids since injury may occur.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

KEEP CONTAINER TIGHTLY SEALED DURING STORAGE. Do not reuse empty container. Wrap container and put in trash collection.

**LAWNS:** Bluegrass, Bentgrass, Fescue, Bermuda, Dichondra, St. Augustine. To control Chinch Bugs, Sod Webworm (Lawn Moth), Ants, Bermuda Mites, Earwigs, Lawn Billbugs, Fleas, Leafhoppers, Clover Mites: Use 2 tbsps. (1 fl. oz.) per 3 gallons of water to treat 125 sq. ft. Use 4 fl. oz. per 15 gals. of water to treat 500 sq. ft. Water lawn thoroughly a few hours before application. For Lawn Billbugs, water thoroughly after treatment.

**ROSES & FLOWERS:** Chrysanthemums, Carnations, Gladiolus, Azalea, Camellia, Rhododendron, Lilac. To control Aphids, Cutworms, Spider Mites, Thrips, Whiteflies, Leafhoppers, Mealybugs: Application

GROW!

THIS PRODUCT MUST NOT BE USED  
ON GOLF COURSES OR SOD FARMS.

## 25% DIAZINON GARDEN INSECT SPRAY

Controls Aphids, Spider Mites, Chinch Bugs,  
and Other Common Garden Insect Pests.

#### ACTIVE INGREDIENTS:

Diazinon [0,0-Diethyl 0-(2-Isopropyl-4-methyl-4-pyrimidinyl) phosphorothioate]	25.0%
INERT INGREDIENTS*	75.0%

TOTAL 100.0%

\*This product contains a xylene-range aromatic solvent.

### KEEP OUT OF REACH OF CHILDREN WARNING

SEE BACK PANEL FOR ADDITIONAL WARNINGS

EPA REG. NO. 34704-531

EPA EST. NO. \_\_\_\_\_

NET CONTENTS 1 QUART (946 ml)



24728



EXP-1Y

**UNUSUAL TREES & SHRUBS:** Arborvitae, Fir, Pine, Juniper, Spruce, Birch, Elm, Hawthorne, Holly, Locust, Maple, Oak, Poplar, Willow. To control Aphids, Spider Mites, Scale Crawlers, Pine Shoot Moths, Webworms, Leafhoppers, Leaf Miners: Apply as a thorough cover spray. Wet leaves and branches to dripping point. Penetrate dense foliage. Use 2 tbsps. (1 fl. oz.) per 3 gals. of water.

**CITRUS FRUITS:** Grapefruit, Lemon, Orange. To control Aphids, Spider Mites, Citrus Thrips, Mealybugs, Leafhoppers, Scale Crawlers: Apply as a full coverage spray when insects are first observed. Repeat as required. Do not spray within 21 days of picking or harvesting. Use 2 tbsps. (1 fl. oz.) per 3 gals. of water.

**Apples (14), Pears (14).** To control Aphids, Scale Crawlers, Codling Moth: Use 2 tbsps. per 3 gals. of water. Start spraying at petal fall. Repeat at 10 to 14 day intervals as needed. Apples should be sprayed at 18 to 21 day intervals for additional sprays.

**Almonds, Pecans, Filberts, Walnuts.** To control Aphids, Scale Crawlers, Spider Mites, Thrips: Use 2 tbsps. per 3 gals. of water. Spray when insects are noticed. Do not apply after husks open or split.

**Apricots (14), Cherries (10), Peaches (21), Nectarines (10), Plums (10), Prunes (10).** To control Aphids, Scale Crawlers, Cherry Fruit Flies, Fruit Tree Leaf Rollers, Spider Mites, Thrips, Peach Twig Borers: Start spraying at petal fall. Repeat at 7 to 10 day intervals as needed for control. Use 2 tbsps. per 3 gals. of water.

**To control Aphids, Cabbageworms on Cabbage (7), Broccoli (5), Cauliflower (5):** Aphids, Diterous Leaf Miner, Colorado Potato Beetle on Tomatoes (1), Potatoes (35), Peppers (5): Aphids, Mexican Bean Beetle, Leaf Miners on Beans (7), Peas (1): Aphids, Leaf Hoppers, Mites on Cantaloupe (3), Cucumber (7), Squash (10): 2 tbsps. per 3 gals. of water covers approximately 1,000 sq. ft. of garden.

Thoroughly spray all plant parts as required. Use no closer than 7 to 10 day intervals.

\*Numbers between parentheses ( ) indicate days required between last application and harvest.

**NOTICE:** Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one is authorized to make any other warranty, guarantee, or direction concerning this product. Because the time, place, rate of application, and other conditions of use are beyond Seller's control, Seller's liability from the handling, storage and use of this product is limited to replacement of product or refund of purchase price.

FORMULATED FOR

PLATTE CHEMICAL CO.

150 SO. MAIN STREET

FREMONT, NEBRASKA 68025

Reviewed. Registrant claims to be in accordance with  
application of Pesticide Product Guidance for Phase 2  
Registration, Page 2.12, Addendum to Insert Change in Status.