

PROD 1668 4104-573 10



UROX-[®]'HX'

GRANULAR WEED KILLER

For General Weed Control on Non-Crop Lands

ACTIVE INGREDIENT:

Bromacil [5-Bromo-3-sec-butyl-6-methyluracil] 4.0%

INERT INGREDIENTS: 96.0%

TOTAL: 100.0%

EPA Reg. No. 2393-297

EPA Est. No. 33560-TN-01

® Registered Trademark of Hopkins Agricultural Chemical Co.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

MAY CAUSE IRRITATION OF EYES,
NOSE, THROAT AND SKIN.

Avoid breathing dust or vapor.

Do not get on skin, in eyes or on clothing.
In case of contact, immediately wash skin
with plenty of water; for eyes, wash with
water for at least 15 minutes and get
medical attention.

ACCEPTED

JUN 13 1979

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, this pesticide
is registered under
EPA Reg. No. 2393-297

NOTE: Do Not apply, allow to drift or drain or flush equipment onto or near croplands, lawns, trees or other desirable plants or to areas where their roots may extend. Do Not contaminate irrigation ditches or water used for domestic purposes. Do Not allow contact with fertilizers, insecticides, fungicides or seeds. Application equipment should be thoroughly cleaned with a detergent and water to prevent corrosion. Equipment which has been used to apply this product should not be used for applying insecticides, fungicides, fertilizers or other agricultural chemicals as plant injury may result.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on the label.

DESTROY CONTAINER WHEN EMPTY

Destroy container so that it may not be used again for any purpose and dispose of it at some location where no attempt to salvage will be made.

DIRECTIONS

For use on non-crop land such as Railroad, Pipeline, and Highway Rights-of-Way, Industrial Plants, Lumber Yards, Tank Farms, Storage Areas and Parking Lots for control of most annual and perennial grasses and broadleaved weeds. Effects are slow to appear and will not become apparent until the chemical has been carried into the root zone of the weeds by moisture. Control will last for an extended period. The effectiveness and length of time control will last depend on type of soil, amount of rain and other conditions.

HOW TO APPLY: Use a mechanical spreader or other device which will spread the required amount evenly over the area to be treated. It may be applied when top growth is dense, but better control will be obtained when weeds are short or when excess growth is mowed and removed before applying.

WHEN TO APPLY: Best results will be obtained if applied shortly before or at the time plant growth begins. Rain or thorough watering after application will cause the material to move to the root zone and thus speed up control. Do Not use when ground is frozen.

GENERAL WEED CONTROL

ANNUAL WEEDS AND GRASSES: For foxtail, crabgrass, cheatgrass, wild oats, brome grass, puncturevine, ryegrass, pigweed, ragweed, turkey mullein, and lambsquarters use 60 to 120 pounds per acre.

PERENNIAL WEEDS AND GRASSES: For quackgrass, bluegrass, smooth brome, Bahiagrass, redtop, purpletop, broomsedge, aster, plantain, dandelion, wild carrot, dog fennel, and goldenrod use 140 to 240 pounds per acre.

For Hard-to-Kill Weeds and Grasses such as Johnsongrass, Bermuda grass, nutgrass, Dallisgrass, Vasey grass, saltgrass, bouncingbet, dogbane, bracken fern, and horse tail use 300 to 600 pounds per acre.

Use the higher level rates given on soils high in organic matter.

BRUSH CONTROL

For control of undesirable woody plants on non-crop land areas such as Railroad Rights-of-Way, Storage Areas, Industrial Plant Sites.

Apply in spring or summer as a broadcast or basal (spot) treatment. **NOTE:** For effective brush control and prevention of damage to desirable vegetation: do not apply to brush standing in water; do not use on rights-of-way or other sites where marketable timber or other desirable trees or shrubs are immediately adjacent to the treated area.


Broadcast Treatment: Apply 140 to 240 pounds per acre to control oak, willow, sweet gum, and pine; apply 300 to 600 pounds per acre to control brush such as American elm, winged elm, hackberry, sumac, and cottonwood. Use the higher levels of the dosage ranges on adsorptive soils (those high in organic matter or carbon).

Basal (Spot) Treatment: Apply 50 pounds at the rate of 1.25 to 2.50 ounces per stem 2" to 4" in basal diameter. Treatment controls woody plants such as cottonwood, hackberry, maple, oak, poplar, red bud, sweet gum, wild cherry, willow, and winged elm.

NOTICE OF WARRANTY

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

NET WEIGHT



MANUFACTURED BY
Hopkins
agricultural chemical co.

50 POUNDS

45A