

## VAPONA<sup>®</sup> INSECTICIDE 2EC cont'd.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

**STORAGE:** Do not store below temperature of 0°F. Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment listed under DANGER when handling open containers.

**SPILLED MATERIAL:** Block or dike to prevent spreading of spill. Cover with absorbent materials such as lime, clay or sawdust. Scoop and sweep into a disposal container. Wash area with strong lye solutions, absorb and place into a disposal container.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### APPLICATIONS

This product is a specially prepared animal insecticide for use in the control of certain insects attacking livestock (face flies, house flies, horn flies, stable flies, mosquitoes and gnats).

### MIXING DIRECTIONS

1% Solution: Mix 5 1/4 fl. ozs. (10 3/4 tbs.) concentrate with 1 gal. of water (1 qt. to 6 1/4 gals. 0.5% solution: Mix 2 3/4 fl. ozs. (5 1/2 tbs.) concentrate with 1 gal. of water (1 qt. to 12 1/2 gallons.)

### RECOMMENDATIONS

**BEEF & DAIRY CATTLE:** Flies (House flies, Horn flies, Stable flies, Face flies) and Mosquitoes—Apply 1 to 2 ozs. of 1% solution per animal as a mist spray daily with a hand or automatic sprayer. Spray to thoroughly cover all parts of the animal, including the legs but do not wet the skin. Do not apply in excess of 2 fluid ounces per animal per day.

**FACE FLY:** For maximum face fly control, add 1 to 2 cups of sugar to each gallon of 1% solution.

**CATTLE FEED LOTS, STOCK YARDS, HOLDING PENS and CORRALS:** Flies, Mosquitoes and Gnats. Apply at the rate of 4 1/2 gallons 0.5% solution per acre with a mist blower or other suitable equipment as an over-all application. The mist spray should be directed over the entire area. Particular attention should be given to areas where flies congregate on fences, spillage areas around feed bunks and walls of buildings. Animals may be present during treatment. Avoid direct application to exposed feed and water. For baited solutions dissolve one pound of sugar per 5-10 gals. of water before adding the Vapona concentrate.

**ANIMALS BUILDINGS (Dairy Barns, Milk Sheds, Shelter Sheds and Horse Barns):** For control of flies, gnats and mosquitoes. Before applying reduce air movement as much as possible by closing doors, windows and other openings.

**SURFACE SPRAY:** Apply 1 qt. of 0.5% solution or 1 pint of 1% solution as a coarse wet spray per 1000 sq. ft. For milk sheds, stables, livestock barns, loafing sheds, pig-pens and outdoor areas, treat around doorways, feed storage rooms, alleyways and window sills. In Poultry Houses apply to manure, window sills, outside of penned enclosures and on floors of feed and storage rooms. Also controls roaches, fleas, ants, sawbugs and brown dog ticks. Repeat as necessary. Do not treat animals or humans.

**FOG OR MIST:** Use 1 qt. of 0.5% or 1 pint of 1% solution per 8000 cubic feet of space. Animals can be left in buildings provided they have not received a direct application within eight hours.

**CAUTION:** Do not contaminate water, feed or foodstuffs, milk or milking utensils.

**FLY MAGGOTS:** In maggot breeding areas on garbage dumps and manure piles. Dilute 1 to 2 pts. of 1% Vapona in equal parts of diesel fuel and apply 1 to 2 qts. of the 0.5% solution per 100 square feet.

### NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to the seller, and buyer assumes the risk of any such use.

FORMULATED FOR  
**PLATTE CHEMICAL CO.**

150 SO. MAIN STREET      FREMONT, NEBRASKA 68025

