

CLEAN CROP
BENOMYL 50% DF SYSTEMIC FUNGICIDE

- A systemic fungicide to be mixed with water for spray application.
- Controls listed diseases on TURF, ROSES, FLOWERS, ORNAMENTALS, BULBS, LISTED GARDEN VEGETABLES AND FRUITS
- Provides Curative (Eradicant) and Protective Action

ACTIVE INGREDIENT:

Benomyl [Methyl 1-(butylcarbamoyl)-
2-benzimidazolecarbamate]

50%

INERT INGREDIENTS:

50%

TOTAL 100%

EPA Reg. No. 34704-

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN (Pt. Size as appropriate
for front panel)

C A U T I O N

(Pt. Size as appropriate
for front panel)

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Remove contaminated clothing and immediately wash affected area with soap and water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

May irritate eyes, nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using.

Protective Measures: Always spray with your back to the wind. Wear long-sleeved shirt, long pants, and rubber gloves. Wash gloves thoroughly with soap and water before removing. Change clothes immediately after using this product and launder separately from other laundry items before reuse. Shower immediately after use.

Do not enter or allow reentry into treated areas until sprays have dried unless wearing the protective clothing specified under "Protective Measures".


ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish in adjacent areas. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from fire or sparks.

NET CONTENTS:



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in original labeled container in a cool, dry, locked place out of reach of children. Do not allow this product to become wet in storage. This may lead to certain chemical changes which will reduce its effectiveness. Keep container tightly closed when not in use.
DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

This product is recommended for the control of certain diseases in turf, flowers, ornamentals, certain vegetables, fruits and fruit trees. It is effective for the control of many fungus diseases, providing both curative (eradicator) action and protective action. This product is used with water for application as a spray. USE LEVEL MEASUREMENTS. Continuous agitation is required to keep the material in suspension. Apply as a spray (except as otherwise directed), using sufficient water to obtain thorough coverage of the plants. Under severe disease conditions use the higher rate and shorter interval specified for each crop; also, for tree crops, use the higher rate for large mature trees. Consult your State Agricultural Extension Specialist for guidance on your particular crop and disease control situation.

TURF

For DOLLAR SPOT (*Sclerotinia homocarpa*), LARGE BROWN PATCH (*Rhizoctonia solani*), and FUSARIUM PATCH (*Fusarium nivale*) apply at the rate of 2 oz. (12 tbsp.) in 2-5 gals. of water per 1000 sq. ft. For FUSARIUM BLIGHT (*Fusarium roseum* and *Fusarium tricinctum*) use 6 oz. (36 tbsp.) in 2-5 gals. of water per 1000 sq. ft. Addition of a wetting agent to the spray mixture enhances curative action and improves distribution of spray. Start treatments at earliest appearance of disease and continue at 10 to 14 day intervals for as long as needed. When conditions are unusually favorable for development of disease, shorten interval to 5 to 7 days. For Stripe Smut use 6 ozs. (36 tbsp.) in 2-5 gals. per 1,000 sq. ft. in October or early Spring before grass begins to grow. After application, water turf sufficiently to carry product to base of plants. Do not graze or feed clippings from treated area to livestock.

ROSES, FLOWERS & ORNAMENTALS (Garden & Greenhouses)

This product is recommended as a foliar spray for control of many foliar diseases, as a plant drench for control of certain stem, crown and root diseases of ornamental plants, and as bulb soak against *Fusarium* and *Penicillium* rot.

FOLIAR SPRAY ON ROSES, FLOWERS & ORNAMENTALS: Except as noted begin applications when diseases first appear. Use 1 tbsp. per 2 gals. of water for Powdery Mildew and Botrytis Gray Mold. Use 2 tbsp. per 2 gals. of water for Black Spot on Roses. The higher rate also controls Anthraxnose (begin applications at bud break); Cercospora, Entomosporium, Ramularia, and Septoria leafspots; Ascochyta and

Phomopsis blights; Didymellina leafspot of Iris; Corynespora leafspot of Ligustrum; Ovulinia blight of Azalea and Rhododendron (begin as flowers open); and Scab of Pyracantha and Flowering Crab. Repeat at 10-14 day intervals throughout growing season for Roses. Repeat as long as needed for flowers and ornamentals. Shorten interval during humid, rainy weather. Use sufficient water to thoroughly wet foliage. Addition of a suitable approved Wetting Agent to the spray mixture enhances curative action of the fungicide and improves distribution of the spray on hard-to-wet plants such as Roses.

PLANT DRENCH on herbaceous annuals, perennials & bedding plants and on Woody Ornamentals such as azaleas, rhododendrons, conifers, and poinsettias: Apply a suspension of 1 tbsp. per gal. of water as a drench or heavy spray to plants after transplanting into propagation beds or containers, utilizing porous growing medium. Use 1 to 2 pints per square foot of bed container surface for control of Stem, Crown and Root Rots caused by Rhizoctonia, Fusarium, Sclerotinia or Botrytis on herbaceous annuals, perennials and bedding plants. Use the same rate for the control of Cylindrocladium and Thielaviopsis rots on Woody Ornamentals such as Azaleas, Rhododendron, Conifers and Poinsettias. Repeat application at 2 to 4 week intervals during disease pressure periods.

As an optional treatment, plants or cuttings may be dipped for 10 to 15 minutes in a suspension of 1 tbsp. per gal. of water, removed and allowed to drain, prior to transplanting and subsequent drench treatments.

NOTE: This product does not control Pythium or Phytophthora. For control of these diseases, consult your local Agricultural Extension Service for recommendations.

BULBS (Daffodil, Easter Lily, Gladiolus, Iris, Tulip): Fusarium and Penicillium Rots - Use 2 tbsp. per gal. of water (1 2/3 lbs. per 100 gals.). Soak cleaned bulbs for 15 to 30 minutes in warm dip (80 to 85 degrees F.) preferably within 48 hours after digging. Dry bulbs after treatment. If bulbs are for forcing, treat after bulbs have been heat cured.

VEGETABLES

BEANS: White Mold (Sclerotinia), Gray Mold (Botrytis) - Apply 3 3/4 - 5 tbsp. in 5 gals. water to 1000 sq. ft. (1 1/2 - 2 pounds per acre), at 25% to 50% bloom; repeat at peak bloom. Do not apply to lima beans within 28 days of harvest. For other beans, do not apply within 14 days of harvest. Do not graze or feed treated bean vines or hay to livestock.

CELERY: Early Blight (Cercospora), Late Blight (Septoria) - Apply 1 7/8 to 3 3/4 tsp. in 5 gals. water to 1000 sq. ft. (4-8 ozs. per acre); begin when disease first appears and repeat at 7 to 10 day intervals; Do not apply within 7 days of harvest.

CUCUMBERS, MELONS, PUMPKINS, SUMMER and WINTER SQUASH: Target Spot (Cucumbers), Gummy Stem Blight, Powdery Mildew, Anthrachnose - Apply 1 7/8 to 3 3/4 tsp. in 5 gals. water per 1000 sq. ft. (4-8 ozs. per acre.). Begin applications when plants begin to run or when disease first appears and repeat at 7 to 14 day intervals as needed. For Target Spot, use 7 day intervals as needed.

MUSHROOMS: Verticillium Spot (Dry Bubble) - Use 1 tbsp. per gallon and apply to bed surface at the rate of 12 1/2 gals. per 1000 sq. ft. Apply immediately after casing and repeat at pinning; alternatively, if

disease has occurred, apply to beds after picking and repeat 10 days later. Do not apply within 2 days of harvest.

TOMATOES (Garden and Greenhouse): Apply 1/2 - 1 tbsp. per gal. of water to control Gray (Botrytis), Leaf (Cladosporium) and White (Sclerotinia) Molds; Cercospora Leaf Spot; Phoma Leaf Spot. Begin applications when disease first appears and repeat at 7-14 day interval as needed.

FRUITS

BLUEBERRIES: Apply 2 1/2 tbsp. in 5 gals. water per 1000 sq.ft. (1 lb. per acre.). Do not make more than 4 applications before harvest; do not apply within 21 days of harvest. Mummy Berry, Botrytis Blossom Blight - Apply at green tip and repeat at 7 to 10 day intervals through petal fall. Anthracoese Leafspot - Apply when disease first appears and make one additional application 14 days later. After harvest, make up to 4 applications to the bushes at 14 day intervals as needed.

CANEBERRIES-RASPBERRIES, BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, OSWBERRIES: Botrytis, Powdery Mildew, Penicillium Rots - Apply 1 7/8 tbsp. in 5 gals. water per 1000 sq. ft. (3/4 lb. per acre) at early bloom [5 to 10%] and at full bloom; make up to 3 additional applications at 14 day intervals as needed. Do not apply within 3 days of harvest.

GRAPES:Botrytis Bunch Rot - Apply 2 1/2 to 3 3/4 tbsp.in 5 gals. water to 1000 sq. ft. (1 - 1 1/2 lbs. per acre) at first bloom [no later than 5% bloom] and repeat 14 days later if severe disease conditions persist. Make an additional application 3 to 4 weeks before harvest or when sugar begins to build; repeat 14 days later if conditions favorable for disease persist. This product does not control bunch rots caused by other organisms such as Rhizopus spp., Alternaria spp., and Diplodia spp.; these rots occur most frequently in high temperature areas such as the San Joaquin and Sacramento Valleys of California. Powdery Mildew, Black Rot, Bitter Rot - East of Rockies - Apply 1 7/8 to 3 3/4 tbsp. in 5 gals. of water per 1000 sq. ft. (3/4 - 1 1/2 lbs. per acre) when foliage first develops and repeat at 14 to 21 day intervals, or as needed, until berries are full size.

NOTE: Do not apply within 7 days of harvest.

STRAWBERRIES: Gray Mold (Botrytis), Powdery Mildew, Leaf Scorch, Leaf Blight, Mycosphaerella Leafspot - Apply 2 1/2 tbsp. in 5 gals. of water per 1000 sq. ft. (1 lb. per acre) at 10% bloom and at full bloom; continue at 10 to 14 day intervals, using 1 1/4 tbsp. in 5 gals. of water per 1000 sq. ft. Anthracoese - Apply 2 1/2 tbsp. in 5 gals. of water per 1000 sq. ft. when plants are established and repeat at 7 day intervals.

Transplants: Botrytis Crown Rot, Mycosphaerella Leafspot - Use 1/2 tbsp. per gal. of water. Immerse plants to give thorough wetting; remove and allow to drain.

TREE FRUITS: Apply this product by using sufficient volume of spray to provide thorough coverage - until sprays begin to drip from leaves. Do not graze livestock in treated areas. Apply at the following rates:

APPLES: For applications through cover sprays, use in tank mixtures as detailed below.

Apply with mancozeb for Scab, Powdery Mildew, Sooty Blotch, Plyspect, Cedar Apple Rust, Quince Rust, Bitter Rot, Black Rot, Brown Rot. - Use 1/2 tsp. of this product + 2 tsp. of 80% mancozeb per gallon water: 1 3/4 tsp. spray oil may be added to mancozeb spray mixture. Apply at 1/2" green tip and repeat at 7 to 14 day intervals (or as needed) through the cover sprays. Do not apply within 30 days of harvest.

Apply with captan for Scab, Powdery Mildew, Sooty Blotch, Flyspeck, Black Rot. - Use 1/2 tsp. of this product + 2 tsp. of captan 50 WP per 1 gallon of water; Note spray injury may result if Captan is used with, immediately before, or closely following an oil spray. Apply at 1/2" green tip and repeat at 7 to 14 day intervals (or as needed) through the cover sprays.

Postharvest Fruit Rots (Botrytis spp., Penicillium spp., Gloeosporium spp.) - Make a single application of 1 tsp. per 1 gal. water anytime from 3 weeks before harvest up to day of harvest. For additional protection of fruit to be held in storage, thoroughly wet harvested fruit by dipping or spraying at 1 1/2 tsp. per 1 gal. water.

CITRUS: Scab - Apply 3/4 to 1 1/2 tbsp. per 1 gal. water (1 1/2 - 3 lbs. per acre). Under conditions of severe disease pressure, apply at pinhead stage (just prior to first flush) and repeat at 2/3 petal fall; otherwise, make a single application at 2/3 petal fall.

Greasy Spot - Make a single application of 3/4 to 1 1/2 tbsp. per 1 gal. water during the period mid-June to mid-July.

Fruit Decay (Green Mold, Blue Mold, Stem-end Rot) - Preharvest Spray - Make a single application of 1/2 to 1 tbsp. per 1 gal. of water (1 - 2 lbs. per acre) anytime from 3 weeks prior to harvest up to day of harvest. Postharvest - Apply as a dip, flood, or spray using 1 to 2 tbsp. per 1 gal. water; do not immerse fruit for more than 5 min. When citrus wax is used, this product may be incorporated into the wax spray. Use the higher rate on more susceptible fruits and when excessive inoculum levels are present. For control of sporulation (penicillium spp.), apply as a spray in citrus wax using 4 tbsp. in 1 gal. water.

PEARS: Scab, Powdery Mildew, Sooty Blotch, Flyspeck - Use 3/4 to 1 1/8 tsps. per gal. of water. Apply at 1/2" green tip and repeat at 7 to 14 day intervals (or as needed) through the cover sprays. If an application is missed during an infection period, use 1 1/8 tsp. per gal. of water and apply as soon as possible after the infection period in order to deactivate scab and to prevent further infection. Postharvest Fruit Rots (Botrytis spp., Penicillium spp., Gloeosporium spp.) - Make a single application of 1 1/8 tsp. per gal. of water anytime from three weeks before harvest up to day of harvest. For additional protection of fruit to be held in storage, thoroughly wet harvested fruit by dipping or spraying at 1 1/2 tsp. per gal. of water.

Overwintering Scab - Apply 1 1/2 tsp. per gal. after harvest but before leaf drop. Thorough wetting of foliage is necessary.

STONE FRUITS: PEACHES, NECTARINES, APRICOTS, CHERRIES, FRUNES and PLUMS for the control of Brown Rot Blossom Blight, Fruit Brown Rot, Peach Scab, Powdery Mildew, Cherry Leaf Spot and Post Harvest Fruit Rots. Brown Rot Blossom Blight, Fruit Brown Rot: Apply 1 tbsp. per gal. of water (in combination with dormant oil) as a delayed dormant treatment prior to bud break. Use 1 tbsp. per 2 gals. water at early bloom (popcorn, red bud or green tip) and at full bloom. Beginning three weeks before harvest, apply 1 or 2 additional sprays of 1 tbsp. per 2 gals. water.

Peach Scab: Use same schedule as for Brown Rot, plus applications of 1 tbsp. per 2 gal. water at shuck split and shuck fall: (two Sprays).

Powdery Mildew: Use same schedule as for Brown Rot, plus one application of 1 tbsp. per gal. water at shuck split and first cover spray. (two Sprays)

Cherry Leaf Spot: Use same schedule as for Brown Rot, plus a single application of 1 tbsp. per 2 gals. water 2 to 3 weeks after harvest.

Post Harvest Fruit Rots: Dip or spray fruit thoroughly as soon as possible after harvest using 1 tbsp. per 2 gals. of water. This treatment does not control fruit rots caused by Rhizop spp. or Alternaria.

RESISTANT FUNGUS STRAINS MAY BE PRESENT

IMPORTANT: If treatment is not effective following use of this product as recommended, a resistant strain of the fungus may be present. If treatment is ineffective due to the presence of a benomyl resistant strain, then neither this product nor any other benzimidazole or thiophanate type fungicide will effectively control that disease; consideration should be given to prompt use of other types of suitable fungicides.

NOTICE

Seller warrants that the product conforms to the chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
150 So. Main Street
Fremont, Nebraska 68025