



Reg # 34704-609 PM-23

Platte Chemical Co.

419 18th Street (80631-5852)

P.O. Box 667

Greeley, CO 80632-0667

Ph. 303-356-4400

Fax 303-356-4418



103

CERTIFICATION OF COMPLIANCE

I, being an authorized representative of Platte Chemical Co., certify that all containers of Salvo

, EPA Reg. No. 34704-609,

produced by:

October 23, 1993 (Products registered for use on sites other than residential or turf, including sod farms)

June 15, 1994 (Products registered for use only on residential or turf sites, excluding sod farms)

October 23, 1993 (Products registered for both types of uses of above)

will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing use products. I further certify that all containers of said product sold or distributed by Platte Chemical Co. by:


April 15, 1994 (Products registered for use on sites other than residential or turf, including sod farms)

January 1, 1995 (Products registered for use only on residential or turf sites, excluding sod farms)


April 15, 1994 (Products registered for both types of uses above)

will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing use products.

William M. Mahlburg 3/1/93
William M. Mahlburg Date
Registration Manager



SALVO



LOW VOLATILE WEED KILLER

ACTIVE INGREDIENT:

Isooctyl (2-ethylhexyl) ester of 2,4-Dichlorophenoxyacetic acid

81.8%*

INERT INGREDIENTS

18.2%

TOTAL

100.0%

*Equivalent to 54.2% or 5 lbs. per gallon of 2,4-Dichlorophenoxyacetic acid.
Isomer specific by AOAC Method 6.275, 13th Ed, 1980.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel For Additional Precautionary Statements.

EPA REG. NO. 34704-609

EPA EST. NO. _____

NET CONTENTS _____ GALLONS

34330

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION. Avoid contact with eyes, skin and clothing. Do not take internally. Do not contaminate feed or foodstuffs. When mixing, loading, or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical-resistant gloves, long-sleeved shirt, long pants, socks and shoes. For aerial applicators in enclosed cockpits and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required.

Wash hands, face and arms with soap and water as soon as possible after mixing, loading, or applying this product. Wash hands, face and arms with soap and water before eating, smoking, or drinking. Wash hands and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower.

Rinse gloves before removing.

For containers over 1 gallon and less than 5 gallons in capacity: Persons engaged in open pouring of this product must also wear coveralls or a chemical-resistant apron.

For containers of 5 gallons or more in capacity: A mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when cleaning of equipment washwaters.

Groundwater Contamination:

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Avoid spray drift to susceptible plants as this product may injure cotton, beans, peas, ornamentals, etc. Coarse sprays are less likely to drift. Do not contaminate domestic or irrigating water supplies. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not seed lawns, gardens or fields until at least four weeks after application of spray to area. Do not mix with liquid fertilizers. Do not apply in any manner not specified on the label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with this labeling.

BEST AVAILABLE COPY

AGRICULTURAL USE REQUIREMENTS

Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to statements on this label about personal protective equipment, restricted-entry interval and notification to workers. For any requirements specific to your State, consult the agency in your State responsible for pesticide regulation.

Do not enter or allow worker entry into treated areas during the restricted-entry interval of 12 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, wear eye protection, chemical resistant gloves, long-sleeved shirt, long pants, socks and shoes.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides, or fungicides. Do not store near heat or open flame. Containers should not be stacked more than six high. Reclose all partially used containers by thoroughly tightening screw cap. Damaged or leaking containers which cannot be used immediately should be transferred to suitable sound containers and properly marked. Absorb any spills with a suitable clay absorbent and dispose of as indicated under "Pesticide Disposal."

To prevent safety and prevention of unauthorized use, all pesticides should be stored in locked facilities.

To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification.

Opened, partially used pesticides should be stored in original container labeled containers when possible. When transfer to another container is necessary because of leakage or damage, carefully mark and identify contents of the new container.

If stored below freezing, it may be necessary to warm product to 40°F. and agitate before using. This does not affect the efficiency of the product.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Do not reuse empty container. Plastic: Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Metal: Triple rinse (or equivalent), adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

GENERAL INFORMATION

When filling spray tank first fill tank 1/4 full of water; add the calculated amount of SALVO, then finish filling tank with water with by-pass open and pump running.

Material may tend to settle when not agitated. When mixture is left in sprayer, allow by-pass pressure to re-agitate before resuming spraying.

PRECAUTION

Because of the difficulties of removing 2,4-D from sprayers do not use the same spray equipment for applying other materials to plants. Injury will result. Use a separate sprayer for this purpose.

Do not store near fertilizers, seeds, insecticides, fungicides or plants.

TIME OF APPLICATION

Annual weeds are most susceptible in the seedling stage and early growth stages of development and when conditions are such as to promote a vigorous growth.

WHEN USED IN CROP SPRAYING AS A SELECTIVE SPRAY, THE STAGE OF GROWTH OF THE CROP MUST ALSO BE CONSIDERED, OTHERWISE SEVERE DAMAGE MAY RESULT.

SOME SUSCEPTIBLE WEEDS

Arrowhead	Dock	Mexican Weed	Shepherd's Purse
Artichoke	Fansweed	Morning Glory	Smart Weed
Beggar Ticks	Figwort	Musk Thistle	Sow Thistle (annual)
Blindweed	Four O'Clock	Mustards	Stinkweed
Blue Thistle	Ground Ivy	Nettles	Sumack
Bull Thistle	Hemp Dogbane	Pennycress	Sunflower
Burdock	Herbit	Pepper Grass	Wild Garlic
Bullseye	Horsetail	Pig Weed	Wild Mustard
Canada Thistle	Jimsonweed	Prickly Lettuce	Wild Onion
Catnip	Kamashweed	Puncture Vine	Wild Parsnip
Chickweed	Lambquarters	Purslane	Wild Radish
Chicory	Loose Weed	Ragweeds	Wild Rape
Cocklebur	Mallow	Russet Thistle	Wild Sweet Potato
Dandelion	Marshall	Quickweed	Yellow Goatbeard

Some weeds above, listed as susceptible may not be completely killed but growth is so retarded by proper application that they are no longer a factor affecting growth and yield of crop.

APPLICATIONS

Application rates may vary slightly from State to State depending upon soil types and weather factors. Consult your State agricultural experiment station for recommendations under local conditions or more specific uses. Do not apply this product through any type of irrigation system.

CORN, Pre-emergence: Apply 1 1/2 to 2 1/4 pints of SALVO per acre. Spray uniformly over the soil surface after planting (preferably 3 to 5 days) but before emergence. Use in 5 to 20 gallons of water. Do not use on light sandy soils, or when moisture is inadequate for normal weed growth.

CORN, MILK; Post-emergence: Apply 1/2 to 3/4 pint SALVO per acre. Make application when crop is small and more resistant to SALVO but when weeds are succulent or in active growth stage. Use lower dosage only against succulent, non-resistant, annual broadleaf weeds. Use drop nozzles when crop is 10 inches tall or higher. Keep spray off leaves. Do not spray after tassels appear. Some injury to corn may occur.

WHEAT, OATS, BARLEY, RYE; Fall: Apply 1/2 to 3/4 pint of SALVO per acre. Make application in the spring when weeds are small and grain is in full tiller stage (4 to 8 inches high) and always before the boot stage. Do not apply during boot or dough stage. Do not forage or graze treated grain fields within two weeks after treatment. Do not feed treated straw to livestock.

WHEAT, OATS, BARLEY, RYE; Spring: Apply 1/2 to 3/4 pint of SALVO per acre. Apply after plants are fully tillered but before stems begin to joint. Do not apply during boot or dough stage. Do not forage or graze treated grain fields within two weeks after treatment. Do not feed treated straw to livestock.

PERENNIAL GRASS PASTURES, MEADOWS: Apply 1/2 pint of SALVO per acre. Apply in the early spring when sufficient weeds have emerged and are actively growing and not too mature. Injury may result to bent grass and to clovers. Usually clovers will recover from early spring applications. Do not apply when grass is in boot to milk stage. Do not apply to newly seeded areas or after heading begins.

Do not graze dairy animals on treated areas within 7 days after application. Do not graze meat animals on treated areas within 3 days before slaughter. Do not cut treated grass for hay within 30 days after application.

PASTURE AND MEADOW USE REQUIREMENTS

Do not enter treatment areas until spray has dried or dust has settled. For early entry to treated areas, wear eye protection, chemical resistant gloves, long sleeved shirt, long pants, socks and shoes.

LAWNS, ORNAMENTAL TURF GRASS, GOLF FAIRWAYS: For large areas apply 1 1/2 pints of SALVO per acre. For small areas 3 1/2 teaspoons in 3 to 5 gallons of water applied to 1000 square feet. Spray 3 to 5 days after mowing. Retreatment of perennial weed areas may be necessary. May cause injury to bent grass, carpet grass, dichondra, St. Augustine and to clovers. Protect ornamentals from spray drift.

TURF USE REQUIREMENTS

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried. The maximum number of broadcast applications per treatment site is 2 per year.

NOTE: For application to sod farms, follow AGRICULTURAL USE REQUIREMENTS on this label.

NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR PURPOSES STATED ON SUCH LABEL, ONLY WHEN USED IN ACCORDANCE WITH DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER. PLATTE MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN STREET FREMONT, NEBRASKA 68025-6807

BEST AVAILABLE COPY