



**COMPANY
NUMBER** _____

34704

**PRODUCT
SERIAL NO.** _____

609

**LABEL
APPROVAL
DATE** _____


3-14-96

SUPPLEMENTAL LABELING

(CONTAINS ADDITIONAL INFORMATION WHICH MAY NOT APPEAR ON PACKAGE LABEL.)



EPA REG. NO. 34704-609



FOR USE IN CROP RESIDUE MANAGEMENT SYSTEMS IN SOYBEANS (Preplant only)

GENERAL INFORMATION

SALVO is a phenoxy-type herbicide that provides postemergence control of many susceptible annual and perennial broadleaf weeds. SALVO may be applied prior to planting soybeans to provide foliar burndown control of susceptible annual and perennial broadleaf weeds and certain broadleaf cover crops such as those listed on this label. SALVO should only be applied preplant to soybeans in situations such as reduced tillage production systems, where emerged weeds are present. Apply only according to the application instructions given below. Do not use any tillage operations between application of SALVO and planting of soybeans.

MIXING INSTRUCTIONS

Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may increase the herbicidal effectiveness of SALVO on certain weeds and may be added to the spray tank. Read and follow all directions and precautions on this label and on the label of each product added to the spray mixture.

APPLICATION PROCEDURES

Apply using air or ground equipment in sufficient gallonage to obtain adequate coverage of weeds. Use 2 or more gallons of water per acre in aerial equipment and 10 or more gallons of spray mixture per acre for ground equipment.

APPLICATION TIMING AND USE RATES

Maximum Rate Per Acre	When to Apply (Days prior to planting soybeans)
1/5 pint (12.8 fl. ozs.)	NOT LESS THAN 7 DAYS
1 3/5 pints (25.6 fl. ozs.)	NOT LESS THAN 30 DAYS

WEEDS CONTROLLED

- | | | |
|------------------------------|----------------------|--------------------------|
| alfalfa* | horsweed or maretail | ragweed, common |
| bindweed* | ironweed | ragweed, giant |
| bullnettle | lambquarters, common | shepherdspurse |
| bittercross, smallflowered | lettuce, prickly | smartweed, Pennsylvania* |
| buttercup, smallflowered | morningglory, annual | sowthistle, annual |
| Carolina geranium | mousetail | speedwell |
| cinquefoil, common and rough | mustard, wild | thistle, Canada* |
| clover, red* | onion, wild* | thistle, bull |
| cocklebur, common | pennycress, field | velvetleaf |
| dandelion* | peppergrass* | vetch, hairy* |
| dock, curly* | plantains | Virginia copperleaf |
| eveningprimrose, cutleaf | purslane, common | |
| garlic, wild* | | |

* These species are only partially controlled.

In general, weeds should be small, actively growing and free of stress caused by extremes in climatic conditions, diseases, or insect damage at the time of treatment. The response of individual weeds species to SALVO is variable. Consult your local county or state Agricultural Extension Service or crop consultant for advice.

APPLICATION RESTRICTIONS AND PRECAUTIONS

Important Notice: Unacceptable injury to soybeans planted in fields previously treated with SALVO may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather (temperature and rainfall) from herbicide application until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present at the time of application. Do not apply SALVO as described on this label unless you are prepared to accept soybean injury, including loss of stand and yield.

Apply a maximum of one application per growing season regardless of the treatment rate.

Do not use on sandy soils with less than 1% organic matter.

Do not replant fields treated with SALVO in the same growing season with crops other than those labeled for use with SALVO.

Do not apply SALVO when weather conditions such as temperature air inversions or wind favor drift from treated areas to susceptible plants.

Livestock Grazing Restriction: Do not feed hay, forage or fodder. Restrict livestock from grazing treated fields. Livestock should be restricted from feeding/grazing of treated cover crops.

In fields previously treated with SALVO, plant soybean seed as deep as practical or at least 1 inch deep. Adjust the planter, if necessary, to ensure that planted seed is completely covered.

FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE PACKAGE LABEL.

FORMULATED FOR
PLATTE CHEMICAL CO.

150 SO. MAIN STREET FREMONT, NE 68025-5697

EXP02/6SHIRATE