

34704-609

10/23/2012

1/22



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Loveland Products Inc.  
Chris Mason, Ph.D.  
P.O. Box 1286  
Greeley, CO 80632-1286

OCT 23 2012

Subject: Label Amendment  
Product Name: Salvo  
EPA Registration Number: 34704-609  
Decision Number: 467919

Dear Dr. Mason,

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

A stamped copy of your label is enclosed for your records. This label supersedes all previously accepted labels. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after eighteen (18) months from the date of this letter or the next printing of the label, whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA §6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions, please contact Emily Hartman of my staff at (703) 347-0189 or [hartman.emily@epa.gov](mailto:hartman.emily@epa.gov).

Sincerely,

A handwritten signature in cursive script that reads "Kathryn V. Montague".

Kathryn Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

2/22



# SALVO®

## POSTEMERGENCE BROADLEAF HERBICIDE

SPECIAL LOW VOLATILE FORMULATION FOR CONTROL OF BROADLEAF WEEDS IN CERTAIN CROPS AND NONCROP AREAS

**ACTIVE INGREDIENT:**

Isooctyl (2-ethylhexyl) ester of 2,4-Dichlorophenoxyacetic acid .....	81.8%*
OTHER INGREDIENTS .....	18.2%**
<b>TOTAL</b>	<b>100.0%</b>

\*Equivalent to 54.2% or 5 lbs. per gallon of 2,4-Dichlorophenoxyacetic acid.  
Isomer specific by AOAC Method 6.275, 13th Ed, 1980.

\*\*Contains petroleum distillates.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

ACCEPTED

OCT 23 2012

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No 34704-609

### FIRST AID

<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

### EMERGENCY INFORMATION

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL 1-866-944-8565.**

**NOTE TO PHYSICIAN:** Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

**See Below For Additional Precautionary Statements.**

**EPA REG. NO. 34704-609**

**EPA EST. NO. 34704-MT-001**

**NET CONTENTS 2½ GALS. (9.46 L)**

**EXP 07/12 TYPO**

3/22

**SALVO®**  
**EPA REG. NO. 34704-609**

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

**CAUTION:** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**Personal Protective Equipment:**

Some materials that are chemical-resistant to this product are made of barrier laminate, nitrile rubber, neoprene rubber or viton. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

**All mixers, loaders, applicators, flaggers, and other handlers must wear:**

- Long-sleeved shirt and long pants,
- Shoes and socks,
- Chemical resistant gloves, except for pilots,
- Chemical-resistant apron for mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls:**

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

**Groundwater Contamination:**

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

**Nontarget plant precautions:**

This herbicide may cause injury to desirable plants by contacting foliage, stems or roots. Use care in all applications to avoid surface water or soil transport to nontarget plant areas. Avoid contamination of irrigation or domestic water supplies. Although this product is a low volatile formulation, at high temperatures (about 85° F or higher), vapors from this product may injure susceptible plants

4/22

**SALVO®**  
**EPA REG. NO. 34704-609**

growing nearby such as cotton, grapes, tobacco, fruit trees, legumes, vegetables, and ornamentals. Avoid applications in the vicinity of susceptible plants or when winds are blowing toward nearby susceptible plants, or when temperature inversions are expected. Avoid direct application or spray drift to susceptible plants since very small quantities of this herbicide can cause severe injury in the growing or dormant period. Plants contacted may be killed or suffer significant injury resulting in grade or yield losses. Do not apply in greenhouses.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls,
- Chemical-resistant gloves made of any water-proof material,
- Shoes plus socks.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people or pets to enter the treated areas until sprays have dried.

**SPRAY DRIFT MANAGEMENT**

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

**Droplet Size**

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

5/  
22

**SALVO®**  
**EPA REG. NO. 34704-609**

**Wind Speed**

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a medium spray, leave one swath unsprayed at the downwind edge of the treated field.

**Temperature Inversions**

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

**Susceptible Plants**

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

**Other State and Local Requirements**

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

**Equipment**

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

*Additional requirements for aerial applications:* The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

**Additional requirements for ground boom application:** Do not apply with a nozzle height greater than 4 feet above the crop canopy.

**General Precautions:** 2,4-D esters may volatilize during conditions of low humidity and high temperatures.

Do not apply during conditions of low humidity and high temperatures.

**GENERAL INFORMATION**

Best results will be obtained when SALVO® is applied during warm weather to young weeds that are actively growing under good moisture conditions. Lowest rates will generally be satisfactory on susceptible annual weed seedlings. For listed perennial or biennial weeds and under certain conditions such as drought or cool temperatures where control is difficult, the higher rates may be required. In general, only weeds emerged at the time of application will be affected.

When SALVO is used for weed control in actively growing crops, the growth stage of the crop must be considered. Proper timing is required to obtain maximum crop tolerance and to avoid crop injury. Weed control and crop tolerance of this product may be affected by local conditions, crop varieties, cultural practices, application methods and other factors. Users should consult Agricultural Extension Service, agricultural experiment station, university weed specialists, seed companies or other qualified crop advisors for information pertaining to local use. In general, weed control and crop tolerance will be best when plants have neither too little nor excessive moisture before or after application, and the crop is not under other stresses.

6/22

**SALVO®**  
**EPA REG. NO. 34704-609**

Certain states have regulations which may affect the use of this product. Contact your state pesticide authority for additional information.

Soil residue of this product may temporarily inhibit seed germination and plant growth.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. W.A.) For further information, please refer to EPA Website:

<http://www.epa.gov/espp/litstatus/eslitig.htm>

**MIXING INSTRUCTIONS**

SALVO is an emulsifiable concentrate formulation intended for dilution in water for many applications. For certain specified applications, liquid fertilizer or oil may replace part or all of the water as diluent.

If dry flowable (DF), wettable powder (WP) or flowable (F) tank mix products are to be used, these should generally be added to the spray tank before SALVO. Refer to mixing directions on tank mix product labels.

For best results, thoroughly clean sprayer immediately after use by flushing system with water and heavy duty detergent such as Loveland Products, Inc. Tank & Equipment Cleaner.

**Water Spray:** To prepare a water spray mixture, fill clean spray tank about 1/2 to 2/3 full with clean water. Add required amount of SALVO with agitation turned on. Continue agitation while adding balance of water and during spray operations. NOTE: This product forms an emulsion in water and can separate upon prolonged standing. If spray mixture is allowed to stand, agitate it before use to assure uniformity.

**Liquid Fertilizer Spray:** Due to increased risk of crop foliage burn with fertilizer, use only as recommended on this label or supplemental labeling distributed for SALVO. Use fertilizer rate recommended locally. Fill clean spray tank about 1/2 to 2/3 full with liquid nitrogen fertilizer (UAN or urea) solution. Add required amount of SALVO with vigorous agitation running. Continue agitation while adding balance of liquid fertilizer and during spray operations. Application should be made immediately. Overnight storage of mixture is not recommended. Application during very cold (near freezing) temperatures is not advisable because of the likelihood of crop injury. SALVO is formulated to be compatible with most liquid nitrogen solutions, however, due to variability in fertilizers, users may wish to perform a jar compatibility test before large scale mixing.

**Oil Spray:** Use only as recommended on this label or supplemental labeling distributed for SALVO. Fill clean spray tank about 1/2 to 2/3 full with diesel oil, fuel oil, stove oil or other suitable oil. Add required amount of SALVO with agitation turned on. Continue agitation while adding balance of oil. The resulting mixture is a solution and will generally remain uniform without agitation once mixed. However, agitation is suggested if available. Do not allow any water to get into the spray mixture to avoid formation of an invert emulsion (mayonnaise consistency).

**Water Spray With Oil:** Use only as recommended on this label or supplemental labeling distributed for SALVO. Where a combination of water and oil diluent is recommended, the use of emulsifiable crop oil or crop oil concentrate is suggested since mild agitation will be sufficient. Mix in the sequence of water, SALVO, and oil.

If diesel or other nonemulsified oils listed above under "Oil Spray" are desired for use with water, add no more than 1 quart of such oil per 1 gallon of water and agitate vigorously until tank is emptied. If possible, premix nonemulsified oil with SALVO and add this premix to a mostly filled spray tank with agitation on. Otherwise, mix in the sequence of water, SALVO, and oil with agitation on. Follow these procedures carefully to avoid formation of an invert emulsion (mayonnaise consistency).



**SALVO®**  
**EPA REG. NO. 34704-609**

**Weeds Controlled:**

Arrowhead  
 Artichoke  
 Blue thistle  
 Blueweed, Texas  
 Boxelder  
 Bittercress, smallflowered  
 Blue lettuce  
 Broomweed, common  
 Bull nettle  
 Burdock, common  
 Burhead  
 Buttercup, smallflowered  
 Carolina geranium  
 Carpetweed  
 Catnip  
 Chickweed  
 Chicory  
 Cinquefoil, common and rough  
 Cocklebur, common  
 Coffeeweed  
 Cornflower  
 Creeping jenny  
 Croton (Texas, woolly)  
 Dogfennel (mayweed)  
 Elderberry  
 Evening primrose, common  
 Evening primrose, cutleaf  
 Fanweed  
 Figwort  
 Four o'clock  
 Galinsoga (elderberry, hairy)

Goatsbeard  
 Healall  
 Horsetail  
 Ironweed  
 Jerusalem artichoke  
 Jewelweed  
 Jimsonweed  
 Klamathweed  
 Ladysthumb  
 Lambsquarters, common  
 Loco, bigbend  
 Mallow (Venice, dwarf, little)  
 Maretail  
 Marshelder  
 Mexican weed  
 Milk vetch  
 Morningglory (annual, common, ivy, woolly)  
 Mousetail  
 Mustards (except blue), prior to bolting  
 Pennycress (fanweed)  
 Pepperweeds (except perennial)  
 Plantains  
 Poison ivy  
 Poorjoe  
 Puncturevine  
 Purslane, common  
 Quickweed  
 Ragweeds (common, giant)

Redstem  
 Rough fleabane  
 Shepherdspurse  
 Sicklepod  
 Sneezeweed, bitter  
 Sowthistle (annual, spiny)  
 Spanishneedles  
 Speedwell  
 Stinkweed  
 Sumacs  
 Sunflower  
 Sweetclover (annual)  
 Tumbleweed  
 Velvetleaf  
 Vetches, except hairy  
 Virginia copperleaf  
 Wild hemp  
 Wild lettuce  
 Wild mustard  
 Wild parsnip  
 Wild radish  
 Wild rape  
 Wild sweet potato  
 Willow  
 Witchweed  
 Wormwood  
 Yellow goatsbeard  
 Yellow rocket  
 Yellow starthistle

**Weeds Partially Controlled (Higher rates and/or repeated applications may be needed):**

Alfalfa  
 Asters  
 Beggarticks  
 Bindweeds (hedge, European)  
 Buckbrush  
 Bull thistle  
 Canada thistle  
 Chamise  
 Clover, red  
 Corn gromwell  
 Coyotebrush  
 Dandelion  
 Docks  
 Dogbanes

Goldenrod  
 Ground ivy  
 Hawkweed  
 Henbit  
 Hoary cress  
 Knotweed  
 Manzanita  
 Musk thistle  
 Nettles  
 Peppergrass  
 Prickly lettuce  
 Rabbitbrush  
 Russian thistle  
 Sage, coastal

Sagebrush (big, sand)  
 Salsify (western, common)  
 Sand shinnery oak  
 Smartweed, annual  
 Smartweed, Pennsylvania  
 Tansyragwort  
 Vervains  
 Vetch, hairy  
 Western ironweed  
 Wild carrot  
 Wild garlic  
 Wild onion

**Weeds Partially Controlled And For Which Locally Resistant Biotypes May Occur:**

Pigweed



9/22

**SALVO®**  
**EPA REG. NO. 34704-609**

**Weeds Suppressed When Another Labeled Herbicide Is Also Applied:**

Bindweed (field)  
Russian knapweed

**TANK MIXES**

Unless otherwise prohibited on this label or the label of an intended tank mix product, SALVO may be applied in combination with any herbicide registered for the same crop, timing, and method of application. Observe the most restrictive label statements of various tank mix products used. **LIABILITY FOR CROP INJURY RESULTING FROM A TANK MIXTURE NOT SPECIFIED ON THIS LABEL, OR SUPPLEMENTAL LABELING DISTRIBUTED FOR SALVO, IS SPECIFICALLY DISCLAIMED BY LOVELAND PRODUCTS INC.**

**COMPATIBILITY**

Before full-scale mixing of this product with other herbicides, fertilizer solutions and adjuvants, it is advisable to determine the compatibility of the proposed mixture. Use proportionate quantities of each ingredient and mix in a small container. Always mix one product thoroughly with the diluent before adding another product. If no incompatibility is evident after 30 minutes, the mixture is generally compatible for spraying.

**PLANTING IN TREATED AREAS**

**Labeled Crops:** Within 29 days following an application of this product, plant only those crops named as use sites on this or other registered 2,4-D labels. Follow more specific limitations, if any, provided in the directions for individual crops. Labeled crops may be at risk for crop injury or loss when planted soon after application, especially in the first 14 days. Degradation factors described below should be considered in weighing this risk.

**Other Crops:** All other crops may be planted 30 or more days following an application without concern for illegal residues in the planted crop. However, under certain conditions, there may be a risk of injury to susceptible crops. Degradation factors described below should be considered in weighing this risk. Under normal conditions, any crop may be planted without risk of injury if at least 90 days of soil temperatures above freezing have elapsed since application.

**Degradation Factors:** When planting into treated areas, the risk of crop injury is less if lower rates of product were applied and conditions following application have included warm, moist soil conditions that favor rapid degradation of 2,4-D. Risk is greater if higher rates of product were applied and soil temperatures have been cold and/or soils have been excessively wet or dry in the days following application. Consult your local Agricultural Extension Service for information about susceptible crops and typical soil conditions in your area.

**APPLICATION INSTRUCTIONS**

Read all preceding general sections of label and NOTICE before use. Unless otherwise specified, applications may be made by ground or air equipment. Ground applications may provide more thorough coverage and better weed control. For selective postemergent weed control in crops, do not add oil, surfactant, fertilizer or other additives unless specifically recommended on this label or supplemental labeling distributed for SALVO.

**CORN (Field, Sweet and Pop):**

**Restrictions**

**Field and pop:**

Do not use treated crop as fodder for 7 days following application. The preharvest interval (PHI) is 7 days. Do not exceed a maximum of 3 lbs ae/acre per crop cycle.

Preplant or preemergence: Limited to one preplant or preemergence application per crop cycle. Maximum of 1.0 lb ae/acre per application.

10/22

**SALVO®**  
**EPA REG. NO. 34704-609**

Postemergence: Limited to one postemergence application per crop cycle. Maximum of 0.5 lb ae/acre per application.

Preharvest: Limited to one preharvest application per crop cycle. Maximum of 1.5 lbs ae/acre per application.

**Sweet Corn:**

Do not use treated crop as fodder for 7 days following application.

The preharvest interval (PHI) is 45 days.

Observe a minimum of 21 days between applications.

Do not exceed a maximum of 1.5 lbs ae/acre per crop cycle.

Preplant or preemergence: Limited to one preplant or preemergence application per crop cycle. Maximum of 1.0 lb ae/acre per application.

Postemergence: Limited to one postemergence application per crop cycle.

Maximum of 0.5 lb ae/acre per application.

SALVO may be applied to corn at several different timings. In all cases, plant corn to a uniform depth of at least 1 1/2 inches. Avoid applying this product with Accent® SP Herbicide because severe grass control antagonism may occur. SALVO should be applied at least 7 days before or 3 days after Accent SP Herbicide.

**Preplant:** To control existing broadleaf weed seedlings or burn down susceptible cover crops prior to planting, apply SALVO from 7 to 14 days before planting. To control grasses and certain other problem weeds, it may be desirable to use a tank mixture with other herbicides. Liquid fertilizers and agriculturally approved surfactants may be added. Observe the most restrictive label statements of various tank mix products used. Use SALVO rates according to the following table:

**CORN PREPLANT APPLICATION RATES**

Soil Texture	Organic Matter	Rate Per Acre
Fine or medium (silt and clay loams)	Less than 1%	Do not apply.
	1% or more	6.4 to 19.2 fluid ounces (.25 to .75 lb ae)
Coarse (sand, sandy loam, loamy sand)	Less than 2%	Do not apply.
	2% or more	6.4 to 12.8 fluid ounces (.25 to .5 lb ae)

**Preemergence:** To control small broadleaf weeds, apply SALVO after planting, but before corn emerges. Liquid fertilizers and agriculturally approved surfactants may be added. Do not apply SALVO preemergence if a preplant application of this product was made. Use SALVO rates according to the following table:

**CORN PREEMERGENCE APPLICATION RATES**

Soil Texture	Organic Matter	Rate Per Acre
Fine or medium (silt and clay loams)	Less than 1%	Do not apply.
	1% or more	6.4 to 16 fluid ounces (.25 to .625 lb ae)
Coarse* (sand, sandy loam, loamy sand)	Less than 2%	Do not apply.
	2% or more	6.4 fluid ounces (.25 lb ae)

\*Partial weed control may result on coarse soils due to lower rate.

11/22

**SALVO®**  
**EPA REG. NO. 34704-609**

**Postemergence:**

**Caution:** Do not apply with liquid fertilizer or oil. Many types of adjuvants will increase risk of crop injury. Where an adjuvant is required because of tank mixing with another herbicide, use the lowest recommended concentration of a nonionic surfactant (often 0.25% vol./vol. or less) to minimize such risk. Treated crop may be brittle and subject to breaking by wind and/or cultivation, especially in the 2 weeks following SALVO application.

**Early Postemergence:** To control small broadleaf weeds, apply SALVO broadcast from spike to 4-leaf stage of crop or up to 8 inches tall, whichever comes first. Avoid spraying just after corn leaves unfold. Postemergence application should not follow a preplant or preemergence application by less than 3 weeks. Use SALVO rates according to the table below.

**Late Postemergence:** Typical timing for this application is when most broadleaf weeds are no more than 4 to 6 inches tall and corn is between 8 and 16 inches tall. The timing can extend until corn is 36 inches tall or to tasseling, whichever occurs first, but weeds usually become too large and hard to control. Perennial weeds should be in the bud to bloom stage for best results. **Apply as a directed spray using drop nozzles to keep spray off crop foliage.** Do not apply from 7 to 10 days before tasseling to hard dough stage. Use SALVO rates according to the following table:

**CORN POSTEMERGENCE APPLICATION RATES**

<b>Crop Stage</b>	<b>Comments</b>	<b>Rate Per Acre*</b>
Spike to 4-leaf, or up to 8 inches tall	Early postemergence over-the-top broadcast spray. Ground or aerial application.	3.2 to 9.6 fluid ounces (.125 to .37 lb ae)
8 to 36 inches tall, before tasseling	Late postemergence directed spray using drop nozzles. Ground application only.	4.8 to 9.6 fluid ounces (.18 to .37 lb ae)

\*Lowest rates may not provide adequate weed control unless used in a tank mixture with another registered herbicide.

**Preharvest:** After the hard dough (or denting) stage when silks have turned brown, apply 12.8 to 25.6 fluid ounces of SALVO per acre (.5 to 1 lb ae/A) to suppress perennial weeds such as hemp dogbane or field bindweed, and many tall weeds such as cocklebur, pigweed and sunflower that interfere with harvest. Weed seed production will also be suppressed if SALVO application is prior to the flowering stage of weeds. For field and sweet corn grown for seed, apply after silks have turned brown. Use the labeled rate of 12.8 to 25.6 fluid ounces of Salvo per acre. The high rate is recommended under dry conditions. Do not forage or feed corn fodder for 7 days following application.

**NOTE:** Hybrid varieties may vary in tolerance to 2,4-D. Some varieties are easily injured. Your local seed company, or Agricultural Experiment Station or Extension Specialist may provide additional information.

**Postharvest:** Following the harvest of corn, certain perennial or biennial weeds produce new fall growth. To aid in suppressing these weeds before a hard freeze, SALVO may be applied at the rate of 12.8 to 25.6 fluid ounces per acre (.5 to 1 lb ae/A) either alone or in combination with other registered herbicides such as certain formulations of dicamba and picloram. See PLANTING IN TREATED AREAS section. Follow more restrictive limitations, if any, for tank mix products used.

**SALVO®**  
EPA REG. NO. 34704-609

**SORGHUM (Milo-Grain):**

**Restrictions:**

The preharvest interval (PHI) is 30 days. Do not permit meat or dairy animals to consume treated crop as fodder or forage for 30 days following application.

Postemergence: Limited to 1 application per crop cycle. Do not exceed a maximum of 0.5 lb ae/acre per application.

Postemergence: To control small broadleaf weeds, apply when sorghum is 6 to 15 inches tall to top of canopy. **If sorghum is taller than 8 inches to top of canopy, use drop nozzles to keep spray off crop foliage.** Do not treat during the boot, flowering or early dough stages. Do not forage or feed fodder for 7 days following application. Use SALVO rates according to the following table:

**SORGHUM (Milo) POSTEMERGENCE APPLICATION RATES**

Crop Stage	Comments	Rate Per Acre*
6 to 8 inches tall	Over-the-top broadcast spray. Ground or aerial application.	3.2 to 9.6 fluid ounces (.125 to .37 lb ae)
8 to 15 inches tall	Directed spray using drop nozzles. Ground application only.	4.8 to 9.6 fluid ounces (.18 to .37 lb ae)

\*Lowest rates may not provide adequate weed control unless used in a tank mixture with another registered herbicide. Highest rates may have increased risk of injury.

**SORGHUM-SUDAN GRASS HYBRIDS (Forage Crop Only):**

Postemergence: To control small broadleaf weeds, apply SALVO when sorghum-sudan has at least 6 leaves, is well established, and is 5 to 10 inches tall. Do not treat crop over 10 inches tall through maturity.

Plant Response: Even when SALVO is sprayed at the proper stage, some crop injury is likely, including reduced seed production. If risk of crop injury is unacceptable, do not use this product. The lower rate may reduce the risk of crop injury, but will result in reduced weed control.

Livestock Feeding Restrictions: Do not permit meat or dairy animals to consume treated crop as fodder or forage for 30 days following application.

**SORGHUM-SUDAN GRASS POSTEMERGENCE APPLICATION RATES**

Crop Stage	Rate Per Acre
At least 6 leaves, well established, 5 to 10 inches tall	6.4 to 12.8 fluid ounces (.25 to .5 lb ae)

**SMALL GRAINS (WHEAT, OATS, BARLEY, RYE) NOT UNDERSEEDED WITH A LEGUME:**

**Restrictions:**

The preharvest interval (PHI) is 14 days. Limited to 1.75 lbs ae/acre per crop cycle.

Postemergence: Limited to one postemergence application per crop cycle. Do not exceed a maximum of 1.25 lbs ae/acre per application.

Preharvest: Limited to one preharvest application per crop cycle. Do not exceed a maximum of 0.5 lbs ae/acre per application.

Apply SALVO to small grains as directed below.

Livestock Feeding Restrictions: Do not permit dairy animals or meat animals being finished for

**SALVO®**  
**EPA REG. NO. 34704-609**

slaughter to forage or graze treated grain fields within 7 days after treatment. Do not feed treated straw to livestock if an emergency and/or preharvest treatment is applied.

**Liquid Nitrogen Fertilizers:** At full tiller, SALVO may be combined with liquid nitrogen fertilizers suitable for foliar application to small grains. Refer to MIXING INSTRUCTIONS section of label for further information. Fertilizers can increase foliage contact burn of herbicides. Reducing the fertilizer rate and concentration will reduce the hazard of foliage burn.

**Spring Wheat and Barley:**

**Onset of Tillering Stage:** Grains are generally tolerant of these treatments, but risk of crop injury is greater than at full tillering stage. Do not make application if the risk of injury is unacceptable. Apply 6.4 to 9.6 fluid ounces of SALVO per acre (.25 to .37 lb ae/A) in the spring when grain has 1 or more tillers as well as 3 or more leaves. Do not apply from boot to dough stage. Apply 6.4 to 12.8 fluid ounces of SALVO per acre (.25 to .5 lb ae/A) when grain is in the full tiller stage (usually 4 to 8 inches tall). Do not apply from boot to dough stage.

**Emergency Weed Control:** Higher rates, up to 25.6 fluid ounces of SALVO per acre, (1 lb ae/A) may be needed to handle difficult weed problems in certain areas, such as under dry conditions especially in western areas. These higher rates increase the risk of crop injury. The severity of the weed problem should be balanced against the possibility of crop injury. Do not apply before the tiller stage nor from boot to dough stage.

**Winter Wheat, Barley and Rye:**

**Onset of Tillering Stage:** Grains are generally tolerant of these treatments, but risk of crop injury is greater than at full tillering stage. Do not make application if the risk of injury is unacceptable. Apply 6.4 to 12.8 fluid ounces of SALVO per acre (.25 to .5 lb ae/A) in the spring when grain has 1 or more tillers as well as 3 or more leaves. Do not apply from boot to dough stage.

**Full Tillering Stage:** For these applications, full tillering stage is defined as follows. Grain should have 3 or more tillers and the flag leaf should not be visible. Apply 6.4 to 12.8 fluid ounces of SALVO per acre (.25 to .5 lb ae/A) when grain is in the full tiller stage (usually 4 to 8 inches tall). Do not apply from boot to dough stage.

**Emergency Weed Control:** For improved control of difficult weeds and heavy weed infestations, apply up to 25.6 fluid ounces of SALVO per acre (1 lb ae/A). These higher rates increase the risk of crop injury. The severity of the weed problem should be balanced against the possibility of crop injury. Do not apply before the tiller stage nor from boot to dough stage.

**Spring Seeded Oats:**

**Full Tillering Stage:** For these applications, full tillering stage is defined as follows. Grain should have 3 or more tillers and the flag leaf should not be visible. Oats are less tolerant to SALVO than wheat or barley and present a greater risk of crop injury. The severity of the weed problem should be balanced against the possibility of crop injury. Larger weeds and hard-to-kill weeds may be poorly controlled, especially under dry conditions. Apply 6.4 fluid ounces of SALVO per acre (.25 lb ae/A) when grain is in the full tiller stage as specified above. Do not apply before the tiller stage nor from boot to dough stage.

**Fall Seeded Oats (Southern) Grown for Grain:**

Apply 6.4 to 12.8 fluid ounces of SALVO per acre (.25 to .5 lb ae/A) after full tillering, but prior to joints forming in the stem. Do not apply until after full tillering nor from jointing to dough stage. Oats are less tolerant to SALVO than wheat or barley and present a greater risk of crop injury. The severity of the weed problem should be balanced against the possibility of crop injury, especially at higher rates. Avoid spraying during or immediately following cold weather.

14/22

**SALVO®**  
**EPA REG. NO. 34704-609**

**Preharvest Treatment (Wheat, Oats, Barley, Rye):**

Apply 12.8 fluid ounces of SALVO per acre (.5 lb ae/A) when grains are in the hard dough stage to control large weeds that may interfere with harvest. In tank mixtures with other herbicides registered for preharvest application, a rate of 6.4 to 9.6 fluid ounces per acre (.25 to .37 lb ae/A) may be desired. Best results will be obtained when soil moisture is sufficient to cause succulent weed growth. Addition of a nonionic surfactant such as LI 700®, Activator 90, or similar product usually improves weed control.

**Postharvest (Wheat, Oats, Barley, Rye):**

Following harvest, a flush of new weed growth may occur. For control of many annual broadleaf species, apply SALVO at up to 12.8 fluid ounces per acre (.5 lb ae/A). Also, certain perennial or biennial weeds may produce new fall growth in stubble grain fields. To aid in suppressing these weeds, SALVO may be applied at the rate of 12.8 to 25.6 fluid ounces per acre (.5 to 1 lb ae/A) either alone or in combination with other registered herbicides such as dicamba or picloram. See PLANTING IN TREATED AREAS section. Follow more restrictive limitations, if any, for tank mix products used.

**FALLOW LAND AND CROP STUBBLE:**

**Restrictions:**

Plant only labeled crops within 29 days following application. Limited to 2 applications per year. Use a maximum of 2.0 lbs ae/acre per application. Wait a minimum of 30 days between applications.

Fallow land or land idle between crops may be subject to unwanted weed growth. For control of many annual broadleaf species, apply SALVO at the rate of 6.4 to 12.8 fluid ounces per acre (.25 to .5 lb ae/A). To aid in suppressing certain perennial or biennial broadleaf weeds, SALVO may be applied at the rate of 12.8 to 25.6 fluid ounces per acre (.5 to 1 lb ae/A). Use the high rate on older plants, drought stressed plants or for hard to kill species. See PLANTING IN TREATED AREAS section. Follow more restrictive limitations, if any, for tank mix products used. SALVO may be used to kill fall alfalfa stands in preparation for spring planting of row crops under conservation tillage. The treated alfalfa crop cannot be grazed, fed to livestock or cut for hay.

**SOYBEANS—PREPLANT ONLY—FOR USE IN CROP RESIDUE MANAGEMENT SYSTEMS:**

**Instructions:**

SALVO is a phenoxy-type herbicide that provides postemergence control of many susceptible annual and perennial broadleaf weeds. SALVO may be applied prior to planting soybeans to provide foliar burndown control of susceptible annual and perennial broadleaf weeds and certain broadleaf cover crops such as those listed on this label. SALVO should only be applied preplant to soybeans in situations, such as reduced tillage production systems, where emerged weeds are present. Apply only according to the application instructions given below. Do not use any tillage operations between application of SALVO and planting of soybeans.

**Restrictions:**

The maximum rate per crop cycle is 1.0 lb ae/acre.

Preplant: Limited to 2 preplant applications per crop cycle. Maximum of 0.5 lb ae/acre per preplant application. Apply not less than 7 days prior to planting soybeans.

or

Preplant: Limited to 1 application per crop cycle. Maximum of 1.0 ae/acre per preplant application. Apply not less than 15 days prior to planting soybeans.

**Mixing Instructions:**

Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may increase the herbicidal effectiveness of SALVO on certain weeds and may be added to the spray tank. Read and follow label directions and precautions on this label and on the label of each product added to the spray mixture.

1.5/22

**SALVO®**  
**EPA REG. NO. 34704-609**

**Application Procedures:**

Apply using air or ground equipment in sufficient gallonage to obtain adequate coverage of weeds. Use 2 or more gallons of water per acre in aerial equipment and 10 or more gallons of spray mixture per acre for ground equipment.

**Application Timing and Use Rates**

Maximum Rate Per Acre	When To Apply (Days prior to planting soybeans)
12.8 fl. ozs. (.5 lb ae)	Not less than 7 days
25.6 fl. ozs. (1 lb ae)	Not less than 15 days

**Weeds Controlled**

Alfalfa*	Horseweed or Marestalk	Ragweed, common
Bindweed*	Ironweed	Ragweed, giant
Bittercress, smallflowered	Lambsquarters, common	Shepherdspurse
Buttercup, smallflowered	Lettuce, prickly	Smartweed, Pennsylvania*
Carolina geranium	Morningglory, annual	Sowthistle, annual
Cinquefoil, common and rough	Mousetail	Speedwell
Clover, red*	Mustard, wild	Thistle, Canada
Cocklebur, common	Onion, wild	Thistle, bull
Dandelion*	Pennycress, field	Velvetleaf
Dock, curly*	Peppergrass*	Vetch, hairy*
Evening primrose, cutleaf	Plantains	Virginia copperleaf
Garlic, wild*	Purslane, common	

\*These species are only partially controlled.

In general, weeds should be small, actively growing and free of stress caused by extremes in climatic conditions, diseases, or insect damage at the time of treatment. The response of individual weeds species to SALVO is variable. Consult your local county or state Agricultural Extension Service or crop consultant for advice.

**Application Restrictions and Precautions:**

**Important Notice:** Unacceptable injury to soybeans planted in fields previously treated with SALVO may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather (temperature and rainfall) from herbicide application until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present at the time of application. Do not apply SALVO as described on this label unless you are prepared to accept soybean injury, including stand and yield.

Do not use on sandy soils with less than 1% organic matter.

Do not replant fields treated with SALVO in the same growing season with crops other than those labeled for use with SALVO.

Do not apply SALVO when weather conditions such as temperature air inversions or wind favor drift from treated areas to susceptible plants.

**Livestock Grazing Restriction:** Do not feed hay, forage or fodder. Restrict livestock from grazing treated fields. Livestock should be restricted from feeding/grazing of treated cover crops.

In fields previously treated with SALVO, plant soybean seed as deep as practical or at least 1 inch deep. Adjust the planter, if necessary, to ensure that planted seed is completely covered.

**SALVO®**  
**EPA REG. NO. 34704-609**

**GRASS PASTURES:**

**Restrictions:**

Postemergence: For susceptible annual and biennial broadleaf weeds: Use 1.0 lbs ae/acre per application. For moderately susceptible biennial and perennial broadleaf weeds: Use 1.0 to 2.0 lbs ae/acre per application.

For difficult to control weeds and woody plants: Use 2.0 lbs ae/acre per application.

Spot treatment: Use 2.0 lbs ae/acre.

Maximum of two applications per year. Maximum of 4.0 lbs ae/acre per year.

Wait a minimum of 30 days between applications.

If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.

To control many emerged broadleaf weeds, apply 6.4 to 19.2 fluid ounces of SALVO per acre (.25 to .75 lb ae/A) . Addition of a nonionic surfactant such as LI 700, Activator 90, or similar product usually improves weed control. Preferred timing is in the early spring when sufficient weeds have emerged, and when weeds are small and actively growing, but before weeds are too mature. Summer applications of SALVO to older, drought stressed weeds are less effective. However, weeds are more susceptible again in the fall when cooler, wetter conditions support active growth before a killing frost. For fall treatment of mature weeds or perennial weed regrowth, use up to 25.6 fluid ounces of SALVO per acre. Several seasons of spring plus fall treatments may be necessary to control certain perennials.

**Plant Response:** Injury may result to bentgrass, other warm season or southern grasses, and alfalfa, clover or other legumes. Do not use SALVO if this risk of injury is unacceptable. Clovers may recover from early spring applications. Do not apply when grass is in boot to milk stage, or after heading begins, if grass seed production is desired. Do not apply to newly seeded areas until grass is well established. Reseeding is not recommended for at least 30 days following SALVO application. Addition of a surfactant may increase the risk of injury to newly seeded grasses.

**Livestock Feeding Restrictions:** Do not graze dairy or meat animals on treated areas within 7 days after application. Do not cut treated grass for hay within 7 days after application.

**GRASS SEED CROPS:**

**Restrictions:** Limited to 2 applications per year. Maximum of 2.0 lbs ae/acre per application. Minimum of 21 days between applications.

To control many emerged broadleaf weeds, apply 6.4 to 19.2 fluid ounces of SALVO per acre (.25 to .75 lb ae/A) Use on established stands of cool season grass seed crops, such as bluegrass, tall fescue and perennial ryegrass. Make applications in the spring from the tiller to early boot stage. Do not spray in boot stage. New spring seedings may be treated after the grasses have more than 5 true leaves. On established stands that have had the seed crop removed, perennial weed regrowth may be treated in the fall at up to 25.6 fluid ounces of SALVO (1 lb ae) per acre. Refer to "Plant Response" and "Livestock Feeding Restrictions" under GRASS PASTURES.

**SOD FARMS:**

**Restrictions:** Limited to 2 applications per year. Maximum of 2.0 lbs ae/acre per application. Minimum of 21 days between applications.

**Instructions:** For best results, do not mow turf 1 to 2 days before or after application. Turf watering should be delayed until the day after application. Do not apply SALVO to newly seeded areas until grass is well established and has been mowed several times. A period of about 30 days after application is usually a sufficient interval before reseeding. Seeding a small area and observing response is recommended before large scale seeding.



**Cool Season Grasses:** To control many emerged broadleaf weeds in cool season turfgrasses such as tall fescue, bluegrass or perennial ryegrass, apply 6.4 to 19.2 fluid ounces of SALVO per acre (.25 to .75 lb ae/A) . Apply when weeds are small and are actively growing under good moisture conditions. Not for use on centipede, carpetgrass, St. Augustine, bentgrass or Dichondra turf, or where desirable clovers are present.

### **RANGELAND PASTURES AND PERENNIAL GRASSLANDS NOT IN AGRICULTURAL PRODUCTION:**

---

#### **Restrictions:**

**Postemergence:** For susceptible annual and biennial broadleaf weeds: Use 1.0 lb ae/acre per application. For moderately susceptible biennial and perennial broadleaf weeds: Use 1.0 to 2.0 lbs ae/acre per application.

For difficult to control weeds and woody plants: Use 2.0 lbs ae/acre per application.

**Spot treatment:** Use 2.0 lbs ae/acre.

Maximum of two applications per year. Maximum of 4.0 lbs ae/acre per year.

Wait a minimum of 30 days between applications.

If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.

---

**Livestock Feeding Restrictions:** Do not graze dairy or meat animals on treated areas within 7 days after application. Do not cut treated grass for hay within 30 days after application. For government program grasslands, follow program grazing restrictions if more restrictive than those given above.

**Instructions:** SALVO can be used to control or suppress a number of susceptible broadleaf weeds in rangeland, or perennial grasslands that are set aside from agricultural use such as in the Conservation Reserve Program (CRP) or similar government programs. Consult program rules to determine whether grass and hay may be used. For best results, apply when broadleaf weeds are small. Adequate moisture is needed for best grass tolerance and weed control. Addition of a non-ionic surfactant such as LI 700, Activator 90, or similar product usually improves weed control.

**Plant Response:** Injury to legumes, bentgrass, and other warm season grasses is likely to occur. Grasses may be discolored following treatment. Do not apply when grass is in boot to milk stage, or after heading begins, if grass seed production is desired.

**New Stands:** Preseeding applications should occur at least 30 days prior to seeding. Newly seeded stands should only be treated after they are well established (more than 5 true leaves) or injury may occur. Apply 6.4 to 12.8 fluid ounces of SALVO per acre (.25 to .5 lb ae/A) when weeds are small and actively growing. Addition of a surfactant may increase the risk of injury to new stands.

**Established Stands:** For best results, weeds must be actively growing. Apply 12.8 to 19.2 fluid ounces of SALVO per acre (.5 to .75 lb ae/A) for annual weeds and up to 25.6 fluid ounces per acre (1 lb ae/A) for biennial or perennial weeds. Treat biennial weeds when they are in the seedling to rosette stage and before flower stalks become apparent. Treat perennial weeds in the bud to bloom stage. For brush species in rangeland, apply up to 51.2 fluid ounces of SALVO per acre (2 lb ae/A) in an oil spray (see MIXING INSTRUCTIONS). Another option is to add 1 gallon of oil per acre to a SALVO water spray (see MIXING INSTRUCTIONS). Repeat applications in the same or subsequent year may be needed to control brush species.

SALVO®  
EPA REG. NO. 34704-609

**FOREST MANAGEMENT:**

**Restrictions:**

Broadcast application: Limited to 1 broadcast application per year.

Maximum of 4.0 lbs ae/acre per broadcast application.

Basal spray, Cut Surface - Stumps, and Frill: Limit of one basal spray or cut surface application per year. Maximum of 8.0 lbs ae per 100 gallons of spray solution.

Injection: Limit to one injection application per year. Maximum of 2 ml of 4.0 lbs ae formulation per injection site.

**Forest Site Preparation:**

**Budbreak Spray:** For control of alder, susceptible broadleaf weeds, and susceptible woody plants before planting forest seedlings, apply up to 96 fluid ounces of SALVO per acre (3.7 lb ae/A) in a minimum of 10 gallons spray mixture per acre. Apply as an oil spray (see MIXING DIRECTIONS) after alder buds break, but before foliage is 1/4 full size. A water spray including 2 to 4 quarts per acre of diesel oil, fuel oil, stove oil or crop oil concentrate may also be used.

**Foliage Spray:** To control alder and susceptible woody plants before planting forest seedlings, apply up to 96 fluid ounces (3.75 lb ae) of SALVO per acre in a minimum of 10 gallons spray mixture per acre. Apply as a water spray including, if desired, up to 1 quart of diesel oil, fuel oil, stove oil or crop oil concentrate per gallon of water (see MIXING INSTRUCTIONS). For best results, apply after alder foliage has reached full size.

**Conifer Release:**

To control alder, susceptible broadleaf weeds, and susceptible woody plants in young conifer stands, apply up to 51.2 fluid ounces (2 lb ae) of SALVO per acre in a minimum of 10 gallons spray mixture per acre. This spring foliage treatment should be applied as a water spray when 3/4 of the brush foliage has full size leaves and before new conifer growth reaches 2 inches in length. Such stages usually occur between early May and mid-June, but application timing should be based on growth stages of brush and conifers. Application may cause leader deformation or other conifer injury, but trees should overcome it during the next growing season.

To control tan oak, madrone, ceanothus, canyon live oak, and manzanita, and to release Douglas fir, hemlock, Sitka spruce or grand fir, apply up to 76.8 fluid ounces of SALVO per acre (3 lb ae/A) in a minimum of 10 gallons spray mixture per acre. This spring foliage treatment should be applied as a water spray including, if desired, up to 1 quart of diesel oil, fuel oil, stove oil or crop oil concentrate per gallon of water (see MIXING INSTRUCTIONS). Make application before new growth on Douglas fir is 2 inches long. To release ponderosa pine from the same species, treat before new pine growth begins in the spring. Addition of oil or oil concentrate may cause unacceptable injury to pines.

For dormant applications in late winter or early spring for control of susceptible woody species such as alder, willow, poplars, cherry, vine maple, ceanothus, tan oak, madrone, and manzanita, apply up to 76.8 fluid ounces (3 lb ae) of SALVO per acre in a minimum of 10 gallons spray mixture per acre. This dormant treatment should be applied in diesel oil, fuel oil, stove oil or other suitable diluent such as water plus crop oil concentrate (see MIXING INSTRUCTIONS). Do not use in plantations where pine and larch are among the desired crop species.

To control hazel brush in the Lake states, apply up to 51.2 fluid ounces of SALVO (2 lb ae) per acre in a minimum of 10 gallons spray mixture per acre. Apply as a water spray when new shoot growth of hazel is complete (usually mid-July).

After conifer species such as white pine, ponderosa pine, jack pine, red pine, black spruce, white spruce, red spruce, and balsam fir cease growth and harden off and brush is still actively growing in late summer, apply up to 74 fluid ounces of SALVO (2.9 lbs ae) per acre in a minimum of 10 gallons spray mixture per acre. Apply as a water spray to control certain competing hardwoods such

19/22

**SALVO®**  
**EPA REG. NO. 34704-609**

as alder, aspen, birch, hazel and willow. Since this treatment may cause conifer injury, do not use if possible injury cannot be tolerated.

**Forest Roadsides:**

To control susceptible broadleaf weeds and woody plants on forest roadsides, apply 25.6 to 76.8 fluid ounces of SALVO per acre (1 to 3 lb ae/A) in a minimum of 10 gallons spray mixture per acre. Apply as a water spray including, if desired, up to 3 quarts per acre of diesel oil, fuel oil, stove oil or crop oil concentrate (see MIXING INSTRUCTIONS). Apply when sufficient foliage is present for absorption of herbicide.

**Established Conifers (Including Christmas Trees):**

***Directed Spray or Spot Spray:***

To control susceptible broadleaf weeds, mix up to 51.2 fluid ounces (2 lb ae) of SALVO per 100 gallons of water and apply to emerged weeds in the spring with ground equipment. Avoid contacting conifer foliage with spray or drift as injury may result. For brush, mix 96 fluid ounces (3.75 lb ae) of SALVO per 100 gallons of water. Thoroughly spray brush in full foliage, but avoid contacting conifer foliage with spray or drift. Do not apply more than the equivalent of 96 fluid ounces of SALVO per acre.

***Over-the-Top Broadcast Application:***

To control susceptible broadleaf weeds, apply 25.6 fluid ounces of SALVO per acre (1 lb ae) in a minimum of 10 gallons spray mixture per acre. To decrease the potential for injury to firs, apply only before budbreak in the spring and/or after complete bud set and hardening in the late summer or fall. Avoid treatment during the year of intended harvest.

**ROADSIDES; MEDIANS; HIGHWAY, RAILROAD, UTILITY, AND PIPELINE RIGHTS-OF-WAY; VACANT LOTS; AROUND UTILITY INSTALLATIONS, TRANSFORMERS, PUMP HOUSES, AND BUILDINGS; STORAGE AREAS; FENCES; GUARDRAILS; LUMBER YARDS; INDUSTRIAL SITES; AIRPORTS; TANK FARMS; FARMSTEADS; AND SIMILAR NONCROP AREAS:**

**Restrictions:**

Postemergence (annual and perennial weeds): Limited to 2 applications per year. Maximum of 2.0 lbs ae/acre per application. Minimum of 30 days between applications.

Postemergence (woody plants): Limited to 1 application per year. Maximum of 4.0 lbs ae/acre per year.

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

For control of many broadleaf weeds and small woody plants, apply 12.8 to 51.2 fluid ounces of SALVO per acre (.5 to 2 lb ae/A). Use the high rate for woody plants. Applications may be as broadcast sprays, small area sprays or spot treatments. For small areas or spot spraying, use 3.2 fluid ounces of SALVO per gallon of water and spray weeds to runoff. Regardless of the method of application, use adequate spray volume for full coverage of weeds. Preferred application timing is in the early spring when sufficient weeds have emerged, and when weeds are small and actively growing, but before weeds are too mature. Summer applications of SALVO to older, drought stressed weeds are less effective. However, weeds are more susceptible again in the fall when cooler, wetter conditions support active growth before a killing frost. For fall treatment of mature weeds or perennial weed regrowth, use up to 25.6 fluid ounces of SALVO per acre (1 lb ae/A). Several seasons of spring plus fall treatments may be necessary to control certain perennials. Use of oil sprays or the addition of spray adjuvants improves weed control, but also increases risk of damage to desirable ground covers.

20/22

**SALVO®**  
**EPA REG. NO. 34704-609**

**Plant Response:** Bentgrass, other warm season or southern grasses, and alfalfa, clover or other legumes may be killed or injured. Do not apply when grass is in boot to milk stage, or after heading begins, if grass seed production is desired. Do not apply to newly seeded areas until grass is well established. Reseeding is not recommended for at least 30 days following SALVO application.

**ORNAMENTAL AND RECREATIONAL TURFGRASSES, LAWNS, GOLF COURSES (Fairways, Aprons, Tees and Roughs), PARKS, CEMETERIES:**

**Restrictions:** Postemergence: Limited to 2 applications per year. Maximum of 1.5 lbs ae/acre per application. The maximum seasonal rate is 3.0 lbs ae/acre, excluding spot treatments.

**Instructions:** Refer to TURF USE REQUIREMENTS in the NON-AGRICULTURAL USE REQUIREMENTS section of this label. The maximum number of broadcast applications per treatment site is 2 per year. For best results, do not mow turf 1 to 2 days before or after application. Turf watering should be delayed for at least 1 hour after application. Avoid contacting desirable trees, shrubs, flowers, or vegetables as plant injury may result. Do not apply to newly seeded areas until grass is well established and has been mowed several times. A period of about 30 days after application is usually a sufficient interval before reseeding grasses (or other plants). Seeding a small area and observing response is recommended before large scale seeding.

**Cool Season Grasses:** To control many emerged broadleaf weeds in cool season turfgrasses such as tall fescue, bluegrass or perennial ryegrass, apply 12.8 to 19.2 fluid ounces of SALVO per acre (.5 to .75 lb ae/A) (0.3 to 0.44 fluid ounces per 1000 square feet). Preferred application timing for broadcast treatment is in the early spring when small weeds have emerged and are actively growing under good moisture conditions. For very weedy turf, a followup broadcast or spot application may be warranted about 2 to 4 weeks later. Summer applications of SALVO are typically spot treatments of individual weeds that have emerged after a spring broadcast treatment. In the fall when cooler, wetter conditions again favor active weed growth, broadcast application may be appropriate for very weedy turf, such as an area that had no spring broadcast treatment. Not for use on centipede, carpetgrass, St. Augustine, bentgrass or Dichondra turf, or where desirable clovers are present.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Open dumping is prohibited. Avoid contamination of fertilizers, seeds, plants, insecticides, and fungicides in storage. It is preferable to store all pesticides in a locked area. Containers with screw caps should be closed tightly when not in use. When transfer to another container is necessary because of leakage or damage, carefully mark and identify contents of new container. If label is damaged or missing, contact dealer or manufacturer. Absorb spills with granular clay absorbent and dispose of as indicated under PESTICIDE DISPOSAL. If this product is stored below freezing, it is suggested that it be allowed to warm to at least 40° F and be agitated before use.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:**

**Nonrefillable container:** Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer

**SALVO®**  
**EPA REG. NO. 34704-609**

**Storage & Disposal cont'd.:**

or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at [www.acrecycle.org](http://www.acrecycle.org). Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**For packages up to 5 gallons. Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 56 gallons: Triple rinse or pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container about ¼ full with water, rinsing down all sides inside the container thoroughly. Recirculate water with the pump for 2 minutes. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**For refillable containers:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**BEFORE BUYING OR USING THIS PRODUCT**, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.


**SALVO®**  
**EPA REG. NO. 34704-609**

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Accent® is a registered trademark of E.I. Du Pont de Nemours & Co., Inc.  
LI 700® and Salvo® are registered trademarks of Loveland Products, Inc.



**FORMULATED FOR**  
**LOVELAND PRODUCTS, INC.**  
**P.O. BOX 1286, GREELEY, COLORADO 80632-1286**