



Reg # 34704-644

PM-23

P 194
%

Platte Chemical Co.

419 18th Street (80631-5852)

P.O. Box 667

Greeley, CO 80632-0667

Ph. 303-356-4400

Fax 303-356-4418

CERTIFICATION OF COMPLIANCE

I, being an authorized representative of Platte Chemical Co.,
certify that all containers of Clean Crop Weed & Feed 2,4-D
Granular, EPA Reg. No. 34704-644,

produced by:

October 23, 1993 (Products registered for use on sites other
than residential or turf, including sod farms)

June 15, 1994 (Products registered for use only on
residential or turf sites, excluding sod farms)

October 23, 1993 (Products registered for both types of uses
of above)

will bear revised labeling in accordance with the revised
labeling required for Task Force technical and manufacturing
use products. I further certify that all containers of said
product sold or distributed by Platte Chemical Co. by:


April 15, 1994 (Products registered for use on sites other
than residential or turf, including sod farms)

January 1, 1995 (Products registered for use only on
residential or turf sites, excluding sod farms)

April 15, 1994 (Products registered for both types of uses
above)

will bear revised labeling in accordance with the revised
labeling required for Task Force technical and manufacturing
use products.

William M. Mahlburg 3/1/93
William M. Mahlburg Date
Registration Manager



234



WEED & FEED 2,4-D GRANULAR

Kills Weeds & Feeds Lawns

Kills Dandelions, Chickweeds, Plantains, Buckhorn and similar susceptible Broadleaf Weeds.

For Cool Season Grasses Such as Bluegrass, Ryegrass, and Fescue. For Warm Season Grasses such as St. Augustine, Bahia, Bermuda and Zoysia.

ACTIVE INGREDIENT:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid.....	1.2%*
INERT INGREDIENTS:	98.8%
TOTAL	100.0%

*Equivalent to 1.0% 2,4-Dichlorophenoxyacetic Acid by isomer specific analysis.

Refer to tag for Fertilizer Labeling.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Below For Additional Precautionary Statements.

EPA REG. NO. 34704-644 EPA EST. NO. _____

NET WEIGHT 18 POUNDS

(Covers 5000 Sq. Ft. Most Grasses)

(Covers 10,000 Sq. Ft. Certain Warm Season Grasses—Treat Twice)


13989

EXP02P93

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Do not breathe dust. Avoid contact with skin or clothing. Do not get in eyes. Wear glasses or goggles to protect eyes from dust. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Wash thoroughly before eating or smoking.



STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Immediately flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water. If irritation persists get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

Groundwater Contamination: Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Do not apply when weather conditions favor wind drift of product from target area since this product will injure beans, cucumbers, grapes, roses, tomatoes, and most vegetables, flowers and ornamental plants. Do not apply where run-off is likely to occur. Do not contaminate water used for irrigation or domestic purposes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in its original container in a cool, dry, locked place out of reach of children.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

HOUSEHOLD: Do not reuse empty container. Wrap and put in trash.

GENERAL INFORMATION

This product may be used for control of many broadleaved weeds in residential and commercial lawns, recreational turf areas, golf courses (except putting greens), and other ornamental turf areas. Do not apply this product on sod farms.


This product is most effective when weeds are small and actively growing and should be applied when the temperature ranges from 65 to 90 degrees F. Do not apply when wind speed is sufficient to cause drift. Wind-drifted product may injure susceptible plants. If applied during extreme temperatures (below 65 or above 95 degrees F.) weed control will be erratic. Avoid applying within 2 hours of expected rain. If product is washed off within 2 hours after application re-treat in 7 to 10 days. Reduced weed control and fertilizer "streaking" will result if product is poorly applied. Deeprooted perennial weeds and locally resistant weeds may not be adequately controlled by this product.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

WHERE TO APPLY: Use on: Bluegrass, Ryegrass, Fescue, Bermuda, St. Augustine, Bahia or Zoysiagrass. St. Augustine is sensitive to most herbicides and may be injured by this product. Do not use on: Newly seeded lawns; Newly sprouted lawns until after the second mowing; Centipedegrass; Dichondra; Bentgrass; or where desirable clovers are present.

WHEN TO APPLY: For best results apply to moist established lawns when temperature ranges from 65 to 90 degrees F. in the spring or early summer when weeds are young and actively growing. Adequate soil moisture must be present for active growth to occur. A late summer or early fall application is also recommended to control broadleaf weeds that have developed from seeds during the summer. The lawn should be mowed two days prior to treatment. Wait 2 to 3 weeks to reseed areas where weeds have died. Dead weeds can be removed by raking or tilling.

The maximum number of broadcast applications per treatment site is 2 per year.



HOW TO APPLY: If grass is not moist, sprinkle it lightly with water before application to hold WEED AND FEED. Select a suitable Lawn Fertilizer Spreader and apply to turf by "marking your spreader path" to avoid double applications or total missing of turf area. During application, close spreader openings when turning. Apply 18 pounds of this product to cover 5000 sq. ft. of turf area. Do not water in. Thoroughly water the lawn the 3rd or 4th day to move the fertilizer into the root zone to stimulate new lawn growth as weeds die. Lawns that are heavily overrun by weeds or are in poor condition may require another application in approximately 30 days.

NOTE: When applying to St. Augustine or Bahia grasses under conditions of inadequate moisture, high temperature and high humidity, use 18 pounds per 10,000 sq. ft. to avoid possible dehydration or browning of grasses. A second application of 18 pounds per 10,000 sq. ft. should be made in about 30 days.

SUGGESTED SPREADER SETTINGS

The following spreader settings are for use with new equipment at average walking speed on level terrain. Your used spreader may need calibrating. Since proper application is based on the spreader's condition with regard to age and use, your walking speed, and the evenness of the terrain, your spreader may require a slight adjustment. If you have reason to believe that your spreader is not calibrated correctly, test apply by placing the amount for 500 sq. ft. into your spreader and apply to your pre-measured 500 sq. ft. (5 ft. wide by 100 ft. long) as follows:

HOW TO MAKE SURE YOUR SPREADER IS OPERATING CORRECTLY

For the 18 lb. rate to cover 5,000 sq. ft. using a Scott Spreader, place 1.8 lbs. of this product in your spreader using No. 4 setting and apply to 500 sq. ft. With a 20 inch spreader this will equal 3 spreader widths wide (3 passes) by 100 feet long. If you run out before you cover the pre-measured area, close the spreader setting by 1/2 the number. If you have granules left over, open (increase) the spreader setting by 1/2 the number. Use similar procedure for other brands of spreaders. For further help refer to your Spreader Guide book.

SPREADER SETTINGS*			
Spreader	18 pounds	Spreader	18 pounds
	per 5,000 sq. ft.**		per 5,000 sq. ft.**
Scott (20 inch)	#4	Jackson	#4 1/2
Central	#6	Modern	#5
Cyclone Model B	#2 3/4	Lawn Beauty	#5

NOTE: Broadcast spreaders such as the Cyclone are dependent on rate of walking and turning speed of the broadcast unit. Carefully calculate the area to be fertilized and make certain the proper number of pounds are applied to the area.

*For settings when using a new spreader.
**18 lbs./5000 sq. ft. is equivalent to 158 lbs./acre.

NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. TURF INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER. EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.

FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN STREET
FREMONT, NEBRASKA 68025