

METHOXYCHLOR 25 EC

ACTIVE INGREDIENTS:	BY WT.
Methoxychlor: 1,1,1-trichloro-2,2-bis (p-methoxyphenyl) ethane*	25.00%
INERT INGREDIENTS:**	75.00%
TOTAL	100.00%

One Gallon contains 2 lbs. Technical Methoxychlor
*Equivalent to 22% of 2,2-bis (p-methoxychlor)-1,1,1-trichloroethane and 3% other isomers and related compounds.
**This product contains xylene range aromatics.

KEEP OUT OF REACH OF CHILDREN CAUTION

See below for additional precautionary statements.

EPA REG. NO. 43789-45

EPA EST. NO. 43789-TX-01

NET CONTENTS _____ GALLONS

25021

EXP-1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes eye and skin irritation. Harmful if swallowed, inhaled, or absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid inhaling of spray mist. Do not use or store near food or feed.

Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work and wash all clothing with soap and hot water before reusing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Do not induce vomiting due to aspiration hazard. Gastric lavage is indicated if material is taken internally. Contact a physician or poison control center immediately.

If Inhaled: Remove victim to fresh air immediately. Apply artificial respiration if indicated.

If On Skin: Remove contaminated clothing and wash affected area with plenty of soap and water.

If In Eyes: Flush with plenty of water. Get medical attention if irritation persists.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT

Do not enter or allow entry into treated areas until spray has dried to perform hand labor tasks. A person may enter the area to perform other tasks only if the person is wearing the personal protective equipment listed on the label for a pesticide handler.

Do not apply this product in a way that will contact unprotected workers, either directly or through drift. Only protected handlers wearing long pants, long sleeved shirts, socks, shoes and chemical-resistant gloves may be in the area during application.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. The area being treated must be vacated by unprotected persons. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers.

Written or oral warnings must include the following information:

CAUTION

(Insert area or field description) treated with Methoxychlor on (insert date of application). Do not enter treated areas without protective clothing until sprays have dried. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Methoxychlor and describe his condition. See STATEMENT OF PRACTICAL TREATMENT.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE: If exposed to prolonged cold, place in warm storage (50 to 60 degrees F.) for several hours and agitate before using. Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PRODUCT INFORMATION

This product is an oil-base emulsifiable concentrate containing 2 pounds technical Methoxychlor per gallon. It may be diluted and mixed with water or certain oils as specified on this label or may be used as a direct application where recommended.

SEED TREATMENT USES

SEED TREATMENT: For control of insects such as cadelle, confused flour beetle, rice weevil, angoumois grain moth, Indian meal moth. Thoroughly mix the recommended amount of this product into the required amount of water for the slurry treater equipment and dilution rate to be used, or apply directly if treater equipment will provide adequate coverage in undiluted state. If used in undiluted state, do not expose treated seed to open flame or excessive heat. Dry seed thoroughly before stacking. If seed is stacked when wet, heat may develop which will impair germination. All treated seed should be stored in a cool, dry place until planted so that excess moisture may evaporate.

TYPE OF SEED

Corn
Cotton
Sorghum

OUNCES PER 100 POUNDS

1/4 fl. oz. - 3 fl. ozs.
1/2 fl. oz. - 3 fl. ozs.
1/2 fl. oz. - 3 fl. ozs.

Use maximum rates where insect infestations are severe.

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

(continued)

METHOXYCHLOR 25 EC

EPA REG. NO. 43789-45

(continued)

IMPORTANT:

Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Seed commercially treated with this product must be labeled in accord with all applicable requirements of the Federal Seed Act. Labels attached to treated seed must state "Treated seed must not be used for or mixed with food or animal feed, or processed for oil," and "Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water."

NOTE TO USER:

The purchaser of this product is responsible for insuring that all seeds treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR Part 2.25. Any dye added to treated seed must be cleared for use under 40 CFR part 180-1001.

STORAGE BINS AND AREAS

GRAIN STORAGE BINS, ELEVATOR TUNNELS (GALLERY FLOOR & HEADHOUSE), FREIGHT CARS, GRAIN TRUCKS AND TRANSPORT VEHICLES, SHIP HOLDS: Aids in control of cadelle, flat grain beetle, saw-toothed grain beetle, lesser grain borer, red flour beetle, long-headed flour beetle, rice weevil, granary weevil, confused flour beetle, foreign grain beetle, and hairy fungus beetle. Use 1 gallon in 10 gallons of water and apply 2 gallons of spray per 1,000 square feet. Treat when empty. Allow bins to air dry before refilling with grain, 2 to 4 weeks.

BUILDINGS AND STORAGE AREAS. Use as a contact and space spray for flies and mosquitoes. To reduce population of houseflies, stable flies, and mosquitoes in farm buildings, in barn yards, alleys, dumps, and other outdoor locations dilute one pint of this product with 2 1/2 gallons of water and apply with mist blower, hydraulic sprayer, hand sprayer, or any other suitable spray equipment. Care should be taken not to contaminate milk handling equipment, foodstuffs, or feed troughs and watering receptacles. Lactating dairy animals should not be present while spraying.

LIVESTOCK USES

BEEF CATTLE, SHEEP, SWINE, NON-LACTATING DAIRY CATTLE, NON-LACTATING GOATS: For control of hornflies, lice, ticks, fleas. Use one quart in 12 1/2 gallons of water and apply as a drenching spray. For fly control on cattle (except dairy animals) dilute 1 gallon with 4 gallons diesel fuel or furnace oil. Apply to backrubber device. Renew every 2 to 3 weeks.

MOSQUITO CONTROL USES

MOSQUITO BREEDING AREAS: Apply in undiluted or diluted form by ground sprayer or aircraft to known nonagricultural mosquito breeding sites such as parks, beaches, public dumps, municipal and residential areas. Prehatch larvicide: make a single application during the winter months at 1 to 2 quarts per acre. Larvicide: apply as needed during the spring and summer at 3/4 to 2 pints per acre.

AGRICULTURAL CROP USES

AGRICULTURAL CROPS: Apply at recommended rates as a spray with ground or aerial equipment to thoroughly cover foliage or fruit. Mix in water, using the appropriate volume of water for equipment to be used (for example, 2 to 20 dilution rate for aircraft or ground concentrate sprayers and greater amounts for high volume sprayers).

IMPORTANT NOTE: Do not apply within 14 days of sulphur-product applications, or within 7 days of Captan or Phaltan products.

Begin application at first sign of infestation (for fruits begin at petal fall) and repeat at 7 to 14 day intervals or as needed. Do not apply to crops within the number of days specified before harvest or grazing as shown by PHL (preharvest limitation).

ALFALFA, CLOVER, COWPEA, FORAGE GRASSES: For leafhopper, spittlebug use 1 to 2 quarts per acre. For alfalfa caterpillar, flea beetles, use 1 1/2 to 2 quarts per acre. For alfalfa weevil (larvae), webworms, fall armyworm, clover leaf weevil, mexican bean beetle, blister beetle, cucumber beetle, pea weevil, soybean caterpillar, velvet bean caterpillar, use 2 to 3 quarts per acre. For cowpea curculio, japanese beetle, use 3 quarts per acre. For armyworm use 3 to 4 quarts per acre. PHL — 7 days. Do not graze or feed treated cowpea vines or cowpea hay to livestock.


PEANUT, SOYBEAN: For velvet bean caterpillar, mexican bean beetle, japanese beetle, blister beetle, webworm, alfalfa webworm, cowpea curculio, leafhoppers, fall armyworm, use 2 to 6 quarts per acre. PHL — 7 days. Do not graze or feed treated soybean hay or straw to livestock.

WARRANTY

Manufacturer and Seller warrant that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors

as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Manufacturer and Seller. In no case shall Manufacturer and Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Manufacturer and Seller make no warranties of Merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Insecticide Liquid NOS RQ Methoxychlor NA1993



FORMULATED FOR
AGRICULTURAL PRODUCTS COMPANY
FT. WORTH, TX 76112