



ATRAZINE 90 WDG TURF AND CONIFER HERBICIDE

For Use on Southern Grasses and in Conifers.

ACTIVE INGREDIENTS:

Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine)	85.5%
Related compounds	4.5%
INERT INGREDIENTS:	10.0%
TOTAL	100.0%

handling.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

EPA REG. NO. 34704-
EPA EST. NO. 2393-M-3
NET WEIGHT 25 POUNDS

26A29

EXPOS#80

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, absorbed through skin, or inhaled. Avoid breathing dust. Causes eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.
 If On Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.
 If In Eyes: Flush with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to lakes, streams, or ponds. Use with care when applying in areas adjacent to any body of water. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes.
 Atrazine leaches readily and accepted label rates have been found to result in contamination of water supplies by way of groundwater. Therefore, users are advised to avoid use of Atrazine in well drained soils, particularly in areas having high groundwater tables.
 This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.
STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Personnel should use clothing and equipment consistent with good pesticide

Do not apply this product through any type of irrigation system.

GENERAL INFORMATION

This product controls most annual broadleaf and grass weeds in the crops included on this label.
 This product may be applied before or after weeds emerge.
 *Important Note: Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. These weeds may include annual bluegrass, yellow foxtail and others. Where this is known or suspected, we recommend the use of this product in combination with other registered herbicides which are not triazines. Consult your State Agricultural Extension Service about specific weed resistance to atrazine in your area and weed control recommendations. Within the specific crop directions on this label, certain weeds are indicated for suppression only. Weed suppression is a visual reduction in weed competition (reduce population and/or vigor) as compared to an untreated area. Degree of suppression will vary with rate used, size of weeds, and environmental conditions following treatments.
 Since this herbicide acts mainly through root absorption, its effectiveness depends on rainfall or irrigation to move it into the root zone, (except for already emerged weeds).
 Care should be taken to avoid using this product where adjacent desirable trees, shrubs, or plants might be injured.
 Note: Manufacturer does not recommend the use of this product, in combination with other herbicides or oils except as specifically described on this label or supplemental labeling distributed by manufacturer.

APPLICATION PROCEDURES

Small hand or power sprayers: This product should be mixed with water and applied as a spray. Pour product into the tank during or after filling. Agitation is recommended during mixing and application to keep the material in suspension. Wash sprayer thoroughly after use. Do not apply this product through any type of irrigation system.
Mixing procedures: (1) Be sure sprayer is clean and not contaminated with other products as plant injury may result. (2) Fill tank 1/2 to 3/4 full with clean water. (3) Add required amount of this product to tank. (4) Finish filling tank with water, and agitate mixture. (5) Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent.
NOTE: Adding other insecticides, herbicides, liquid fertilizer, or other materials is not recommended because they may cause compatibility problems or plant injury.
Commercial Ground application: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and of the same size. Calibrate sprayer before use.
 Use a pump with capacity to (1) maintain 35-40 psi at nozzle, (2) provide sufficient agitation in tank to keep mixture in suspension, and (3) to provide a minimum of 20% bypass at all times. Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gal./minute/100 gal. tank size circulated through a correctly positioned sparger tube or jets.
 Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or

BEST AVAILABLE COPY

ATRAZINE 90 WDG TURF & CONIFER HERBICIDE

EPA REG. NO. 34704-

coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For band applications, calculate amount to be applied per acre as follows:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

Aerial application: Use aerial application only where broadcast applications are specified. Apply in a minimum of 1 gal. of water for each 1 lb of this product applied per acre. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin, and should wash thoroughly before eating and at the end of each day's operation.

Commercial Ground and Aerial Application in water: Mixing procedures: (1) Be sure sprayer is clean and not contaminated with other materials. (2) Fill tank 1/2 to 2/3 full with clean water. (3) Start agitation. (4) Pour this product directly from package into tank. Let it wet and settle into the water. (5) Finish filling tank with water. Continue agitation during mixing and application.

NOTE: Adding other insecticides, herbicides, liquid fertilizer or other materials is not recommended because they may cause compatibility problems or plant injury.

Chemigation: Do not apply this product through any type of irrigation system.

TURFGRASSES FOR SOD (FLORIDA ONLY)

St. Augustinegrass, Centipedegrass, and Zoysia Grass
Broadcast 2.2-4.4 lbs./A according to soil texture. Apply in at least 15 gallons of water per acre, or 1 gallon per 1,000 sq. ft., for ground applications.

Broadleaf and Grass Weeds Controlled

Barnyardgrass* (watergrass)	cocklebur	nightshade*
giant foxtail*	groundcherry	pigweed
green foxtail*	jimsonweed	purslane
large (hairy) crabgrass	kochia	ragweed
wild oats	lambquarter*	sicklepod
witchgrass* (Panicum capillare)	annual	velvetleaf*
yellow foxtail*	morningglory	(burtonweed)
	mustards	

Muck or peat soil	4.4 lbs.	Old beds: Within 2 days after lifting sod New beds: 3-4 days after springing or plugging
Sandy soil	2.2 lbs.	Old beds: Within 2 days after lifting sod New beds: 7-10 days after springing or plugging

If weeds regrow, apply an additional 2.2 lbs./A on muck or peat or 1.1 lbs./A on sandy soil.

Precautions: To avoid crop injury, (1) Do not apply within 30 days prior to cutting or lifting. (2) Do not apply in combination with surfactants or other spray additives. (3) Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases. (4) On newly sprigged turfgrass, temporary slowing of growth may follow application.

TURFGRASS FOR FAIRWAYS, LAWNS, SOD PRODUCTION* AND SIMILAR AREAS

*In states other than Florida. For use on turfgrass for sod in Florida, see "Turfgrasses for Sod (Florida only)" section above.

Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysia Grass

Apply this herbicide after October 1 before emergence of winter annual weeds for control of annual bluegrass*, burclover, carpet burweed, chickweed, corn speedwell, henbit, hop clover, and spurge. This herbicide will control annual bluegrass even if it is emerged at time of treatment. For control of summer annual weeds listed in Turfgrasses for Sod (Florida Only), also apply this product in late winter before the weeds emerge. Apply in a minimum of 15 gals. of water per acre or 1 gal. per 1,000 sq. ft. Where annual bluegrass is the major weed, use 1.1 lbs. of this product per acre (0.4 oz. per 1,000 sq. ft.) Use 2.2 lbs./A (0.8 oz. per 1,000 sq. ft.) for control of the other weeds named above. Do not exceed 1.1 lbs./A per treatment on newly sprigged turfgrass or on hybrid bermudagrass such as Triflawn, Tifway, and Ormond.

(*Local resistance may result in suppression only or lack of control. Refer to IMPORTANT NOTE within GENERAL INFORMATION SECTION of this label).

For continued summer annual weed control, apply another 1.1 lbs./A at least 30 days after the previous application, but not after April 15. How-

ever, do not make more than two applications of this product per year.
Precautions: On newly sprigged turfgrass and hybrid bermudagrass, temporary slowing of growth and yellowing may occur following application. To avoid turf injury, (1) Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases. (2) Do not use on golf greens. (3) Do not use north of NC (except may be used in Virginia Coastal Plains) or west of the high rainfall areas of eastern OK and eastern TX. (4) Do not use on muck or alkaline soils. (5) Do not apply over the rooting area of trees or ornamentals not listed on this label. (6) Do not overseed with desirable turfgrass with 4 months before or 6 months after treatment. (7) Do not apply this product to newly seeded bermudagrass until it has overwintered and has a well-developed rhizome system. Do not exceed 2.2 lbs. product/A within 12 months of seeding bermudagrass.

Note: Do not graze or feed turf clippings to animals, or illegal residues may result.

CONIFERS

For control of annual broadleaf and grass weeds prior to transplanting, after transplanting or in established conifers (including Douglas fir, grand fir, noble fir, white fir, Austrian pine, bishop pine, Jeffrey pine, knobcone pine, loblolly pine, lodgepole pine (shore pine), Monterey pine, ponderosa pine, Scotch pine, slash pine, blue spruce, and Sitka spruce): Broadcast 2.2-4.4 lbs. in a minimum of 5 gals. of water per acre by air or 10 gals. by ground before weeds are 1.5 inches tall. Apply to established trees between fall and early spring while trees are dormant. For new transplants, apply during or soon after transplanting. For applications prior to transplanting, allow sufficient precipitation to activate herbicide before transplanting. In areas where spring and summer rainfall is inadequate to activate herbicide, apply during fall prior to spring transplanting.

Barnyardgrass* (watergrass)	cocklebur	nightshade*
giant foxtail*	groundcherry	pigweed
green foxtail*	jimsonweed	purslane
large (hairy) crabgrass	kochia	ragweed
wild oats	lambquarter*	sicklepod
witchgrass* (Panicum capillare)	annual	velvetleaf*
yellow foxtail*	morningglory	(burtonweed)
	mustards	

Quackgrass* control: Broadcast 4.4 lbs. in a minimum of 5 gals. of water per acre by air or 10 gals. by ground between fall and early spring while trees are dormant and before quackgrass is more than 1.5 inches tall.

(*Local resistance may result in suppression only or lack of control. Refer to IMPORTANT NOTE within GENERAL INFORMATION SECTION of this label.)

Precautions: (1) In areas west of the Rocky Mountains (except the Great Basin), grazing may begin 7 months after a fall application or 3 months after a winter or spring application. (2) To prevent illegal residues, do not graze treated areas of the Great Basin, or areas east of the Rocky Mountains. (3) Temporary injury to trees may occur following use of this product on coarse-textured soil. (4) To avoid crop injury, do not apply to seedbeds. (5) Also apply only once per year.

Aerial application: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft. above vegetation, using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

NOTE: In very hilly or mountainous terrain where the 10 foot flying height is unsafe, fly as low as possible. There may be increased risk of spray drift and uneven application.

Conifer Tank Mixes—Reforestation Areas: This product may be tank mixed with other herbicides, such as certain 2,4-D formulations, labeled for conifer site preparation and dormant conifer over-the-top applications. Users should carefully follow all label directions and precautions for each product, and obtain local use information from regional, state or university extension foresters, or similar reliable sources. The risk of conifer injury from tank mixes is generally unacceptable for Christmas trees and ornamentals. **IMPORTANT:** Tank mixes may increase potential for damage to conifers. Use only where the risk of damage can be tolerated. New combinations should be evaluated in limited tests before large scale use. Read WARRANTY AND LIMITATION OF DAMAGES for limit of manufacturer's warranty.

NOTICE

The Directions For Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Platte Chemical Co. or the Seller. All such risks shall be assumed by the Buyer.

**ATRAZINE 90 WDG
TURF & CONIFER HERBICIDE**

EPA REG. NO. 34704-

WARRANTY AND LIMITATION OF DAMAGES

Platte Chemical Co. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, and Conditions of Sale, subject to the inherent risks referred to therein. Platte Chemical Co. makes no other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Platte Chemical Co. is authorized to do so except in writing with a specific reference to this warranty. Any damages, arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or of value, etc.

ACCEPTED

United Federal Association of Pesticide and
Rodenticide Act, as amended, for the
pesticide register, 1975
EPA Reg. No. 34704-680

FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN ST. FRENONT, NEBRASKA 68025

TEST RESULTS 6-15-75