

yellow copy
p11
13

JUN 08 1992

Mr. J. Allen Dunlap III
Platte Chemical Co.
150 South Main Street
Fremont, NE 68025

Dear Mr. Dunlap:

Subject: Revised labeling
Azinphos-methyl 2EC (Sniper 2-E)
EPA Registration No. 34704-691
Amendment application of May 21, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable. A stamp copy is enclosed for your records.

Add the following restrictions statement:

"Do not graze treated fields."

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (H-7505C)

Enclosure

RESTRICTED USE PESTICIDE

Due to Acute Toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING CONTAINING INFORMATION WHICH MAY NOT APPEAR ON PACKAGE LABELS.

FOR DISTRIBUTION AND USE ONLY WITHIN LOUISIANA.



SNIPER™ 2-E AZINPHOS METHYL INSECTICIDE

EPA REG. NO. 34704-691

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE PACKAGE LABEL ARE TO BE FOLLOWED.

THIS SUPPLEMENTAL LABEL MUST BE IN POSSESSION OF USER AT THE TIME OF APPLICATION WHEN USED ON SUGARCANE IN THE STATE OF LOUISIANA.

THIS SUPPLEMENTAL LABEL REPLACES THE DIRECTIONS FOR USE ON THE CONTAINER AND BOOKLET OF THIS PRODUCT ON SUGARCANE ONLY IN THE STATE OF LOUISIANA. THESE DIRECTIONS FOR USE MUST BE FOLLOWED WHEN APPLYING THIS PRODUCT TO SUGARCANE IN LOUISIANA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Sugarcane (Conventional Spray) Louisiana only	Sugarcane borer	3	Apply specified dosage per acre by air application using a minimum of 2 gallons of water per acre. For best results applications should be made at approximately cane top level and not more than 10 feet above cane top level. Do not apply more than 3 times per season nor within 30 days of harvest. Bagasse from sugarcane treated with SNIPER 2-E may be used for feed of cattle, goats, and sheep. Consult your State Agricultural Extension Service or Experiment Station for specific use information.
--	--------------------	---	---

<p>Sugarcane (1) (Ultra Low Volume Spray) Louisiana only</p>	<p>Sugarcane borer</p>	<p>3</p>	<p>Apply specified dosage per acre undiluted. Apply SNIPER 2-E undiluted 1:1 (water to product). Aircraft must be equipped with low volume device that will produce droplets with the range of 30 to 100 microns. Apply at cane top level not to exceed 10 feet above cane top level. Do not apply more than 3 times per season nor within 30 days of harvest. Bagasse from sugarcane treated with SNIPER 2-E may be used for feed of cattle, sheep and goats. Consult your State Agricultural Extension Service or Experiment Station for specific use information.</p>
--	----------------------------	----------	--

- (1) Note: This formulation, when used undiluted, may cause spotting of automobile finishes if prolonged exposure is permitted. Do not spray directly over automobiles. If accidental exposure does occur, automobiles should be washed immediately.

RESTRICTIONS

Do not apply in the rain.

Do not make applications during temperature inversions.

A temperature inversion is a stable atmospheric condition characterized by an increase in air temperature with increased height above the ground until at some height a "ceiling" or barrier of colder air is met.

Make applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). In Louisiana do not apply when the wind velocity exceeds 10 mph.

For applications, the spray boom must be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vertices. Boom length must not exceed 75% of wing span or rotor diameter.

Buffer Zone: Do not apply within 75 feet of lakes, reservoirs, rivers, permanent streams, marshes or ponds, canals, estuaries and commercial fish farm ponds. Do not apply if the soil is saturated with water.

Do not apply under conditions that favor runoff.

Allow at least 21 days between applications and at least 30 days between the last application and harvest.

FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN STREET FREMONT, NE 68025

