

PM 25

34704-692

PM 184

PROMETRYNE



ACCEPTED
 APR 3 1996
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 34704-692

HERBICIDE

FOR SELECTIVE WEED CONTROL IN COTTON, CELERY, AND PARSLEY

ACTIVE INGREDIENT:	
Prometryne: 2,4-bis (isopropylamino)-6-(methylthio)-s-triazine	44.4%
INERT INGREDIENTS:	55.6%
TOTAL	100.0%

PROMETRYNE 4L contains 4 lbs. active per gallon

**KEEP OUT OF REACH OF CHILDREN
 WARNING AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

EPA REG. NO. 34704-692

EPA EST. NO. 34704-NB-2

NET CONTENTS 2 1/2 GALS. (9.46 L)

Promet[®] trademark of American Cyanamid for pendimethalin
 Reflan[®] trademark of Dow Blanco
 X-77[®] trademark of Chevron Chemical Co.
 Triton[®] is a Registered T.M. of Kalo, Inc.
 Triton[®] is a Registered T.M.

5392

EXP03899

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if inhaled. Harmful if swallowed or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin or clothing.

First Aid: If inhaled, move to clean air. If swallowed, give milk or water; if conscious, induce vomiting by placing finger in back of throat. Get medical attention. In case of contact with the eyes, flush with plenty of water for 15 minutes; for skin, wash with soap and water. Get medical attention if irritation persists.

Personal Protective Equipment:

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:
 Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink 1 or two glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.
If on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.
If in Eyes: Flush with plenty of water for 15 minutes. Get medical attention.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
 For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: if the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes plus socks.

Shake well before using.

284

PROMETRYNE 4L HERBICIDE

EPA REG. NO. 34704-692

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage of disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

STORAGE: Store in a safe manner. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang can affect package strength.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide and Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PROMETRYNE 4L is a selective herbicide that may be applied either before or after weeds emerge for control of most annual broadleaf weeds and grasses, including groundcherry, lambquarters, annual morning-glory, malva, mustard, black nightshade, pigweed (carelessweed), purslane, Florida pusley, ragweed, smartweed, teasel (prickly sida), barnyardgrass (watergrass), crabgrass, foxtail, goosegrass, Jungieria, Panicum spp., signalgrass (and other Brachiaria spp.), and wild oats. PROMETRYNE 4L also controls shallow-germinating seedlings of cocklebur, coffeeweed, and sandbur. It does not control johnsongrass, bermudagrass, other established perennials, or sprangletop at selective rates.

When applied before weeds emerge, PROMETRYNE 4L enters weeds through their roots. Thus, its effectiveness depends on moisture to move it into the soil. Under very dry soil conditions after application, a shallow cultivation or rotary hoeing will generally result in better weed control.

When applied to emerged weeds, PROMETRYNE 4L provides foliar knockdown and/or residual control of later germinating weeds, depending on the rate applied.

Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected, we recommend the use of this product in combination with other registered herbicides which are not triazines. Consult with your State Agricultural Extension Service for specific recommendations.

FAILURE TO FOLLOW DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR PERFORMANCE, AND/OR ILLEGAL RESIDUES.

Equipment and Mixing

Thoroughly clean sprayer prior to use. Do not use a sprayer contaminated with 2,4-D or other materials, as crop damage or sprayer clogging may result.

Use conventional spray equipment with hydraulic or mechanical agitation. Screens in nozzles and in suction and in-line strainers should be no finer than 50-mesh. Use a pump with capacity to maintain 35-40 psi at the nozzles, and provide sufficient agitation in tank to keep mixture in suspension. A centrifugal pump which provides propeller shear action is recommended for dispersing and mixing this product. Spray solution should be bypassed through a correctly positioned sparger tube or jets.

Nozzles: For preplant incorporated or preemergence application, use flat fan nozzle tips. For postemergence band application, use off-center nozzle tips. For postemergence broadcast application, use flat fan or off-center nozzle tips. Use flood nozzle tips only in Arizona and California for lay-by treatment in cotton at least 18 inches tall.

Mixing: (1) Fill spray tank 1/2 to 3/4 full with clean water. (2) Start agitation. (3) Pour the product directly from the container into the spray tank partially filled with water, and then add the rest of the water. (4) Provide agitation during mixing and application to maintain a uniform suspension.

Cleaning: Wash sprayer thoroughly with clean water immediately after use. Do not use the same sprayer on sensitive crops, as even small residues of PROMETRYNE 4L in the tank may cause injury to these crops.

CHEMIGATION:

Refer to supplemental labeling entitled "APPLICATION THROUGH IRRIGATION SYSTEMS—CHEMIGATION" for use directions for chemigation. Do not apply this product through any irrigation systems unless the supplemental labeling on chemigation is followed.

Ground Application (All uses):

Use conventional ground sprayer equipped with nozzles that provide accurate and uniform application.

Calibrate sprayer before use. Unless otherwise specified, use a minimum of 20 gals. of spray mixture per acre for all preplant incorporated, preemergence, and postemergence applications with ground equipment.

Aerial Application (Cotton only):

Use aerial application only where broadcast applications are specified. Use a minimum of 5 gals. of spray mixture per acre. Avoid applications under conditions where uniform coverage cannot be obtained or where spray drift may occur.

To assure that spray will be controllable within the target area when used according to the label directions, make applications at a maximum height of 10 ft. above vegetation, using low drift nozzles at a maximum pressure of 40 psi., and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply this product at a minimum upwind distance of 400 ft. from sensitive plants.

Seedbed Preparation

To insure proper placement of PROMETRYNE 4L, seedbeds must be well prepared and as free as possible from trash and clods. A firm seedbed is best for obtaining effective weed control. Uniformity in height and width of seedbed is essential for proper postemergence applications of PROMETRYNE 4L. Beds should be low and flat. Take care to avoid planter marks. Wide planter packing wheels or rollers are recommended. Wheel furrows should be uniform in depth. Mount the sprayer so that it follows the same rows as the planter.

Band Treatment

For band treatment, calculate the amount of PROMETRYNE 4L needed by the formula:

$$\frac{\text{Band width in inches}}{\text{row width in inches}} \times \frac{\text{broadcast rate}}{\text{per acre}} = \frac{\text{amount needed}}{\text{per acre of field}}$$

COTTON

PROMETRYNE 4L may be applied preplant incorporated or preemergence and/or postemergence as recommended in the following tables. The postemergence applications may follow preplant incorporated or preemergence treatments of PROMETRYNE 4L.

Do not use on glandless cotton varieties or crop injury will occur.

Note: Do not feed treated forage to livestock, or graze treated areas.

Preplant Incorporation (Arizona, California, and New Mexico)

Apply PROMETRYNE 4L at the appropriate rate in Table 1 as a broadcast or band treatment. If broadcast, treat the flat soil surface prior to listing. If banded, apply over partially finished or finished beds. Incorporate up to 4 inches deep immediately after application with PTO-driven equipment, double disk, rolling cultivator, or bed conditioner.

Table 1: Preplant Incorporation

Region	Soil texture	Broadcast rate per acre
Arizona, California, and New Mexico	sand, loamy sand	Do not use
	sandy loam (AZ & CA only)	2.4-3.2 pts.
	sandy loam, loams (NM only)	3.2 pts.
	silt loam, clay	4.8 pts.

Precautions: Do not use PROMETRYNE 4L in cut areas of newly leveled fields, in areas of excess salt, or in areas where flooding over the beds is likely to occur or crop injury may result. Do not plant cotton in tractor wheel depressions or crop injury may result. On mulch planted cotton, water back only after cotton seedlings are well established. In New Mexico, apply either preplant incorporated or preemergence (not both)—see Preemergence section.

Preemergence

Apply at planting or shortly after planting at the appropriate rate in Table 2. PROMETRYNE 4L may be used on cotton planted flat, on beds, or in furrows. To avoid concentration of PROMETRYNE 4L in the seed furrow, do not make broadcast applications to cotton planted in furrows deeper than 2 inches. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow. If banded, do not cover treated bands with soil while cultivating untreated row middles. Do not use on sand or loamy sand, on shallow soils with caliche subsoils, or in areas with caliche overcropping.

Cotton may be replanted in soil previously treated with PROMETRYNE 4L. Do not apply a second preemergence application of PROMETRYNE 4L or injury may occur.

Table 2: Preemergence

Region	Soil texture	Broadcast rate per acre
Mid-South and Southeast other than Mississippi River Delta in Mississippi	sandy loam	3.2-4 pts.
	silt and clay loam	4.8 pts.
	Sharkey clay (Arkansas only)	5.6 pts.
Mississippi River Delta in Mississippi	sandy loam	4.4-5 pts.
	silt and clay loam	5.6 pts.
	Sharkey clay	Do not use
Blacklands of Texas and Okla., Texas Gulf Coast, and Texas Coastal Bend	loam	2.4 pts.
	clay	4.8 pts.
Rio Grande Valley of Texas	loam	3.2 pts.
	clay	4.8 pts.
High Plains, Rolling Plains and Edwards Plateau of Texas	sand, loamy sand	Do not use
	sandy loam	1.6 pts.
	loam, sandy clay loam	2.4 pts.
Southwest Texas and New Mexico	other clay soils	3.2 pts.
Arizona and California		Do not use

This * portion was reset

Added this me-too portion to label

484

PROMETRYNE 4L HERBICIDE

EPA REG. NO. 34704-692

Trifluralin Split Application (Arizona and California)

Apply a preplant incorporated application of trifluralin as directed on that label, except use the appropriate rate from Table 6. Do not apply trifluralin before January 1. Follow at planting or just before planting with a preplant incorporated treatment of PROMETRYNE 4L as directed in the Cotton section of this label, except use the appropriate rate from Table 6.

DSMA or MSMA

For faster knockdown of the weeds controlled by PROMETRYNE 4L alone, apply 1-1.3 pts. of PROMETRYNE 4L plus 3 lbs. active ingredient of DSMA or 2 lbs. active ingredient of MSMA per acre, following the same directions, precautions, and limitations as given on this label for PROMETRYNE 4L applied alone postemergence directed (chemical hoe.) If needed, make a second application 1-3 weeks after the first application. Do not apply after first bloom.

Several formulations of DSMA and MSMA are available under various trade names from several manufacturers. Observe the directions, limitations, and precautions on the label of the product used.

CELERY

Seedbeds (Florida)

Broadcast 1.2-1.6 pts. in a minimum of 20 gals. of water per acre after celery has 2-5 true leaves. Application may be made over the celery. Apply only after seedbed covers have been removed from seedbeds for at least one week. Apply only once per year to seedbeds.

Direct-seeded Celery (California only)

Apply PROMETRYNE 4L at rates given below in a minimum of 20 gals. of water per acre. Within the rate ranges given, use the lower rates on coarse-textured soils and soils low in organic matter; use the higher rates on fine-textured soils and soils high in organic matter.

Preemergence: Broadcast 2.4-3.2 pts. per acre at planting or shortly after planting before celery emerges.

Postemergence: Broadcast 1.6-2 pts. per acre after celery has 2-5 true leaves. Application may be made over the celery. Apply before weeds are 2 inches tall.

To Avoid Injury to Direct-seeded Celery: (1) Make either one preemergence or one postemergence application (not both) per celery crop. (2) Do not use on sand or loamy sand. (3) Do not apply if celery is under water stress. (4) Do not apply postemergence treatments of PROMETRYNE 4L with other pesticides. Apply only after foliar applications of other pesticides are dry. (5) Do not apply within two weeks after an application of a herbicidal oil, such as "carrot" oil.

Transplants

Apply one application at the appropriate rate from Table 7 in a minimum of 20 gals. of water per acre during the 2-8 week period after transplanting. Within the rate ranges given, use the lower rate on relatively coarse-textured soils and soils low in organic matter; use the higher rate on relatively fine-textured soils and soils high in organic matter. Application may be made over the celery. Apply before weeds are 2 inches tall.

Table 7: Transplanted Celery

State	Soil	Broadcast rate per acre
Florida	sandy or muck	1.6-3.2 pts.
California	coarse-textured	2-3.2 pts.
	fine-textured	3.2-4 pts.
Hawaii	coarse-textured	3.2-4.8 pts.
	fine-textured	4.8-6.4 pts.
Michigan	fine-textured or muck	2-4 pts.
Wisconsin	fine-textured	3.2-4 pts.

Rotational Crops

The following crops may be seeded 5 months after applying no more than 4 pts. PROMETRYNE 4L per acre on celery: Cabbage, celery, corn, onions, peas, and red beets.

PARSLEY FOR USE IN CALIFORNIA ONLY

For control of cheeseweed, burning nettle and shepherd's purse, apply by ground, 1-2 quarts of product per acre (or 1-2 pounds ai/acre) in a minimum of 30 gallons of water per acre. Use the lower rate on coarse texture soils and soils low in organic matter. Use the higher rate on fine texture soils and soils high in organic matter.

To Avoid Injury to Parsley: (1) Make one preemergence application per parsley crop. (2) Do not use on sand or loamy sand soil. (3) Do not apply if parsley is under water stress. (4) Do not apply within two weeks after an application of a herbicidal oil, such as "carrot" oil.

Preemergence: Apply as a single application following planting, before parsley emerges. Do not enter treated areas until spray residues have dried. Do not apply within 84 days of harvest.

NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

FORMULATED FOR
PLATTE CHEMICAL CO.

150 SO. MAIN STREET FREMONT, NEBRASKA 68025-5697