

Reg # 34704-712

PM-14

126

# RESTRICTED USE PESTICIDE

DUE TO ACUTE ORAL AND DERMAL TOXICITY; AND BIRD TOXICITY:  
FOR RETAIL SALE TO AND APPLICATION ONLY BY CERTIFIED APPLICATORS OR  
PERSONNEL UNDER THEIR DIRECT SUPERVISION.

## RAMPART 10-G

### SOIL AND SYSTEMIC INSECTICIDE (24/48 MESH)

ACCEPTED	
APR 20 1987	
Under the Federal Insecticide, Fungicide, and Rodenticide Act,	
as amended, this product is registered under	
EPA Reg. No.	2749-121

→ Transferred to 34704-712.

ACTIVE INGREDIENT .....	10%
Phorate (O,O-Diethyl S-[(ethythio)-methyl] phosphorodithioate)	
INERT INGREDIENTS .....	90%
TOTAL .....	100%

### DANGER POISON



### PELIGRO

KEEP OUT OF REACH OF CHILDREN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada completamente.

NET WEIGHT \_\_\_\_\_ POUNDS

EPA REG. NO. 2749-121

EPA EST. NO. \_\_\_\_\_

Oral warnings must include the same information that is in the written warnings.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

**STORAGE:** Store in safe manner. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### GENERAL DIRECTIONS

WHEN USING RAMPART 10-G USE BY POURING DOWN WIND TO MINIMIZE EXPOSURE TO HANDLERS. APPLY RAMPART 10-G WITH A GRANULAR PESTICIDE APPLICATOR PROPERLY CALIBRATED TO ASSURE ACCURATE PLACEMENT AND PROPER DOSAGE.

FOODFEED CROPS OTHER THAN THOSE REGISTERED FOR USE WITH PHORATE MAY NOT BE PLANTED IN PHORATE-TREATED SOILS.

#### NOTICE OF WARRANTY

Acute warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label, only when used in accordance with directions under normal use conditions.

**SOIL AND SYSTEMS**

APR 20 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under EPA Reg. No. 2749-121

**ACTIVE INGREDIENT** ..... 10%  
**Phorate (O,O-Diethyl S-[(ethylthio)-methyl] phosphorodithioate)**  
**INERT INGREDIENTS** ..... 90%  
**TOTAL** ..... 100%

**DANGER POISON**  **PELIGRO**

**KEEP OUT OF REACH OF CHILDREN**

**PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.**

**NET WEIGHT** \_\_\_\_\_ **POUNDS**

**EPA REG. NO. 2749-121**

**EPA EST. NO.** \_\_\_\_\_

EXP-11

**PRECAUTIONARY STATEMENTS  
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER**

toxic by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Do not get in eyes, on hair or clothing. Wear freshly laundered long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton). If a skin glove is used, they must be laundered after each use. Do not wear the same gloves for other work. Destroy and replace gloves immediately. In case of contact, immediately remove contaminated clothing. Wash thoroughly with soap and water before eating or smoking. Do not breathe dust. While emptying bags into equipment, pour downward and as far from face as possible. Do not pour at face level and do not allow dust to reach breathing zone. Do not use or store in or around the house. Keep away from domestic animals and feedstuffs.

**ENVIRONMENTAL HAZARD**

This product is toxic to fish, shrimp, crabs, and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crabs may be killed at the recommended dosage rates. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

**STATEMENT OF PRACTICAL TREATMENT**

**SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.

**ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

**IN EYES:** Flush with plenty of water for 15 minutes. Get medical attention.

**CASE OF POISONING, CONTACT PHYSICIAN IMMEDIATELY**

**NOTE TO PHYSICIANS:** Acetate is Antidote. S-PHOS is also Antidote and may be administered in conjunction with Acetate. Worsening symptoms include weakness, headache, dizziness in chest, blurred vision, nausea, vomiting, diarrhea, and abdominal pain.

Oral warnings must include the same information that is in the written warnings.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

**STORAGE:** Store in safe manner. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixtures, or wastes is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Compulsorily empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**GENERAL DIRECTIONS**


**WHEN USING RAMPART 10-G USE BY POURING DOWN: WIND TO MINIMIZE EXPOSURE TO HANDLERS. APPLY RAMPART 10-G WITH A GRANULAR PESTICIDE APPLICATOR PROPERLY CALIBRATED TO ASSURE ACCURATE PLACEMENT AND PROPER DOSAGE.**

**FOODFEED CROPS OTHER THAN THOSE REGISTERED FOR USE WITH PHORATE MAY NOT BE PLANTED IN PHORATE-TREATED SOILS.**

**NOTICE OF WARRANTY**

Aceto warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label, only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aceto. In no case shall Aceto be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Aceto makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

FORMULATED FOR  
**ACETO AGRICULTURAL CHEMICALS CORPORATION**  
 125-82 Northern Blvd., Flushing, New York 11358



**POISON**

NET WEIGHT \_\_\_\_\_ POUNDS

EPA REG. NO. 2749-121

EPA EST. NO. \_\_\_\_\_

15845

EXP-1R

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### DANGER

Poisonous by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Do not get in eyes, on skin or clothing. Wear freshly laundered long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton.) If cotton gloves are used, they must be laundered or discarded after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently. In case of contact, immediately remove contaminated clothing. Wash thoroughly with soap and water before eating or smoking. Do not breathe dust. While emptying bags into equipment, pour downward and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach breathing zone. Not for use or storage in or around the home. Keep away from domestic animals and foodstuff.

#### ENVIRONMENTAL HAZARD

This product is toxic to fish, shrimp, crabs, and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crabs may be killed at the recommended dosage rates. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of waste. Apply this product only as specified on this label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.  
**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.  
**IF IN EYES:** Flush with plenty of water for 15 minutes. Get medical attention.  
**IN CASE OF POISONING, CONTACT PHYSICIAN IMMEDIATELY.**  
**NOTE TO PHYSICIAN:** Atropine is Antidote. S-PAM is also Antidote and may be administered in conjunction with Atropine. Warning symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

#### DIRECTION FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### RE-ENTRY STATEMENT

Do not reenter treated fields before 24 hours after application. If entry prior to 24 hours is required, take the same precautions as an applicator of the product including use of protective clothing, gloves, and goggles. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

#### DANGER

(insert area or field description) treated with Phorate on (insert date of application.) Do not reenter treated fields before 24 hours after application. If reentry prior to 24 hours is required, take the same precautions as an applicator of the product including use of protective clothing, gloves, and goggles. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Phorate and describe his condition. For further information see the "NOTE TO PHYSICIAN."


ACCURATE PLACEMENT AND PROPER DOSAGE.

FOODFEED CROPS OTHER THAN THOSE REGISTERED FOR USE WITH PHORATE MAY NOT BE PLANTED IN PHORATE-TREATED SOILS.

#### NOTICE OF WARRANTY

Aceto warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label, only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aceto. In no case shall Aceto be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Aceto makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

FORMULATED FOR  
ACETO AGRICULTURAL CHEMICALS CORPORATION  
125-82 Northern Blvd., Flushing, New York 11368



# POISON

6

Organic Phosphate Compound  
Mixture, Dry NA2783

# RAMPART 10-G

## SOIL AND SYSTEMIC INFECTICIDE

### DIRECTIONS FOR USE

Apply with a properly calibrated granular pesticide applicator.

GROUP	PESTS CONTROLLED	RATE RAMPART 10-G	APPLICATION	REMARKS
BEANS	Minion bean beetle	10 to 20 lbs. per acre (20-inch row spacing) or 9 to 19 oz. per 1000 ft. of row for any row spacing (minimum 20-inch spacing.)	Distribute the granules evenly in the row in the site of the seed at planting time. Do not place RAMPART 10-G in direct contact with the seed.	Do not feed the foliage of treated beans within 60 days of treatment.
	Leafhopper, aphids, legume bugs, thrips, mites, and seed corn maggot	10 to 15 lbs. per acre (20-inch row spacing) or 9 to 14 oz. per 1000 ft. of row for any row spacing (minimum 20-inch spacing.)		
FIELD and BURET CORN At Planting Time	Corn rootworm, mites	10 lbs. per acre (40-inch row spacing) or 10 oz. per 1000 ft. of row for any row spacing (minimum 20-inch row spacing)	Place granules in a 3-inch band over the row directly behind the planter shoe in front of the press wheel.	Do not place RAMPART 10-G granules in direct contact with seed. *On corn grown only in Iowa and Illinois
	After Plant Emergence	Corn rootworm	Apply granules in a band of cultivation in a band at base of plants just ahead of cultivator should be so to cover granules with soil. Apply in June as soon as rootworm activity is observed.	Do not make any later applications after plant emergence treatment.
FIELD CORN After Plant Emergence	Corn leaf aphid, mite	10 lbs. per acre (40-inch row spacing) or 10 oz. per 1000 ft. of row for any row spacing (minimum 20-inch row spacing.)	Apply granules into the whorl of the plant.	Do not graze or cut for silage within 30 days of treatment.
	Prior to "head emergence (silks only one application after plant emergence)	European corn borer (1st brood)**	Apply granules into the whorl of the plant.	Consult state experiment station or or state extension service for proper timing of application. **On corn grown only in North Central States
COTTON	Early season thrips, aphids, mites, and leaf miners	5 lbs. per (40-inch row spacing) or 5 oz. per 1000 ft. of row for any row spacing (minimum 20-inch spacing.)	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture, and pH should be favorable for germination and emergence.	Recommended for areas where a limited period of protection against early season pests is necessary for good cotton growth. Use 2 lb. RAMPART 10-G per acre for late-dropped cotton.
	Mites, thrips, aphids, black cutworm, leafhopper, white fly and leaf miners	10 to 15 lbs. per acre (40-inch row spacing) or 10 to 15 oz. per 1000 ft. of row for any row spacing (minimum 20-inch spacing.)	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture, and pH should be favorable for good germination and emergence. Under adverse conditions plant stand may be affected.	Recommended for areas requiring an extended period of early season protection against other cotton pests. The higher rate is suggested for heavier soils.
	After Plant Emergence (Original cotton only)	Mites from spring to early summer	10 to 15 lbs. per acre (40-inch row spacing) or 10 to 15 oz. per 1000 ft. of row for any row spacing (minimum 20-inch spacing.)	Incorporate granules into soil as a side-dress application deep enough to avoid disturbance by future cultivation.
PEANUTS At Planting Time	Thrips, leafhoppers	10 lbs. per acre (40-inch row spacing) or 10 oz. per 1000 ft. of row for any row spacing (minimum 20-inch spacing.)	Distribute the granules evenly in the furrow at planting time. For best results, plant peanuts when soil temperatures are favorable for establishing good peanut stands.	Do not graze or feed treated hay or forage.
PEANUTS At Planting Time	Stem-borer and leafhoppers	10 lbs. per acre (40-inch row spacing) or 10 oz. per 1000 ft. of row for any row spacing (minimum 20-inch spacing.)	Distribute the granules in a band over the planting zone at planting time. Work into the top two inches of soil immediately.	This treatment will also control the incidence of leafhoppers. Do not graze or feed treated hay or forage.
PEANUTS	Aphids, leafhoppers, leaf miners, grubs, cutworms, and other soil insects and mites that injure the roots	10 lbs. per acre (40-inch row spacing) or 10 oz. per 1000 ft. of row for any row spacing (minimum 20-inch spacing.)	Distribute the granules evenly in the furrow at planting time. Work into the top two inches of soil immediately.	Use 30 days after treatment before harvesting peanuts.

96n

After Plant Emergence	See evidence	10 lbs. per acre (40 bush row spacing) or 12 lbs. per 1000 ft. of row for any row spacing (minimum 20-bush row spacing)	Apply granules in a band 4 inches of plants just closed of cut-water channels or in to lower granules with soil. Apply in June to meet to maximum ability to absorb.	Do not make any later applications after plant emergence treatment.
WHEAT CORN After Plant Emergence	See leaf aphids, etc.	10 lbs. per acre (40 bush row spacing) or 12 lbs. per 1000 ft. of row for any row spacing (minimum 20-bush row spacing)	Apply granules in the wheel of the plan.	Do not grow or cut for forage within 30 days of treatment.
Wheat to Small Emergence (Make only one application after plant emergence)	Emergence soon later (at bread)		Apply granules into the wheel of the plan.	Consult state experiment station or or state extension service for proper timing of application. **Do not grow only in North Central States
COTTON At Planting Time	Early season thrips, aphids, mites, and leaf miners	5 lbs. per (40-bush row spacing) or 6 lbs. per 1000 ft. of row for any row spacing (minimum 20-bush spacing)	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture, and pH should be favorable for good germination and emergence.	Suggested for areas where a limited period of protection against early season pests is necessary for good cotton growth. Use 5 lbs. R445-PWV 10-6 per acre for 40-bush rows.
At Planting Time	Mites, thrips, aphids, leaf miners, leafhoppers, white flies and leaf miners.	10 to 15 lbs. per acre (40-bush row spacing) or 12 to 15 lbs. per 1000 ft. of row for any row spacing (minimum 10-: in spacing)	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture, and pH should be favorable for good germination and emergence. Under adverse conditions plant stand may be affected.	Recommended for areas requiring an extended period of mite control as well as early season protection against other cotton pests. The higher rate is suggested for heavier soils.
COTTON At Side-Dressing (Delayed or Non only)	Mites from spring to early summer.	3.5 to 4 lbs. per acre, 40-bush row spacing or 4 to 5 lbs. per 1000 ft. of row for any row spacing (minimum 20-bush spacing)	Incorporate granules into soil as a side-dress application deep enough to avoid disturbance by future cultivation.	Integrate as soon as possible after treatment. Do not apply later than 60 days before harvest.
PEANUTS At Planting Time	Thrips, leafhoppers	10 lbs. per acre (20-bush row spacing) or 12 lbs. per 1000 ft. of row for any row spacing (minimum 20-bush spacing)	Distribute the granules evenly in the furrow at planting time. For best results, plant peanuts when soil temperatures are favorable for establishing good percent stands.	Do not grow or feed treated hay or forage.
PEANUTS At Planting Time	Stem-borer root-miners and leafhoppers	10.5 lbs. per acre (20-bush row spacing) or 12 lbs. per 1000 ft. of row for any row spacing (minimum 20-bush spacing)	Distribute the granules in a band over the planting zone at planting time. Work into the top few inches of soil immediately.	This treatment will also control the infestation of leafhoppers. Do not grow or feed treated hay or forage.
POPPLES At Planting Time	Aphids, leafhoppers, leaf miners, psyllids, thrips, etc. (See leafhoppers and other pests of the beetle white)	20 lbs. per acre (40-bush row spacing) or 24 lbs. per 1000 ft. of row for any row spacing (minimum 20-bush spacing). 20 lbs. for heavy soil or clayey soil or 24 lbs. for sandy soil per 1000 ft. of row for any row spacing (minimum 20-bush spacing)	Distribute the granules evenly in the furrow or granules may be banded on each side of the row at planting time.	Wait 30 days after treatment before harvesting produce.
SOYBEANS At Planting Time	Aphids, greenbugs	10 lbs. per acre (40-bush row spacing) or 12 lbs. per 1000 ft. of row for any row spacing (minimum 20-bush spacing)	Form granules 2 inches to side and 1 inch below the seed at planting or in a 2-inch band over the row just prior to press wheel.	Do not place R445-PWV 10-6 in direct contact with the seed. Use material in Western States.
SOYBEANS (Early)	Aphid, greenbugs	10 lbs. per acre	Apply granules manually by air or ground equipment.	Do not apply within 30 days of harvest. Make only one post-emergence application. Use material in Western States.
SOYBEANS (Late)	Aphids, leafhoppers, etc.	10 lbs. per acre (40-bush row spacing) or 12 lbs. per 1000 ft. of row for any row spacing (minimum 20-bush spacing)	Application should be made by killing or broadcast at planting time.	Do not place R445-PWV 10-6 in direct contact with the seed.  Do not apply within 30 days of harvest.
	Aphids, etc.	10 to 15 lbs. per acre	Apply to foliage when plants are dry. Use the higher rate on dense foliage for good coverage.  Foliar application by air may be used for good control when soil conditions prevent ground application.	Do not feed sugar beet tops or other to dairy cattle.  Consult your state experimental station or state extension service for proper timing of application in your area.

965

Prior to Seed Emergence (Make only one application after plant emergence)	Emergence and later (No broadcast)	20-inch row spacing	Apply granules to the seed of the plant.	Consult state experiment station or state extension service for proper timing of application. **On one grower only in North Central States
<b>COTTON</b> At Planting Time	Early season borers, aphids, white, and leaf miners	5 lbs. per acre (40-inch row spacing) or 6 lbs. per 1000 ft. of row for any row spacing (minimum 20-inch spacing)	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture, and pH should be favorable for good germination and emergence.	Suggested for areas where a limited period of protection against early season pests is necessary for good cotton growth. Use 2 lbs. FAMBART 10-G per acre for late-planted cotton.
<b>COTTON</b> At Flowering Time	Miss,蓟, aphids, black cutworms, leafhoppers, white fly and leaf miners	10 to 15 lbs. per acre (40-inch row spacing) or 12 to 18 lbs. per 1000 ft. of row for any row spacing (minimum 20-inch spacing)	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture, and pH should be favorable for good germination and emergence. Under adverse conditions plant stand may be affected.	Recommended for areas requiring an extended period of white control as well as early season protection against other cotton pests. The higher rate is suggested for heavier soils.
<b>COTTON</b> At Seed-Setting (Delayed cotton only)	Miss from spring to early summer	15 to 20 lbs. per acre (40-inch row spacing) or 18 to 24 lbs. per 1000 ft. of row for any row spacing (minimum 20-inch spacing)	Incorporate granules into soil as a side-dress application deep enough to avoid disturbance by future cultivation.	Irrigate as soon as possible after treatment. Do not apply later than 60 days before harvest.
<b>PEANUTS</b> At Planting Time	Termites, leafhoppers	10 lbs. per acre (20-inch row spacing) or 11 lbs. per 1000 ft. of row for any row spacing (minimum 24-inch spacing)	Distribute the granules evenly in the furrow at planting time. For best results, plant peanuts when soil temperature are favorable for establishing good peanut stands.	Do not graze or feed treated hay or forage.
<b>PEANUTS</b> At Filling Time	Southern stem weevils and leafhoppers	12.5 lbs. per acre (20-inch row spacing) or 12 lbs. per 1000 ft. of row for any row spacing (minimum 24-inch spacing)	Distribute the granules to a band over the hilling area at pegging time. Work into the top few inches of soil immediately.	This treatment will also control the infestation of leafhoppers. Do not graze or feed treated hay or forage.
<b>POPPLES</b>	Aphids, leafhoppers, leaf miners, psyllids, whiteflies, and beet beetles larvae and reduction of the beetle adults	25 lbs. per acre (20-inch row spacing) or 28 lbs. per 1000 ft. of row for any row spacing (minimum 20-inch spacing) or 23 lbs. per 1000 ft. of row for heavy soil or clay loam soils per 1000 ft. of row for any row spacing (minimum 20-inch spacing)	Distribute the granules evenly in the furrow or granules may be banded on each side of the row at planting time.	With 60 days after treatment before harvesting potatoes.
<b>SORGHUM (SILC)</b> At Planting Time	Aphids (greenbug)	10 lbs. per acre (40-inch row) or 12 lbs. per 1000 ft. of row for any row spacing (minimum 20-inch spacing)	Place granules 2 inches to side and 1 inch below the seed at planting or as a 7-inch band over the row just prior to press wheat.	Do not place FAMBART 10-G in direct contact with the seed. Use restricted in Western states.
<b>SORGHUM (SILC)</b>	Aphids (greenbug)	10 lbs. per acre	Apply granules directly by air or ground equipment.	Do not apply within 30 days of harvest. Make only one post-emergence application. Use restricted in Western states.
<b>SUGAR BEETS</b>	Aphids, beet root maggot, leafhoppers, white, and leaf miners	10 lbs. per acre (20-inch row spacing) or 9 lbs. per 1000 ft. of row for any row spacing (minimum 20-inch spacing)	Application should be made by drilling or broadcast at planting time.	Do not place FAMBART 10-G in direct contact with the seed.  Do not apply within 30 days of harvest.
	Aphids, white	10 to 15 lbs. per acre	Apply to foliage when plants are dry. Use the higher rate on dense foliage for good coverage.  Foliar application by air may be used for rapid control when soil will not permit ground application.	Do not feed sugar beet tops or silage to dairy cattle.  Consult your state experimental station or state extension service for proper timing of application in your area.
<b>SUGAR BEET (Purple)</b>	Whiteflies	40 lbs. per acre	Apply in a 10 to 12 inch band directly on the food plants and surrounding soil in the open furrow immediately before covering.	
<b>WHEAT</b>	Greenbugs, aphids, Hessian fly	10 lbs. per acre (20-inch row spacing) or 12 lbs. per 1000 ft. of row for any row spacing (minimum 20-inch spacing)	Apply the granules to the seed furrow at planting time.	Do not feed or graze forage within 45 days of treatment. For aphid and greenbug control use only in Western and North Central states.

989