



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

JAN 25 1994

J. Allen Dunlap, III
WILLIAM M. MAHLBURG
Agent for: PLATTE CHEMICAL CO., INC. •
P. O. Box 667
Greeley, CO 80632

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Subject:

Label Amendment Submission of 09/16/93 in Response to PR Notice 93-7

EPA Reg. No. 34704-724

VALOR BRAND PRODUCTS DUSTING SULPHUR

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
 AND
- WITHIN one year from date of this acceptance.

PROPOSED LABEL

ACTIVE BACKSONENTS

VALOR BRAND PRODUCTS DUSTING SULFUR

Sui ERT MOREDIENTS: .. 2% KEEP OUT OF REACH OF CHILDREN CAUTION See Selow For Additional Procautionary Statements **EPA REG. NO. 34704-724** EPA EST NO. KG) 3447 PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION Hermful is swellowed, inheled or absorbed through skin. Avoid contact ith skin, eyes, or clothing. Avoid breathing dust, vapor. Applicators and other hannels imist west USER SAFETY RECOMMENDATIONS STATEMENTS OF PRACTICAL TREATMENT IF CWALLOWED: Induce vomiting by sticking finger down throat. He give anything by Mouth to an unconscious person. Get medical atten-

PHYSICAL HAZARD

Sulphur dust suspended in air ignites easily. Keep away from heat, sparts or flame. Do not emote white applying this product. MOTICE: Before using this product, read the entire Precautionary Statements, close and Statements, Directions for Use, Use Restrictions and Storage and Disposal Instructions on back panel and attached use directions.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a menner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

STORAGE AND DISPOSAL

STORAGE Keep away from tire and sparks. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or diffus material into food or drink containers. Do not contaminate other pseticides, tertifizers, water, food or feed by storage or disconse!

PESTICIDE DISPOSAL Do not conterminate water, food or feed by storage or disposed. Weeter resulting from the use of this product may be disposed of on site or at an approved weste disposed lacility.

CONTAINER DISPOSAL Completely empty bag into application

equipment. Then dispose of it in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of emole.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED: Sweep or should into containers for disposal or reworking. Keep dusting to a minimum, Refer to Proceedinary Statements.

GENERAL USE INFORMATION

USE PLATE DETERMINATION: Carefully read, understand, and tollow tabel use rotes and restrictions.

Under low disease conditions, minimum label rates per application can be used white maximum label rates and shortened spray schedules are resommended for severe or threatening disease conditions.

GROUND OR AIR APPLICATION: Dusters should be adjusted to give uniform and therough coverage. For letter details 22 to time, fructionary and rate of application colouist yelly Agricultural Extension Provide of Experimental Station authorities. When growing capes for a ecusions

SEST AVAILABLE COPY

k, or to drade where surface water is present or to

IF IM EYES: Flush eyes with large amounts of water for at les minutes. Conput a physician if britishon persists. IF ON SIGHT: Which affected skin areas with scap and water.

ENVIRONMENTAL HAZARDS

Do not apply when weather conditions lever drift from the areas tracted. Do not apply where run-off is fitsby to occur. Apply this product any so

led on this label. Do not conteminate mater when disposing of more westweeps. For terrestrial uses, do not apply directly to

DUSTING SULFUR EPA REG. NO. 34704-724

consult processor before applying sulphur. Almende, Cherries, Nesterines, Pesshes, Plume, Prunes: Brown Rot, Cherry Leef Spot (Cherries only), Prune Rust and Powdery Milder. Use 40 to 80 lbs. per acre in pre-bloom, bloom and postbloom periods. Apply use threatens during rain periods and repeat at 7 to 10 day ubon diese

Al'alle and Sugar Boots: Powdery Mildow and Rust. Use 30 to 50 lbs. per acre, beginning when disease first appears, and repeat at 10 to 14 days intervals through asseson.

Assertigue: Rust. Use 15 to 35 lbs. per acre. Dust after cutting stops, irrigate, cultivate, then apply the sulphur. Repeat at 7 to 10 day intervals through season.

ie and Peers: Scob and Powdery Milders. Use 40 to 50 lbs. per acre. Apply in pre-bloom, bloom and post-bloom periods. On pears do not apply after the fruit t agins to turn down.

ns and Peas: Powdery Mildew. Use 15 to 40 lbs. per acre. Apply when disease first appears and repeat as necessary.

Carrele and Onlone: Petrobis Mite. Use 30 to 35 to. per acre. Apply

when post first appears and repost as necessary.

Grapes: Powdery Milden. Use 8 to 20 lbs. per acre. Make first application when shoots are \$ in. to 8 in. long. Make second application when shoots are 12 in. to 15 in. long and third about 14 days later. Repeat application at 10 to 14 day intervals to protect new growth. Avoid over-application in hot weather. Concord and other Labruaca type grapes may be injured by autphur.

Strawberries, Respherries and Boysenberries: Powdery Mildew. Use 10 to 30 lbs. per acre. Apply when disease first appears and repeat as

Tomatees: Tomato Russet Mits. Use 30 to 50 lbs. per acre. Apply when post first appears and repost as necessary.

CONDITIONS OF SALE AND WARRANTY

PLATTE CHEMICAL CO. WARRANTS THAT THE PRODUCT CON-FORMS TO ITS CHEMICAL DESCRIPTION AND IS REASONABLY FIT FOR THE PURPOSE STATED ON THE LABEL ONLY WHEN USED IN ACCORDANCE WITH LABEL DIRECTIONS UNDER NORMAL CONCI-TIONS OF USE. PLATTE CHEMICAL CO. MAKES NO OTHER EX-PRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. HANDLING, STORAGE AND USE OF THE PRODUCT BY BUYER OR USER ARE BEYOND THE CONTROL OF PLATTE CHEMICAL CO. AND SELLER. RISKS SUCH AS CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES RESULTING FROM, BUT NOT LIMITED TO, WEATHER OR SOIL CONDITIONS, PRESENCE OF OTHER MATERI ALS, DISEASE, PESTS, DRIFT TO OTHER CROPS OR PROPERTY OR FAILURE TO FOLLOW LABEL DIRECTIONS WILL BE ASSUMED BY THE BUYER OR USER. IN NO CASE WILL PLATTE CHEMICAL CO. OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR IN-DIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.

BEST AVAILABLE COPY

The state of the s

PORMULATED FOR PLATTE CHEMICAL CO.

2