
34704-724

1





DUSTING SULFUR FUNGICIDE/MITICIDE EPA REG. NO. 34704-724

Agricultural Use Requirements cont'd.:

ł

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear.

STORAGE AND DISPOSAL

STORAGE: Keep away from fire and sparks. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Do not contaminate water, tood or feed by storage or disposal. Waster resulting from the use of this product maybe disposed of on site or at an approved waste disposal facility

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of it in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

STEPS TO BETAKEN IN CASE MATERIAL IS RELEASED OR SPILLED: Sweep or shovel into containers for disposal or reworking. Keep dusting to a minimum. Refer to Precautionary Statements.

GENERAL USE INFORMATION

USE RATE DETERMINATION: Carefully read, understand, and follow label use rates and restrictions.

Under low disease conditions, minimum label rates per application can be used while maximum label rates and shortened spray schedules are recommended for severe or threatening disease conditions.

GROUND OR AIR APPLICATION: Dusters should be adjusted to give uniform and thorough coverage. For further details as to time, frequency and rate of application consult your Agricultural Extension Service or Experimental Station authorities. When growing crops for processing consult processor before applying sulfur.

Almonds, Cherries, Nectarines, Peaches, Plums, Prunes: Brown Rot Biossom Blight, Cherry Leaf Spot (Cherries only), Prune Rust and Powdery Mildew. Use 40 to 50 lbs. per acre in pre-bloom, bloom and postbloom periods. Apply when disease threatens during rain pends and repeat at 7 to 10 day intervals.

Alfalfa and Sugar Beets: Powdery Mildew and Rust. Use 30 to 50 lbs. per acre, beginning when disease first appears, and repeat at 10 to 14 days intervals through season.

Asparagus: Rust. Use 15 to 35 lbs. per acre. Dust after cutting stops, irrigate, cultivate, then apply the sulfur Repeat at 7 to 10 day intervals through season.

Apples and Peers: Scab and Powdery Mildew. Use 40 to 50 lbs. per acre. Apply in pre-bloom, bloom and post-bloom periods. On peers do not apply after the fruit begins to turn down.

Beans and Peas: Powdery Mildew. Use 15 to 40 lbs, per acre. Apply when disease first appears and repeat as necessary.

Carrots and Onlone: Petrobia Mite, Use 30 to 35 lbs, per acre. Apply when pest first appears and repeat as necessary.

Grapes: Powdery Mildew, Use 8 to 20 lbs. per acre. Begin application when shoots are 6-8 inches long and repeat as necessary. Avoid over-application in hot weather, Concord and other Labrusca type grapes may be injured by sulfur.

Strawberries, Raspberries and Boysenberries: Powdery Mildew. Use 10 to 30 lbs. per acre. Apply when disease first appears and repeat as necessary.

Tomatoes: Tomato Russet Mite. Use 30 to 50 lbs, per acre. Apply when pest first appears and repeat as necessary.

NOTICE

IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATE, THE MANUFACTURER OR SELLER. IN NO CASE SHALL PLATE, THE MANUFACTURER OR SELLER BE LIABLE FOR CONSE-QUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE, THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTA- TIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

> FORMULATED FOR PLATTE CHEMICAL CO. 419 18TH STREET, GREELEY, COLORADO 80631-53C2

.

•