

SEP 16 1993

Mr. William M. Mahlburg
Platt Chemical Co.
P.O. Box 667
Greeley, CO 80632-0667

Dear Mr. Mahlburg:

Subject: Deletion of Crops
Casoron G-4 Herbicide
EPA Reg. No. 34704-738
Your submission dated Sept. 3, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

At the next printing of your label, or within 6 months, whichever comes first, please revise your label as follows:

1. Revise your "disposal of wastes statement" under the Environmental Hazards heading to read, "Do not contaminate water when disposing of equipment washwaters".

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR part 158, subpart K, § 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely yours,

BEST AVAILABLE COPY

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

CASORON® G-4

HERBICIDE

DICHLORBENIL WEED & GRASS KILLER

A SELECTIVE HERBICIDE FOR CONTROL OF CRABGRASS, FOXTAIL, SHEPHERDSPURSE, HORSETAIL, PIGWEED, QUACKGRASS, CANADA THISTLE, LEAFY SPURGE, FESCUE, AND CERTAIN OTHER ANNUAL AND PERENNIAL WEEDS.

ACTIVE INGREDIENT:

Dichlobenil® 4.0%

INERT INGREDIENTS: 96.0%

TOTAL 100.0%

*2,6-dichlorobenzonitrile

CASORON is a Registered Trademark of Duphar B.V., Weesp, the Netherlands

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 34704-738

EPA EST. NO. _____

30302

NET WEIGHT _____ LBS.

00F03

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or poison control center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If inhaled: Remove victim to fresh air.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Keep away from food, feedstuffs, and domestic water supplies. Keep container closed and out of reach of children and pets.

Pesticide Disposal: Pesticide that cannot be used according to instructions must be disposed of according to applicable Federal, state or local procedures.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

GENERAL DIRECTIONS

PERENNIAL WEED CONTROL

This product is recommended for control of certain perennial weeds such as ARTEMISIA, CANADA THISTLE, CURLY DOCK, FALSE DANDELION (CATSEAR), FESCUE, LEAFY SPURGE, ORCHARDGRASS, QUACKGRASS, RUSSIAN KNAWEED, TIMOTHY, WILD ARTICHOKE, WILD ASTER, WILD CARROT, AND YELLOW ROCKET IN ESTABLISHED WOODY ORNAMENTALS, FILBERTS AND FRUIT and for control in non-crop areas such as roadways, around buildings and fences, railroad and utility rights-of-way and installations.

SOIL SURFACE TREATMENT: Apply as a soil surface treatment in late fall from November 15 to February 15. Old weed growth should not be removed before application. Do not make surface application to areas which have been cultivated during the fall or summer prior to application.

INCORPORATED TREATMENT: Apply and incorporate immediately after application in late fall or very early spring before May 1.

ANNUAL WEED CONTROL

This herbicide is recommended for control of annual grassy and broad-leaved weeds and certain perennial weeds such as: ANNUAL BLUEGRASS (POA ANNUA), BLUEGRASS, BULL THISTLE, CAMPGRASS, CARPETWEED, CHICKWEED, CITRUS MELON, COFFEEWEED, CRABGRASS, CUDWEED, DANDELION, DOGFENNEL, EVENINGPRIMROSE, FIDDLENECK, FLORIDA PURSLANE (PURSLEY), FOXTAIL, GISEKIA, GROUNDSEL, HENBIT, HORSETAIL, JERUSALEM OAK, GOOSEFOOT, KNOTWEED, LAMBSQUARTERS, MAYPOPS, MILKWEED VINE (MORFENA ODORATA), MINERS LETUCE, NATALGRASS, OLD WITCHGRASS, PEPPERGRASS, PINFAPLEWEED, PLANTAIN, PURSLANE, RAGWEED, RED DEADNETTLE, REDROOT PIGWEED, ROSARYPEA, RUSSIAN THISTLE, SHEPHERDSPURSE, SMARTWEED, SPANISH NEEDLES, SPURGE, TEAWEED, TEXAS PANICUM (HURRAHGRASS), WILD BARLEY, WILD MUSTARD, WILD RADISH, AND YELLOW WOODSORREL.

Apply either in early spring before seeds or annual weeds germinate or after cultivation has removed the growing weeds. Distribute granules uniformly over the soil surface. If treatment is followed by application of 1/2 to 1 inch of water by overhead irrigation the lower rates are recommended. Shallow incorporation immediately after application is recommended in southern areas. In other areas, shallow incorporation or sprinking irrigation is suggested when applications are made during periods of high temperatures.

FRUIT AND NUT CROPS IN ORCHARDS AND NURSERIES

This product may be used on the following crops; Do not allow livestock to graze treated orchards:

A. Apple, Blueberry, Cherry, Grape, Nectarine, Peach, Pear, Prune Plum, and Filbert. May be used on bearing, non-bearing and nursery stock. Do not apply within 4 weeks after transplanting. Do not make application within one month of harvest of nuts or stone fruits. Apply 100 to 110 pounds per acre for annual weed control in areas actually tested. For control of perennial weeds apply 150 pounds per acre. Follow General Directions above.

B. Blackberry and Raspberry; for control of annual weeds follow General Directions above except use only on established plantings. Do not apply more than 100 pounds per acre. Do not apply during new shoot emergence.

BEST AVAILABLE COPY