

Reg # 62719-129
634704-746

PM-25

LABELING

1216



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

DEC 9 1993

Douglas M. Roby
DOWELANCO
QUAD IV 9002 PURDUE RD
Indianapolis, IN 46203

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 01/28/93 in Response to PR Notice 93-7
EPA Reg. No. 62719-129 → 34704-746
BALAN DRY FLOWABLE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 60% recycled fiber

2916

Submit the final printed labeling via the U.S. Postal Service to:


**Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001**

Hand or courier deliveries of final printed labeling may be made to:

**Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202**

BEST AVAILABLE COPY

Sincerely,


**Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)**

Attachment

3 of 16

DOWELANCO
EPA REG NO.: 62719-129 07/28/93
BALAN DRY FLOWABLE

Original Submission

Please add the Agricultural Use Requirements box to your main label that refers to the Agricultural Use Requirements box in your proposed supplemental labeling. Use the exact text and placement as shown on page 43 in Supplement Three to PR Notice 93-7.

6/10/93

4916
7

WPS Labeling

(Editor's Note: Deleted text indicated by strike-through and added text underlined)

(Base label, Balan DF, 12.5 lb):

(logo) DowElanco

**Balan*
Dry Flowable**

BEST AVAILABLE COPY

A selective herbicide for preemergence control of annual grasses and broadleaf weeds

Active ingredient:

N-butyl-N-ethyl-3,3,3-trifluoro-2,3-dihydro-2H-isoxaline . 60%

Inert ingredients . 40%

Contains 0.6 pounds of active ingredient per pound.

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

CAUTION

Harmful if Swallowed, Inhaled Or Absorbed Through The Skin

Avoid breathing dust or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. ~~Remove contaminated clothing and wash before reuse.~~

with COMMENTS
In EPA Letter Dated

DEC 9 1993

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

62719-129 →

Transfered to
34704-746

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Flush with plenty of water. Get medical attention if irritation develops.

If swallowed: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoons) with a large quantity of water. Do not give anything by mouth to an unconscious person. Immediately transport to a medical care facility.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

If inhaled: Remove individual to fresh air. If breathing difficulty occurs, get medical attention.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, or marshes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when conditions favor runoff or drift from the target area. Do not contaminate water when disposing of equipment washwaters.

BEST AVAILABLE COPY

6/10/93

5916

Refer to attached label booklet for additional precautionary information and Directions for Use including STORAGE AND DISPOSAL.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet. In case of an emergency endangering life or property involving this product, call collect 517-636-4400. Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 62719-129

ADE DL MM
EPA Est. 1471-IN-5; 37429-GA-01; 464-MI-1
Superscripts correspond to first letters of lot number.
900-000xxx

20108

*Trademark of DowElanco

20108 20108 20108 20108 20108 20108

20108

Net Weight 1.11 lb

BEST AVAILABLE COPY

BEST AVAILABLE COPY

Table of Contents	Page
Precautionary Statements	
Hazards to Humans and Domestic Animals	
First Aid	
Environmental Hazards	
Directions for Use	
Storage and Disposal	
General Information	
General Use Precautions	
Soil Texture Guide for Application Rates	
Soil Preparation	
Mixing Directions	
Dry Flowable and Liquid Fertilizers	
Dry Flowable and Liquid Fertilizers	
Dry Flowable and Liquid Fertilizers	
Application and Incorporation Directions	
Annual Grasses and Broadleaf Weeds Controlled by Balan Dry Flowable	
Approved Crops (no dots or page number after this heading)	
Alfalfa, Birdfoot Trefoil, Clover (Alsike, Ladino and Red)	
Lettuce (Direct Seeded)	
Warranty Disclaimer	
Inherent Risks of Use	
Limitation of Remedies	

BEST AVAILABLE COPY

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

CAUTION

PRECAUCION:

~~Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.~~

Harmful If Swallowed, Inhaled Or Absorbed Through The Skin

Avoid breathing dust or spray mist. Avoid contact with skin, eyes or clothing. Wash

~~Person using protective equipment (PPE)~~

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid

If in eyes: Flush with plenty of water. Get medical attention if irritation develops.

If swallowed: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoons) with a large quantity of water. Do not give anything by mouth to an unconscious person. Immediately transport to a medical care facility.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

If inhaled: Remove individual to fresh air. If breathing difficulty occurs, get medical attention.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, or marshes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when conditions favor runoff or drift from the target area. Do not contaminate water when disposing of equipment washwaters.

BEST AVAILABLE COPY

6/10/93

6

9716

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, safety, and labeling, and for farm and workplace assistance. It also contains specific instructions and instructions on the use of personal protective equipment (PPE) during the restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. Do not store in direct sunlight. Do not store at temperatures above 120°F. If stored above 120°F, Balan Dry Flowable herbicide could lose some of its dispersing properties, resulting in difficult application, poor weed control or crop injury. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Plastic Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY

General Information

Balan Dry Flowable (DF) herbicide is a preemergence herbicide for control of annual grasses and broadleaf weeds. Incorporation of Balan DF is required to assure effective season long weed control. Balan DF may be applied in liquid sprays of water or liquid fertilizer. Balan DF controls weeds as they germinate. Balan DF does not control established weeds.

General Use Precautions

When applied according to use directions and under normal growing conditions, Balan DF will not harm the treated crop. Over-application can result in crop injury, delayed emergence or soil residue. Uneven application or improper soil incorporation of Balan DF can result in erratic weed control or crop injury. Seedling disease, cold weather, deep planting, excessive moisture, high salt concentration or drought conditions and the possibility of damage to the crop may result. Under these conditions, delayed crop development or reduced yields may result.

In arid, irrigated areas of the Western United States - Arizona, California, Idaho, Montana, Nevada, Oregon, Utah, Washington and Wyoming - certain crops are susceptible to injury when planted in soil previously treated with Balan DF. To avoid such injury, do not plant wheat, barley, rye, other grasses, or onions within 10 months following an application of Balan DF. Do not plant milo (grain sorghum), corn, oats, spinach, red beets, sugar beets, or other root crops for 12 months following an application of Balan DF.

Balan DF should not be applied to soils that are wet or subject to prolonged periods of flooding as poor weed control may result.

Chemigation: Do not apply Balan DF through any type of irrigation system.

Soil Texture Guide for Application Rates

Balan DF application rates are based on soil texture. A fine textured clay soil will require a higher application rate than a coarse textured sandy soil. Application rates for Balan DF are determined by "soil texture groupings" (coarse, medium or fine) which correspond to the various field soil textures to be treated. Using the table below, determine if the field soil to be treated is coarse, medium or fine and use the application rate recommended for that soil texture grouping.

Soil Texture Class	Soil Classification
Coarse (light) soils	Sand, loamy sand, sandy loam
Medium soils	Loam, silty clay loam†, silt loam, silt, sandy clay loam†
Fine (heavy) soils	Clay, clay loam, silty clay loam†, silty clay, sandy clay, sandy clay loam†

†Silty clay loam and sandy clay loam soils are transitional soils and may be classified as either medium or fine textured soils. If silty clay loam or sandy clay loam soils are predominantly sand or silt, they are usually classified as medium textured soils. If they are predominantly clay, they are usually classified as fine textured soils.

Soil Preparation

Destroy existing weeds before application of Balan DF. Chop and thoroughly mix surface residues into the soil to a depth of 4 to 6 inches by plowing or disking. Use tillage equipment to break up large clods before application.

BEST AVAILABLE COPY

Mixing Directions

Balan Dry Flowable - Alone

Balan DF may be applied in water and liquid fertilizer materials. Prior to mixing Balan DF in liquid fertilizer, refer to the label section entitled "Testing for Compatibility in Liquid Fertilizers" for test procedures to determine compatibility with the fertilizer product to be used.

Equipment Requirements: Vigorous continuous agitation is required for Balan DF. Agitation should be sufficient to create a rippling or rolling action at the liquid surface. Centrifugal pumps are preferred and should have sufficient capacity to provide good bypass volume at all times. Screens placed on the suction side of the pump should be at least 16 mesh or coarser. Use 50 mesh or coarser screens between the pump and boom and when required at the nozzles. Before mixing, be sure equipment is clean and does not contain oil or residues from emulsifiable concentrates.

Start with a clean spray tank. Fill the sprayer to 1/3 of the required spray volume. Start agitation. Add the required amount of Balan DF in a controlled fashion to allow gradual wetting of the product and prevent clogging of screens and outlet ports. Continue agitation and fill spray tank to required spray volume. Maintain continuous agitation from mixing through application.

Precaution: Do not allow the mixture to back siphon into the water source.

Balan Dry Flowable in Tank Mix Combinations

Balan DF may be tank mixed with other products when applied with water or liquid fertilizer materials. Prior to mixing Balan DF and another product in liquid fertilizer, refer to the tank mix product manufacturer's label to determine if application in liquid fertilizer is recommended. Also refer to the section in this label entitled "Testing for Compatibility in Liquid Fertilizers" for testing procedures to determine tank mix compatibility with the liquid fertilizer product to be used. Read and follow all label instructions for each material to be added to the spray tank.

Vigorous continuous agitation is required for all tank mixes. Sparger pipe agitators generally provide the best agitation in spray tanks. To prevent foaming in the spray tank during filling, keep the end of the fill pipe below the surface of the water. This prevents air from being stirred or splashed into the mixture.

Mixing Order: Add Balan DF to spray tank according to mixing directions for "Balan DF - Alone", on page-2. Fill the spray tank to 90% of the final spray volume. Add different tank mix formulation types in the following order: dry flowables (DF); wettable powders (WP); aqueous suspensions (AS), flowables (F), liquids (L), solutions (S), and emulsifiable concentrates (EC, LC). Allow time for complete mixing and dispersion after addition of each product. Allow extra mixing and dispersion time for dry flowable products. Continue agitation and fill tank to final spray volume.

Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be resuspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled materials may be more difficult to resuspend than when originally mixed.

Premixing: When tank mixing, initial mixing and dispersion of certain dry flowable or wettable powder products may be improved by premixing with water (slurrying). Where recommended, follow product label instructions for each material. Adding the slurried material to the spray tank through a 20 to 35 mesh wetting screen will help assure good initial dispersion.

Line screens in the tank should be no finer than 50 mesh (100 mesh is finer than 50 mesh).

BEST AVAILABLE COPY

Testing for Compatibility in Liquid Fertilizers

Balan DF alone or in combination with other products may be mixed with liquid fertilizer. Vigorous continuous agitation is always required when Balan DF is applied in liquid fertilizer. In some cases, Balan DF or tank mixes containing Balan DF may not combine properly with some liquid fertilizer materials. Small quantities of such mixtures should always be tested before full-scale mixing. Use the procedure provided below to determine if a compatibility agent is needed or which available agent performs best with the mixture being applied. The compatibility agents listed at the end of this section have been thoroughly tested. Other surfactant materials available may not be suitable for use with liquid fertilizers. Individual state regulations relating to liquid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale. The combination of Balan DF with solution and suspension-type fertilizers provides annual weed control equal to Balan DF applied in water.

PROCEDURE

1. Add 1 pint of liquid fertilizer to a 1 quart glass jar.
2. Add equal amounts (1 to 4 teaspoonfuls depending on application rate) of Balan DF and other products to be tank mixed in the liquid fertilizer. Add materials in the order recommended in the "Mixing Directions" section of this label. Mix until evenly dispersed after addition of each formulation. If dry flowable or wettable powder formulations do not disperse well, it may be necessary to slurry the materials in a small amount of water before addition to the liquid fertilizer.
3. Observe the jar for about 10 minutes. If materials rise to the surface and form a thick layer (oil curds) which will not disperse again when agitated, a compatibility agent is needed. If the mixture is easily dispersed to its original state with slight agitation, no compatibility agent is needed; however, good agitation must be provided to maintain dispersion in the spray tank from mixing through application.
4. If the need for a compatibility agent is demonstrated in step 3, it may be tested as follows: Using a clean clear plastic or glass container, start at step 1, above. Add 1/2 teaspoonful of the compatibility agent to the liquid fertilizer, mix well, and then repeat steps 2 and 3.

An effective compatibility agent will cause the mixture to remain uniformly mixed with little or no separation or oil rising to the surface for 1/2 hour or longer. If slight separation occurs, 2 to 3 inversions of container should provide a uniform mixture. If oily curds form which do not disperse, try adding additional compatibility agent or test the effectiveness of an alternative compatibility agent using the procedure above. A clean jar should be used for each test.

Compatibility Agents

The phosphate ester-type surfactants listed below are designed for use with liquid fertilizers and can be mixed at rates as low as 1 1/2 to 2 pints per ton of liquid fertilizer. They usually do not work well as compatibility agents for tank mixes in plain water. Add the compatibility agent just before adding herbicides. Read and follow label directions for the compatibility agent.

1. Sponto 168D (Wilco Chemicals Co., Chicago, IL)
2. Compat (Farm Chemicals, Inc., Aberdeen, NC)
Not for use in California.
3. Unite (Hopkins Ag Chemical, Madison, WI)
4. T-Mulz 734-2 (Thompson-Hayward Chemical Co., Kansas City, KS)
Not for use in California.
5. Amoco Spray Mate (Amoco Oil Co., Chicago, IL)
Not for use in California.

BEST AVAILABLE COPY

Application and Incorporation Directions

Application

Apply Balan DF anytime after January 1 when soil can be worked. Apply in 5 to 40 gallons of water or liquid fertilizer per acre (broadcast basis). For band application, apply proportionately less. Use any properly calibrated low pressure spray that will apply the spray uniformly. As the amount of spray volume decreases, the importance of accurate calibration and uniform application increases. Avoid boom overlaps that will increase rates above those recommended. Check the spray daily to insure proper calibration and uniform application.

Incorporation

Before planting Balan DF must be incorporated one time within 4 to 8 hours after application. A second incorporation is not recommended. If a second incorporation is necessary, it should be applied to a different area. If Balan DF is applied to a wet, firm soil surface, or if the wind velocity is consistently greater than 10 mph, variable weed control may result. Variable weed control may also result if incorporation is delayed more than 8 hours in the eastern United States or 4 hours in the western United States. Operate equipment to mix Balan DF into the top 2 to 3 inches of the final seedbed. Generally, incorporation equipment will mix Balan DF approximately half as deep as equipment is operated. For example, operating equipment 4 inches deep will mix Balan DF into approximately the upper 2 inches of the seedbed.

Special Note: In the western United States, extremely high temperatures and intense sunlight may be present at the time of application. Under such conditions, Balan DF must be incorporated within 4 hours after application to prevent loss of herbicidal activity.

Incorporation Equipment

Use equipment that thoroughly mixes Balan DF into the soil. Shallow incorporation less than 2 inches deep in the final seedbed may result in erratic weed control. Use of equipment not listed below may result in poor or erratic weed control and/or crop injury. The following types of equipment are recommended:

- Tandem disc: Set equipment to cut 4 to 6 inches and operate at 4 to 6 mph. Two incorporation passes in different directions are required. A single incorporation pass with a tandem or double disc does not provide adequate soil mixing.
- P.T.O. driven equipment (tillers, cultivators, hoes): Set equipment to operate 4 to 6 inches deep and operate no more than 4 mph. Only one incorporation pass is necessary with this type of equipment.

Precaution: Use of P.T.O. equipment alone for incorporation in coarse (sandy) soils may result in an uneven distribution of Balan DF. This occurs when soil containing Balan DF is deposited into wheel tracks that are deeper than the operating depth of the tiller. This results in an excessive amount of Balan DF in tractor wheel depressions and poor weed control in intervening areas containing less than the intended rate of Balan DF.

Cultivation after Planting

Soil treated with Balan DF may be shallowly cultivated or rotary hoed without loss of weed control activity. Do not cultivate deeper than the Balan DF treated layer or untreated soil may be exposed resulting in loss of weed control. In bedded culture, if weeds germinate in the bottom of the furrow due to the removal of treated soil, these weeds should be controlled by cultivation.

BEST AVAILABLE COPY

14 7 16

6/10/93

Annual Grasses and Broadleaf Weeds Controlled by Balan Dry Flowable

Annual Grasses

<u>Common Name</u>	<u>Scientific Name</u>
annual bluegrass	<i>Poa annua</i>
barnyardgrass (watergrass)	<i>Echinochloa crus-galli</i>
broadleaf signalgrass	<i>Brachiaria platyphylla</i>
(spreading panicgrass, See special use instruction on pages 5 and 6)	
crabgrass	<i>Digitaria spp.</i>
(large crabgrass)	
(smooth crabgrass)	
crowfootgrass	<i>Dactyloctenium aegyptium</i>
foxtails	<i>Setaria spp.</i>
(giant foxtail)	
(green foxtail)	
(pigeongrass)	
(robust white foxtail)	
(yellow foxtail)	
Johnsongrass (seedling only)	<i>Sorghum halepense</i>
jungerice	<i>Echinochloa colonum</i>
panicum, fall	<i>Panicum dichotomiflorum</i>
(See special use instruction on pages 5 and 6)	
panicum, Texas	<i>Panicum texanum</i>
(buffalograss)	
(Coloradograss)	
ryegrass, annual (Italian)	<i>Lolium multiflorum</i>
sandbur, field	<i>Cenchrus incertus</i>

Annual Broadleaf Weeds

<u>Common Name</u>	<u>Scientific Name</u>
carpetweed	<i>Mullugo verticillata</i>
chickweed	<i>Stellaria media</i>
Florida pusley	<i>Richardia scabra</i>
(Florida purslane)	
(Mexican clover)	
(pusley)	
knotweed	<i>Polygonum aviculare</i>
lambquarters, common	<i>Chenopodium album</i>
pigweeds	<i>Amaranthus spp.</i>
redroot pigweed	
(carelessweed)	
(rough pigweed)	
prostrate pigweed	
spiny amaranth	
purslane, common	<i>Portulaca oleracea</i>
redmaids	<i>Calandrinia caulescens</i>

BEST AVAILABLE COPY

Approved Crops**Alfalfa, Birdsfoot Trefoil, Clover (Alsike, Ladino and Red)**

Apply and incorporate Balan DF before seeding.

Broadcast Application Rates per Acre**Soil Texture**Coarse
Medium
Fine**Balan Dry Flowable
(Pounds/Acre)**2
2
2 1/2**Alfalfa (Birdfoot Clover)**

Apply and incorporate before seeding.

Broadcast Application Rates per Acre**Soil Texture**Coarse
Medium
Fine**Balan Dry Flowable
(Pounds/Acre)**2
2
2 1/2**Warranty Disclaimer**

DowElanco warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. DOWELANCO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of DowElanco or the seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at DowElanco's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

DowElanco shall not be liable for losses or damages resulting from handling or use of this product unless DowElanco is promptly notified of such loss or damage in writing. In no case shall DowElanco be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of DowElanco or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Copyright© 1993 DowElanco

BEST AVAILABLE COPY

16 9 16
7

6/10/93

13

Label text is based on EPA accepted copy dated 02/08/93
Revised 06/09/93

Amendments:

- 1) Added labels statements required by the Worker Protections Standard (PR Notice 93-7).

BEST AVAILABLE COPY