

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

April 26, 2024

Brien O'Loughlin Regulatory Consultant Loveland Products, Inc. c/o Pyxis Regulatory Consulting Inc. 4110 136th St CT NW Gig Harbor, WA 98332

Subject: Label Amendment - Registration Review Mitigation for Oxadiazon

Product Name: Napropamide-oxadiazon 4-2 granules

EPA Registration Number: 34704-771

Application Date: 08/18/2022 Decision Number: 586935

Dear Brien O'Loughlin:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Oxadiazon Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable. As part of the registration review, EPA identified the need for this product to be classified as Restricted Use. The labeling approved today incorporates the necessary changes for classification as Restricted Use, therefore the classification is effective on the date of this letter.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved

Page 2 of 2 EPA Reg. No. 34704-771 Decision No. 586935

registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. Due to the classification of this product as Restricted Use, in accordance with 40 CFR 152.167, after 120 days, any of this product you sell or distribute must:

- 1) Bear an approved amended label containing the terms of the Restricted Use classification;
- 2) Bear a sticker containing the product name, EPA registration number, and any terms of restricted use; **or**
- 3) Be accompanied by supplemental labeling bearing the product name, EPA registration numbers, and any terms of restricted use.

After 270 days, any of this product you sell or distribute must bear the approved amended labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Kelsi Grogan by phone at (202)566-2228, or via email at grogan.kelsi@epa.gov.

Sincerely,

Jun 1 Postello

Kevin Costello, Branch Chief Risk Management and Implementation Branch 2 Pesticide Re-Evaluation Division Office of Pesticide Programs

ENCLOSURE: Stamped label

RESTRICTED USE PESTICIDE

Due to potential human health risks of concern (cancer) from chronic dietary or dermal exposure when used in accordance with label directions.

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

NAPROPAMIDE	GROUP	15	HERBICIDE
OXADIAZON	GROUP	14	HERBICIDE

[Optional Loveland Products, Inc Company or Brand Logo]

Napropamide-Oxadiazon 4-2 Granules

ACTIVE INGREDIENTS:

Napropamide		4.0%
Oxadiazon: 2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)-z-1,3,4 oxadiazolin-5-one		2.0%
INERT INGREDIENTS:		
	TOTAL	100.0%

Selective, Pre-emergent Granular Herbicide for Control of Annual Grasses and Broadleaf Weeds in Container, Field Grown Nursery Stock, Liner Stock, Ground Covers, Shade/Lath Houses, and Landscape Ornamentals[.]

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or	Take off contaminated clothing.
clothing:	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	Call a poison control center or doctor for treatment advice.
If Inhaled:	Move person to fresh air.
	• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth -to-mouth, if possible.
	Call a poison control center or doctor for further treatment advice.

1

ACCEPTED

04/26/2024

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565

For Chemical Emergency:

Spill, Leak, Fire, Exposure, or Accident,

Call CHEMTREC Day or Night

Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

[For Product Use Information CALL: 1-888-LPI-CUST (1-888-574-2878)]

EPA Reg. No.: 34704-771 EPA Est. No.: _____ Net Weight:

Reg Rev EXP ID 08/22 re 08/23 re 10/23 re 02/24

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt and long pants, socks, and shoes with waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Waterproof gloves,
- And shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use a closed system or enclosed cabs in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Apply only as specified on this label.

Runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

[Hazard Statement for Fish and Aquatic Invertebrates]

This pesticide is toxic to fish and aquatic invertebrates, including oysters and shrimp.

Groundwater Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of napropamide and oxadiazon from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is expected to occur within 48 hours.

NON-TARGET ORGANISM ADVISORY

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize drift.

[Covering or Collecting Spilled Granules]

Spilled napropamide granules exposed on soil surface may be hazardous to wildlife. Cover or collect granules spilled during loading.

DIRECTIONS FOR USE RESTRICTED USE PESTICIDE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

Do not apply by air.

After application, user must thoroughly irrigate the site as soon as possible on the same day of application.

WEED RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, please note that Napropamide-Oxadiazon 4-2 Granules contains both a Group 14 and a Group 15 herbicide. Any weed population may contain plants naturally resistant to Group 14 and/or Group 15 herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of Napropamide-Oxadiazon 4-2 Granule or other Group 14 and Group 15 herbicide within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical

information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to these MOAs have been found in your region. Do not assume that each listed weed is being controlled by multiple mechanisms of action. Co-formulated active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredients in this product.
- For further information or to report suspected resistance contact Loveland Products, Inc. retailer, representative or call 1-888-LPI-CUST [(574-2878)]. You can also contact your pesticide distributor or university extension specialist to report resistance.
- Plant into weed-free fields and keep fields as weed-free as possible.
- Fields with difficult to control weeds should be rotated to crops that allow the use of herbicides with alternative mechanisms of action or different management practices.
- To the extent possible do not allow weed escapes to produce seeds, roots or tubers. Manage weed seeds at harvest and post-harvest to prevent a buildup of the weed seedbank.
- Prevent an influx of weeds into the field by managing field borders.
- Apply this herbicide at the correct timing and rate needed to control the most difficult weed in the field.
- Use a broad-spectrum soil-applied herbicide with a mechanism of action that differs from this product as a foundation in a weed-control program. Do not use more than two applications of this or any other herbicide with the same mechanism of action within a single growing season unless mixed with an herbicide with another mechanism of action with an overlapping spectrum for the difficult-to-control weeds.
- If resistance is suspected, treat weed escapes with an herbicide with a different MOA or use non-chemical methods to remove escapes.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls,
- Shoes plus socks,
- Waterproof gloves,
- And protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protections Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets out of treated areas until dusts have settled.

GENERAL INFORMATION

This product is a selective, pre-emergent herbicide for control of many annual grasses and broadleaf weeds on a wide variety of woody ornamental shrubs, vines, trees, and conifer nurseries. May not be applied to ornamentals in residential properties. This product controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

Apply with a properly calibrated granular applicator that will apply the granules uniformly (uniform application is essential for satisfactory weed control). Calibrate the applicator prior to use, according to the manufacturer's directions, and check frequently during the application to be certain that the equipment is working properly.

Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic purposes.

This product can be applied to *newly planted* container stock after soil has settled from first watering, field-grown nursery stock, dichondra, and to established plants.

This product is recommended on several different container potting soil mixtures. These mixtures include those containing 30% soil and/or sand and mixtures containing no soil (e.g., perlite, peat, bark, sawdust, and sand).

This product does not control established weeds. It must be applied to freshly weeded soil before weeds germinate or during fall and winter months.

After application, user must thoroughly irrigate the site as soon as possible on the same day as application.

For all use sites:

- The minimum retreatment interval (RTI) between applications is 60 days.
- Total applications of oxadiazon of all physical forms must not exceed 6.0 lbs. a.i./A/year if this product is used on the same site with liquid formulations.

For ornamental landscaping, trees, shrubs, and nurseries:

- The maximum single application rate is 150 lbs. of Napropamide-Oxadiazon 4-2 Granule product/Acre. (Equivalent to 6.0 lbs. of Napropamide a.i. / A and 3.0 lbs. of Oxadiazon a.i./A).
- The maximum annual application rate is 225 lbs. of Napropamide Oxadiazon 4-2 granule/ Acre/ Year. (Equivalent to 9.0 lbs. of Napropamide a.i./ A/yr and 4.5 lbs. of Oxadiazon a.i./A/yr).

GENERAL PRECAUTIONS

Read all instructions before using. Use only for recommended purposes and at recommended rates. Do not overdose. Treated plants may not be used for food or feed for one year after the last application of this product.

Not for use on home lawns.

HOW TO USE

Woody Ornamental Shrubs, Vines, Trees, and Conifer Nursery Application*

For maximum weed control, apply prior to weed seed germination. This product can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. After Application, user must thoroughly irrigate the site as soon as possible on the same day of application. This product must not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of this product. *Do not apply in greenhouses* as plant injury may result from codistillation of this product's active ingredients onto leaf surfaces in condensation. *Do not apply to wet foliage*. Under normal conditions, expect control for 90-120 days.

RTI – Minimum Retreatment Interval is 60 days, however, control of weeds may last up to 3 months. Consult with your local extension service or certified crop advisor for specific retreatment interval recommendations. May not be applied to ornamentals in residential properties.

This product may be used on the following established ornamentals and ground covers.

Agapanthus	Ivy, Algerian
Ajuga	Ivy, English
Arborvitae	Jasmine, orange
Ash, green	Jasmine, primrose
Azalea**	Jasmine, star
Bay, or Bay-rum tree	Jasmine, showy
Birch, paper	Juniper
Black-laurel	Lantana, trailing or Weeping lantana
Boxwood, common	Larch, European
Boxwood, Japanese	Leucothoe, coast
Bougainvillea	Leucothoe, drooping
Camellia, common	Lily-of-the-Nile
Camelia, Japanese	Magnolia, southern
Camellia, sasanqua	Maple
Cedar, atlas	Myrtle, wax
Cedar, red	Nandina or Sacred bamboo
Citrus, ornamental (lime, tangelo, temple, Cuban	Oak
shaddock)	Oleander including Dwarf oleander
Cotoneaster**	Osmanthus
Crape myrtle**	Palm (bamboo, fishtail, licuala Macarthur, manac,
Croton	maniala, parlor, rattan, sealing wax, sugar)
Cypress	Palmetto, wax
Cypress, Leyland	Photinia, fraser
Dogwood, common	Pine including Spruce Pine or Cedar Pine
Dogwood, kousa	Pittosporum, Japanese
Dogwood, red osier	Plum, natal
Douglas fir	Podocarpus, yew
Eucalyptus	Poplar
Euonymous, spreading	Pyracantha

^{*}Conifer Nursery Application is Not a Registered Use in California.

Euonymous, wintercreeper	Raphiolepis
Forsythia	Redbud, eastern
Gardenia	Rose
Gazania trailing	Rhododendron**
Heather, Scotch	Sedum
Hemlock, eastern**	Shower-of-gold
Hibiscus, Chinese or Hawaiian hibiscus (crooked Celia,	Spruce**
Jane Cowell crooked Rose, hulagirl)**	Trailing African daisy
Holly	Viburnum including Sandankwa**
Honeysuckle	Xylosma
Ice plant	Yew
Ixora	

^{**}Varietal Sensitive, Refer to Sensitive Varieties Section for Additional Information

Sensitive Varieties

Under certain growing conditions, the plant varieties listed below may be sensitive to this product. Growers should evaluate usage on a small scale prior to large scale applications.

Plant Species	Varieties
Andromeda, Japanese	Snowdrift, Temple bell
(Pieris japonica)	
Azalea	Pink champagne, Hershey,
(Rhododendron spp.)	Leuchtenfuer,
,,,,	Memoire John Hearrens, Hino-crimson,
	White grandeur, Everest, Girard Roberta
Cotoneaster	Eichholz, Himalayan, Shangri-la
(Cotoneaster spp.)	
Crape Myrtle, common	Petite red imp, Seminole bush
(Lagerstroemia indica)	
Heath, spring	Mediterranean pink
(Erica carnea)	
Hemlock, Eastern	pendula
(Tsuga canadensis)	
Hibiscus	Red heart
(Hibiscus syriacus)	
Rhododendron	Daphnoides, Nova zembla,
(Rhododendron spp.)	Catawba album
Spruce, Norway	Pumila
(Picea abies)	
Spruce, white	Densata
(Picea glauca)	

Viburnum	Davidii
(Viburnum spp.)	

Users who wish to apply this product on ornamental plant species not recommended on this label may determine the suitability for such uses by making a trial application at a recommended rate to a small number of plants. Prior to using this product on a large number of plants, the treated plants should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to deter- mine if the treatment is non-injurious to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from the use of this product on plant species not recommended on this label.

Amount to Use

Ground Application - Apply broadcast or banded to ornamentals at 100 to 150 pounds of product per broadcast acre (or 2.25 to 3.44 pounds per 1000 square feet). Use the higher rate for harder-to-control weeds and the lower rate for easier-to-control weeds. Remove existing weed/grass growth before application. In a single application, do not apply more than 3.44 pounds per 1000 square feet.

The maximum single application rate is 150 lbs. of Napropamide-Oxadiazon 4-2 Granule product/Acre. (Equivalent to 6.0 lbs. of Napropamide a.i. / A and 3.0 lbs. of Oxadiazon a.i./A).

The maximum annual application rate is 225 lbs. of Napropamide Oxadiazon 4-2 granule/ Acre/ Year. (Equivalent to 9.0 lbs of Napropamide a.i./ A/yr and 4.5 lbs. of Oxadiazon a.i./A/yr).

Total applications of Oxadiazon of all physical forms must not exceed 6.0 lbs a.i./A/yr if this product is used on the same site with liquid formulations.

Annual Broadleaf Weeds		
Bittercress	Liverwort	
Bristly oxtongue	Niruri	
Carpetweed	Pennsylvania smartweed	
Coast fiddleneck	Petty spurge	
Common chickweed	Pineapple-weed	
Common groundsel	Prickly lettuce	
Common mallow (Cheeseweed)	Prostrate knotweed	
Common purslane	Prostrate spurge	
Common ragweed	Redroot pigweed	
Eveningprimrose	Sensitiveplant, (Patridgepea)	
Filaree	Shepherdspurse	
Fireweed	Sowthistle	
Gallnsoga	Speedwell	
Garden spurge	Spiny amaranth	
Golden ragwort	Spotted catsear	
Horse purslane	Swinecress	
Lambsquarters	Yellow woodsorrel	
Little mallow		

Annual Grasses

Annual bluegrass
Annual sedge
Barnyardgrass
Bristly foxtail

Cheatgrass (Bromus tectorum) Ripgut brome
Downy brome (Bromus tectorum) Sandbur

Fall panicum Smooth crabgrass

Giant foxtail Soft brome (Bromus mollis)
Goosegrass Southwestern cupgrass

Green foxtail Stinkgrass
Guineagrass Texas panicum
Italian ryegrass Wild barley
Johnsongrass, seeding Wild oats
Large crabgrass Witchgrass
Red spangletop Yellow foxtail

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in its original bag, in a cool, dry, locked place out of the reach of children. Keep container closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration. Do not burn, unless allowed by state and local ordinances.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reason- ably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT

NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

FORMULATED FOR

[Optional Graphic: Loveland Products, Inc Company or Brand Logo]

LOVELAND PRODUCTS, INC.

P.O. BOX 1286, GREELEY, COLORADO 80632-1286