



PM 23 34704-777 10/23/97 PS 1/86

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PESTICIDES AND TOXIC
SUBSTANCES

Ms. Jeannie K. Smith
Platte Chemical Company
150 South Main Street
Fremont, NE 68025

OCT 23 1997

Dear Ms. Smith:

Subject: Crabgrass Preventer with Barricade® .20% Plus
EPA Registration No. 34704-781
Crabgrass Preventer with Barricade® .22% Plus
EPA Registration No. 34704-780
Crabgrass Preventer with Barricade® .25% Plus
EPA Registration No. 34704-779
Crabgrass Preventer with Barricade® .295% Plus
EPA Registration No. 34704-778
Crabgrass Preventer with Barricade® .325% Plus
EPA Registration No. 34704-777
Crabgrass Preventer with Barricade® .375% Plus
EPA Registration No. 34704-776
Applications and Letters Dated July 10, 1997,
Requests To Update Labeling of Products To Match
Barricade® F Label; and to Revise Size of Retail Containers

Your request to amend the labeling of the subject pesticide product registrations under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, to reflect the subject proposed amendments has been reviewed and is acceptable under the Act, provided that you:

1. Correct the statement on the front panel that reads: "Barricade® is a registered trademark of Sandoz, Inc. to read: "Barricade® is a registered trademark of Novartis Crop Protection, Inc.".
2. Submit one (1) copy of your final printed labeling for each of these products before you release it for shipment under the proposed label, reflecting the above comment.

If these conditions are not complied with, these registrations

2/6

will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of each of these products constitutes acceptance of these conditions.

A stamped copy of each of the labels is enclosed for your records.

Sincerely yours,

for Eugene M. Wilson

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosures

376

ACCEPTED
with COMMENTS
In EPA Letter Dated

OCT 23 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
34704-777

CRABGRASS PREVENTER WITH BARRICADE® .325% PLUS

ACTIVE INGREDIENT:

Prodiamine, (N³, N³-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine] 0.325

INERT INGREDIENTS: 99.675
TOTAL: 100.0%

See Fertilizer Tag for Guaranteed Fertilizer Analysis
BARRICADE® is a registered trademark of Sandoz, Ltd.

For selective preemergence control of grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings)
- covers up to x,xxx square feet

KEEP OUT OF REACH OF CHILDREN

CAUTION

See the following section,
Precautionary Statements,
Hazards To Humans and Domestic Animals,
for additional information.

EPA Reg. No. 34704-777
EPA Est. No.
Net Weight:

IHT EXP07/97

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

STATEMENTS OF PRACTICAL TREATMENT

- If in eyes:** Flush thoroughly with water for several minutes. Get medical attention if irritation persists.
- If on skin:** Wash with soap and water. Rinse thoroughly. Get medical attention if irritation persists.
- If inhaled:** Remove victim to fresh air.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of soil for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to soil. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
STORAGE: Store in original container away from other fertilizer, food, or food stuffs and separated from other pesticides.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Paper or plastic containers: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or, at an incineration facility, or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- established perennials & wildflower plantings

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application.


**CRABGRASS PREVENTER WITH
BARRICADE® .325% PLUS**
EPA REG. NO. 34704-777

476

- May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 160 lbs/A. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.
- Do not apply more than 460 lbs of this product per calendar year.
- Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates Table.

See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES



Contact your UHS Sales Representative or local Agricultural Extension Agent for crabgrass germination information for your area.

Use Precautions

The following precautions apply to the use of this product in turf grasses and lawns:

Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.

Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as diking may result in reduced weed control.

Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

† These are the maximum rates per calendar year by species limitations.

Maximum Application Rate/Calendar Year of Crabgrass Preventer with BARRICADE .325% Plus by Turf Grass Species†

Turf Species	lbs product/A	lbs a/A
Bermudagrass	460	1.5
Bahlegrass	308	1.0
Centipodegrass	230	0.75
Seashore Paspalum	200	0.65
St. Augustinegrass	308	1.0
Tall Fescue (including turf-type)	308	1.0
Zoysia	308	1.0
Buffalograss	308	1.0
Kentucky Bluegrass	308	1.0
Perennial Ryegrass	308	1.0
Creeping Red Fescue	230	0.75
Creeping Bentgrass	200	0.65

Spreader Settings

Spreader	Spreader Setting*			
	Rates Lb/A			
460	308	230	200	
Spreader and setting to be determined for a particular fertilizer blend.				

* The above spreader settings are approximate. Condition of your spreader, turf conditions and walking speed with spreader can affect feeding rate. Walk at normal rate. Check your spreader at the rate of application you wish to use and adjust accordingly.

Weeds Controlled

When used in accordance with this label this product will provide control of the following weeds:

- | | |
|--|---------------------------------|
| Barnyardgrass | Kochia |
| Bluegrass, Annual (<i>Poa annua</i>) | Lambsquarter, Common |
| Carpetweed | Lovegrass |
| Chickweed, Common | Panicum (Texas, Fall, Browntop) |
| Chickweed, Mouseear (from seed) | Pigweed |
| Crabgrass (large, smooth) | Purslane, Common |
| Crowfootgrass | Pusley, Florida |
| Cupgrass, Woolly | Rescuegrass† |
| Foxtails, Annual | Shepherd's Purse† |
| Goosegrass† | Signalgrass, Broadleaf |
| Honbit | Speedwell, Persian |
| Itchgrass | Sprangletop |
| Johnsongrass (from seed) | Spurge, Prostrate |
| Junglerice | Witchgrass |
| Knotweed | Woodsorrel, Yellow (from seed) |

- 1 In many areas a single application of 200 to 460 lbs/A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 200 to 308 lbs/A followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate Table above.
- 2 Applications for this weed should be made in late summer, fall or winter prior to germination.
- 3 Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

**CRABGRASS PREVENTER WITH
BARRICADE® .325% PLUS
EPA REG. NO. 34704-777**

Crabgrass Preventer with BARRICADE .325% Plus Rate (lb/A)		Months Before Overseeding		
Product	Rate	North	Transition	South
155	.50	4	4	4
200	.65	5	4	4
230	.75	6	5	5
245	.80	—	6	6
308	1.00	—	7	7
350	1.14	—	—	9
400	1.30	—	—	10
460	1.50	—	—	12

LANDSCAPE ORNAMENTALS

(including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

Use Rates

Apply this product at 200 to 460 lbs/A in fall and/or spring. Use higher rates of application for longer control periods. Sequential or single applications are allowed so long as the total amount of product applied does not exceed the maximum annual application rate of 460 lbs/A/yr.

Application Timing and Information

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate postemergence herbicide.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

Newly Transplanted Landscape Ornamentals

This product may be applied to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific Name	Common Name
<i>Abies</i> spp.	Fir species** (Balsam, Fraser, Noble, etc.)
<i>Abelia grandiflora</i>	Abelia; Sherwood
<i>Acer palmatum</i>	Japanese Maple
<i>A. platanoides</i>	Norway Maple
<i>Achillea</i> , sp.	Yarrow; King Edward
<i>Actinidia chinensis</i>	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Agapanthus orientalis</i>	
<i>Akebia quintata</i>	Five-Leaf or Chocolate Vine
<i>Allium cernuum</i>	Lady's Leek, Nodding Onion
<i>Anemone hybrida</i>	Japanese Anemone
<i>Aquilegia</i> sp.	Aquilegia; Red and Gold
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Artemisia</i> sp.	Wormwood; Silver Mound, Castle
<i>Aster</i> sp.	Aster; Bonny Blue, Purple Dome
<i>Aster X Frikartii</i>	
<i>Athrium Filix-lemnia</i>	Lady Fern; Fern Lady
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Begonia</i> sp.	Fibrous Begonia; Hardy Grandis

Scientific Name

- Berberis gladiwomensis*
- B. julianae*
- B. mentorensis*
- B. thunbergii*
- B. verruculosa*
- Bergenia cordifolia*
- Boltonia asteroides*
- Bougainvillea* spp.
- Buddleia davidii*
- Buxus microphylla*
- Callistemon viminalis*
- Calluna vulgaris*
- Campanula carpatica*
- Campis X Tagliabuana*
- Carpobrotus edulis*
- Cassia artemiscoides*
- Ceanothus rigidus*
- Ceratostigma plumbagonoides*
- Chamaecyparis pisifera*
- Chrysanthemum nipponicum*
- Cleyera japonica*
- Citrus* spp.
- Coreopsis* sp.
- Cornus florida*
- C. stolonifera*
- Cortaderia selloana*
- Cotoneaster apiculatus*
- C. buxifolius*
- C. dammeri*
- C. microphyllus*
- Crataegus* spp.
- Crocosmia* sp.
- Cupressus sempervirens*
- Delosperma alba*
- Delosperma* sp.
- Delphinium* sp.
- Dianthus deltoides*
- D. gratianopolitanus*
- Dodonea viscosa*
- Echinacea purpurea*
- Elaeagnus pungens*
- Euonymus fortunei*
- E. japonica*
- E. katuschovica*
- Fatsia japonica*
- Forsythia intermedia*
- Forsythia suspensa*
- F. viridissima*
- Gaillardia* sp.
- Gardenia jasminoides*
- Gaura* sp.
- Gentiana dahurica*
- Geranium cinereum*
- Gladiolus* spp.
- Gypsophila repens*
- Hedera helix*
- Hellanthemum* sp.
- Homocallis* sp.
- Hibiscus*
- Hibiscus* sp.
- Hibiscus rosa-sinensis*
- Hosta plantaginosa*
- H. Sieboldiana*
- Houttuynia cordata* var. *variegata*
- Hydrangea macrophylla*
- Ilex cornuta*
- I. crenata*
- I. opaca*
- I. peryi*
- I. vomitoria*
- Inula ensifolia*
- Iris ensata*
- Iris siberica*
- Iris* spp.
- Jasminium nudiflorum*
- Juniperus chinensis*

Common Name

- Barberry
- Wintergreen Barberry
- Menlor Barberry
- Japanese Barberry
- Warty Barberry
- Snowbank
- Bougainvillea
- Butterfly-Bush (Dwarf Blue); Royal Red
- Japanese Boxwood
- Weeping Bottlebrush
- Scotch Heather
- Tussock Bellflower, (White Clips)
- Trumpet Creeper, Trumpet Flower, Madame Galen
- Hottentot Fig (Ice Plant)
- Feathery Cassia
- Wild Lilac
- False Cypress
- Cleyera
- Citrus species*
- Coreopsis (Callopsis): Early Sunrise, Moonbeam
- Flowering Dogwood
- American Dogwood
- Pampas Grass
- Cranberry Cotoneaster
- Cotoneaster
- Bearberry Cotoneaster
- Rockspray Cotoneaster
- Hawthorne
- Lucifer
- Italian Cypress
- White Trailing Ice Plant
- Cooperi Pink
- Larkspur; Blue Elf
- Dianthus; Maiden Pinks "Zing"
- Cheddar Pink
- Hop Bush
- Coneflower, Purple; Magnus
- Silverberry
- Wintercreeper
- Japanese Spindle Tree (Evergreen Euonymus)
- Spreading Euonymus
- Japanese Aralia
- Border Forsythia
- Weeping Forsythia
- Greenstem Forsythia
- Gaillardia, Blanket Flower; "Goblin"
- Gardenia, Cape-Jasmine
- Gentian
- Cranesbill
- Gladiolus species**
- Baby's Breath
- English Ivy
- Sunrose
- Daylily; Aztec Gold, Stella De Oro, Tender Love
- Rose of Sharon**
- Mallow; Disco Bello White
- Chinese Hibiscus
- Hosta, Plantain Lily (Fragrant)
- Hosta, "Giant Hosta"
- Bigleaf Hydrangea
- Chinese Holly
- Japanese Holly; Helleri
- American Holly
- Holly
- Yaupon Holly; Schillings
- Sword-Leaved Iris; Jodlesong
- Siberian Iris; Cabernet
- Iris species**
- Winter Jasmine
- Chinese Juniper; Nick's
- Compact. Parsonii

586

*Do not use on food producing trees, vines, or plants.

**Not for use on container grown plants.

**CRABGRASS PREVENTER WITH
BARRICADE® .325% PLUS
EPA REG. NO. 34704-777**

696

Scientific Name	Common Name
<i>J. conferta</i>	Shore Juniper; Blue Pacific
<i>J. davurica</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans</i> sp.	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagrostromia indica</i>	Crape Myrtle
<i>Lagerstromia indica X Fauriei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavandea</i> sp.	Lavender; Munstead
<i>Leontopodium alpinum</i>	Edelweiss
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>Ligustrum amurense</i>	Amur Privet
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lilium</i> sp.	Lily; Jazz
<i>Liriope muscari</i>	Big Blue Lily; Turf; Silver Mound;
	Evergreen Giant
<i>Liriope Muscari var. variegata</i>	Liriope, Variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Loropetalum chinense</i>	Burgundy
<i>Lythrum</i> sp.	Loosestrife; Modern pink
<i>Magnolia</i> spp.	Magnolia species**
<i>Malephora luteola</i>	Ice Plant
<i>Malus</i> sp.	Crabapple*
<i>Miscanthus sinensis</i>	Yaka Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus</i> spp.	Narcissus species**
<i>Nerium oleander</i>	Oleander
<i>Olea europaea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osmanthus heterophyllus</i>	Osmanthus (False Holly); Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum amoereum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fourtan Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado*
<i>Photinia fraseri</i>	Fraser's Photinia (Redtip)
<i>Physotogia Virginiana</i>	Dragonhead, False; Vivid
<i>Picea</i> spp.	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-Valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. elliotti</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine
<i>P. palustris</i>	Longleaf Pine
<i>P. radiata</i>	Monterey Pine
<i>P. thunbergiana</i>	Japanese Black Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>P. taeda</i>	Loblolly Pine
<i>P. virginiana</i>	Virginia Pine
<i>Pistachio</i> sp.	Pistachio*
<i>Pittosporum trichophyllum</i>	Queensland Pittosporum
<i>P. tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus</i> sp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>P. fortuneana</i>	Firethorn
<i>P. koidzumii</i>	Firethorn
<i>Pyrus</i> sp.	Bradford Pear sp.
<i>Quercus shumardii</i>	Oak, Shumard's Red
<i>Quercus</i> spp.	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Raphiolepis umbellata</i>	Yedda Hawthorn
<i>Rhododendron</i>	'Coral Bells'
(including Azalea)	'Delaware Valley White'
	'Flame Creeper'
	'Formosa'
	'Glard Crimson'
	'George L. Tabor'

Scientific Name

Common Name

<i>Rosa banksia</i>	'Hino-crimson'
<i>Rosmarinus officinalis</i>	'PJM'
<i>Rudbeckia</i> sp.	'Roseum Elegans'
<i>Santolina virens</i>	'Wakelebisu'
<i>Saxifraga</i> sp.	'White Gumbo'
<i>Scabiosa</i> sp.	Lady Bank's Rose
<i>Seum album</i>	Rosemary
<i>Seum caucicola</i>	Black-Eyed Susan; Goldstrum
<i>Sedum dasyphyllum</i>	Santolina
<i>Sedum spurium</i>	Saxifrage; Purple Dome
<i>Spiraea Bumalda</i>	Pincushion Flower
<i>Syzygium paniculatum</i>	Stoncrop
<i>Taxus cuspidata</i>	Stoncrop; Lidakense
<i>T. media</i>	Stoncrop
<i>Toucrum</i> sp.	Stoncrop
<i>Thalictrum dipterocarpum</i>	Stoncrop; Dragon's Blood
<i>Thuja occidentalis</i>	Spirea: Anthony Waterer
<i>Trachelospermum asiaticum</i>	
<i>Tsuga canadensis</i>	
<i>Tulipa</i> spp.	Japanese Yew
<i>Veronicac</i> sp.	Yew
	Germander
<i>Viburnum japonicum</i>	Meadow Rue
<i>V. odoratissimum</i>	American Arborvitae
<i>V. plicatum</i>	Star Jasmine
<i>V. rigidum</i>	Canada Hemlock
<i>V. suspensum</i>	Tulip species
<i>V. tinus</i>	Veronica, Speedwell; Sunny
<i>V. trilobium</i>	Border
<i>V. wrightii</i>	Japanese Viburnum
<i>Vinca major</i>	Sweet Viburnum
<i>Vinca minor</i>	Japanese Snowball
<i>Vitis</i> sp.	Canary Island Viburnum
<i>Weigela florida</i>	Arrowood Viburnum
<i>Yucca aloifolia</i>	Laurustinus
<i>Y. filamentosa</i>	Cranberry Bush
	Leatherleaf Viburnum
	Vinca
	Periwinkle
	Grape*
	Old Fashioned Welgeia
	Spanish Bayonet
	Adam's Neeckle

*Do not use on food producing trees, vines, or plants.

**Not for use on container grown plants.

NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. PLANT INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER. EXCEPT AS EXPRESSLY DESCRIBED ABOVE, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

FORMULATED FOR
PLATTE CHEMICAL CO.

150 SO. MAIN STREET, FREMONT, NEBRASKA 68025-5697