



3/5/2002
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

1/6

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Ms. Glenda Haage
Registration Manager
Platte Chemical Company
418 18th St.
Greeley, CO 80631

MAR 5 2002

Dear Ms. Haage:

Subject: Crabgrass Preventer with Barricade® .375% Plus
EPA Registration Number 34704-776
Crabgrass Preventer with Barricade® .295% Plus
EPA Registration Number 34704-778
Crabgrass Preventer with Barricade® .20% Plus
EPA Registration Number 34704-781
Crabgrass Preventer with Barricade® 0.50% Plus
EPA Registration Number 34704-820
Applications and Letters Dated February 19, 2002,
Request To Amend Registrations To: "Respond to
PR Notice 2001-1, First Aid Statements Required

The proposed labeling amendments have been reviewed and found acceptable for the subject pesticide product registrations under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, provided that you revise the label of each these products as described below:

1. Revise the statement that reads: "Barricade® is a registered trademark of Novartis Crop Protection, Inc." to read: "Barricade® is a registered trademark of Syngenta Crop Protection, Inc."
2. Revise the "Notice", regarding conditions of sales and warranty which clearly states remedies if these products cause any damage when used as directed or in their handling, by adding a statement that instructs the using public to return the product after reading the labeling of the product. A copy of the Uniform Commercial Codes that are applicable to the remedies claimed are enclosed with an example of an acceptable warranty with limitation of remedies.

Submit one (1) copy of the final printed label prior to your shipment of each of these pesticide products under its respective revised stamped labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment under this labeling constitutes acceptance of these conditions. Stamped copies of these labels are enclosed for your records.

Sincerely yours,

Eugene W. Wilson
for Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosures (4 stamped labels
3 pages re: UCC Codes)

CRABGRASS PREVENTER WITH BARRICADE® .20% PLUS

ACTIVE INGREDIENT:

Prodiamine, [N³, N³-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine]0.20

INERT INGREDIENTS:99.80
TOTAL: 100.0%

See Fertilizer Tag for Guaranteed Fertilizer Analysis
BARRICADE® is a registered trademark of Syngenta.

For selective preemergence control of grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings)
- covers up to x,xxx square feet

KEEP OUT OF REACH OF CHILDREN CAUTION

See the following section, Precautionary Statements, Hazards To Humans and Domestic Animals, for additional information.

EPA REG. NO. 34704-781

EPA EST. NO.

NET WEIGHT:

IHT

EXP 01/02

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 -20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:
1-800-228-5635, EXT. 136, OR CALL COLLECT, 612-851-8180, EXT. 136.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Paper or plastic containers: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or, at an incineration facility, or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- established perennials & wildflower plantings

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application.

**ACCEPTED
with COMMENTS
In EPA Letter Dated**

MAR 5 2002


**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

34704-781

CRABGRASS PREVENTER WITH BARRICADE® .20% PLUS

EPA REG. NO. 34704-781

See the map below for approximate crabgrass seed germination dates.



Contact your UHS Sales Representative or local Agricultural Extension Agent for crabgrass germination information for your area.

Use Precautions

The following precautions apply to the use of this product in turf grasses and lawns:

Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.

Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table.

1) These are the maximum rates per calendar year by species limitations.

Turf Species:	Maximum Application Rate/Calendar Year of Crabgrass Preventer with BARRICADE .20% Plus by Turf Grass Species ¹⁾	
	lbs/A (lbs/1000 sq.ft)	lbs ai/A
Bermudagrass ²⁾	750 (17.2)	1.5
Bahiagrass		
Centipedegrass		
Seashore Paspalum		
St. Augustinegrass		
Tall Fescue (including turf-type)		
Zoysia		
Buffalograss	500 (11.5)	1.0
Kentucky Bluegrass		
Perennial Ryegrass		
Creeping Red Fescue	370 (8.5)	0.75
Creeping Bentgrass	330 (7.6)	0.65

2) May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 260 lbs/A. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

- Do not apply more than 750 lbs of this product per calendar year.
- Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates Table.

Spreader Settings

Spreader	Spreader Setting [*]			
	Rates lbs/A (lbs/1000 sq.ft.)			
	750 (17.2)	500 (11.5)	370 (8.5)	330 (7.6)
Spreaders and setting to be determined for a particular fertilizer blend.				

* The above spreader settings are approximate. Condition of your spreader, turf conditions and walking speed with spreader can affect feeding rate. Walk at normal rate. Check your spreader at the rate of application you wish to use and adjust accordingly.

Weeds Controlled

When used in accordance with this label this product will provide control of the following weeds:

Barneygrass	Kochia
Bluegrass, Annual (<i>Poa annua</i>)	Lambsquarter, Common
Carpetweed	Lovegrass
Chickweed, Common	Panicum (Texas, Fall, Browntop)
Chickweed, Mouseear (from seed)	Pigweed
Crabgrass (large, smooth)	Purslane, Common
Crowfootgrass	Pusley, Florida
Cupgrass, Woolly	Rescuegrass ³⁾
Foxtails, Annual	Shepherd's Purse ²⁾
Goosegrass ¹⁾	Signalgrass, Broadleaf
Henbit	Speedwell, Persian
Itchgrass	Sprangletop
Johnsongrass (from seed)	Spurge, Prostrate
Junglerice	Witchgrass
Knotweed	Woodsorrel, Yellow (from seed)

1 In many areas a single application of 330 to 750 lbs/A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 330 to 500 lbs/A followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate Table above.

2 Applications for this weed should be made in late summer, fall or winter prior to germination.

3 Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

Crabgrass Preventer with BARRICADE .20% Plus Rate (lb/A)	Product	ai	Months Before Overseeding		
			North	Transition	South
250		.50	4	4	4
330		.65	5	4	4
370		.75	6	5	5
400		.80	—	6	6
500		1.00	—	7	7
570		1.14	—	—	7
650		1.30	—	—	10
750		1.50	—	—	12

LANDSCAPE ORNAMENTALS

(including Established Perennials and Wildflower Plantings)

This product may be applied for residual pre-emergence weed control in ornamentals.

Use Rates

Apply this product at 330 to 750 lbs/A in fall and/or spring. Use higher rates of application for longer control periods. Sequential or single applications are allowed so long as the total amount of product applied does not exceed the maximum annual application rate of 750 lbs/A/yr.

**CRABGRASS PREVENTER WITH
BARRICADE® .20% PLUS
EPA REG. NO. 34704-781**

5/6

Application Timing and Information

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate postemergence herbicide.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

Newly Transplanted Landscape Ornamentals

This product may be applied to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grfts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific Name

Abies spp.

Abelia grandiflora
Acer palmatum
A. platanoides
Achillea sp.
Actinidia chinensis
Agapanthus africanus
Agapanthus orientalis
Akebia quinata
Allium cernuum
Anemone hybrida
Aquilegia sp.
Arctostaphylos densiflora
Arctotheca calendula
Artemisia sp.
Aster sp.
Aster X Frikartii
Athnum Filix-femina
Aucuba japonica
Begonia sp.
Berberis gladwynensis
B. julianae
B. mentorensis
B. thunbergii
B. verruculosa
Bergenia cordifolia
Boltonia asteroides
Bougainvillea spp.
Buddleia davidii

Buxus microphylla
Callisternon viminalis
Calluna vulgaris
Campanula carpatica
Campis X Tagliabuana

Carpobrotus edulis
Cassia artemisoides
Ceanothus rigidus
Ceratostigma plumbagonoides
Chamaecyparis pisifera
Chrysanthemum nipponicum
Cleyera japonica
Citrus spp.
Coreopsis sp.

Cornus flonda
C. stolonifera
Cortaderia selioana
Cotoneaster apiculatus

Common Name

Fir species**
(Balsam, Fraser, Noble, etc.)
Abelia. Sherwood
Japanese Maple
Norway Maple
Yarrow. King Edward
Kiwi*
Lily-of-the-Nile (African Lily)

Five-Leaf or Chocolate Vine
Lady's Leek, Nodding Onion
Japanese Anemone
Aquilegia: Red and Gold
Vine Hill Manzanita
Cape Weed
Wormwood; Silver Mound, Castle
Aster: Bonny Blue, Purple Dome

Lady Fern; Fern Lady
Japanese Aucuba
Fibrous Begonia: Hardy Grandis
Barberry
Wintergreen Barberry
Mentor Barberry
Japanese Barberry
Warty Barberry

Snowbank
Bougainvillea
Butterfly-Bush (Dwarf Blue);
Royal Red
Japanese Boxwood
Weeping Bottlebrush
Scotch Heather
Tussock Bellflower, (White Clips)
Trumpet Creeper, Trumpet
Flower, Madame Galen
Hottentot Fig (Ice Plant)
Feathery Cassia
Wild Lilac

False Cypress

Cleyera
Citrus species*
Coreopsis (Calliopsis): Early
Sunrise, Moonbeam
Flowering Dogwood
American Dogwood
Pampas Grass
Cranberry Cotoneaster

Scientific Name

C. buxifolius
C. dammeri
C. microphyllus
Cretaegus spp.
Crocsmia sp.
Cupressus sempervirens
Deliosperma alba
Deliosperma sp.
Delphinium sp.
Dianthus deltooides
D. gratianopolitanus
Dodonea viscosa
Echinacea pupurea
Elaeagnus pungens
Euonymus fortunei
E. japonica

E. kiautschovica
Fatsia japonica
Forsythia intermedia
Forsythia suspensa
F. viridissima
Gaillardia sp.
Gardenia jasminoides
Gaura sp.
Gentiana dahurica
Geranium cinereum
Gladiolus spp.
Gypsophila repens
Hedera helix
Helianthemum sp.
Hemerocallis sp.

Hibiscus
Hibiscus sp.
Hibiscus rosa-sinensis
Hosta plantaginea
H. Sieboldiana
Houttuynia cordata var. *variegata*
Hydrangea macrophylla
Ilex cornuta
I. crenata
I. opaca
I. pernyi
I. vomitoria
Inula ensifolia
Iris ensata
Iris siberica
Iris spp.
Jasminium nudiflorum
Juniperus chinensis

J. conferta
J. davurica
J. horizontalis
Juglans sp.
Justicia brandegeana
Lagerstromia indica
Lagerstomia indica X Fauriei
Lantana montevidensis
Lavender sp.
Leontopodium alpinum
Ligustrum sinense
Ligustrum amurense
L. japonicum
L. lucidum
Lilium sp.
Liriope muscari

Liriope Muscari var. *variegata*
Liriope spicata
Lobelia Cardinalis
Lonicera japonica
L. tatarica
Loropetalum chinense
Lythrum sp.
Magnolia spp.
Maleophora luteola
Malus sp.
Miscanthus sinensis
Nandina domestica
Narcissus spp.
Nerium oleander
Olea europaea
Ophiopogon japonicus
Osmanthus heterophyllus
Osteospermum fruticosum
Oxydendrum arboreum

Common Name

Cotoneaster
Bearberry Cotoneaster
Rockspray Cotoneaster
Hawthorne
Lucifer
Italian Cypress
White Trailing Ice Plant
Cooperi Pink
Larkspur: Blue Elf
Dianthus: Maiden Pinks "Zing"
Cheddar Pink
Hop Bush
Coneflower. Purple; Magnus
Silverberry
Wintercreeper
Japanese Spindle Tree
(Evergreen Euonymus)
Spreading Euonymus
Japanese Aralia
Border Forsythia
Weeping Forsythia
Greenstem Forsythia
Gaillardia, Blanket Flower: "Goblin"
Gardenia, Cape-Jasmine

Gentian
Cranesbill
Gladiolus species**
Baby's Breath
English Ivy
Sunrose
Daylily; Aztec Gold, Stella De
Oro, Tender Love
Rose of Sharon**
Mallow: Disco Belle White
Chinese Hibiscus
Hosta, Plantain Lily (Fragrant)
Hosta, "Searsucker"

Bigleaf Hydrangea
Chinese Holly
Japanese Holly; Helleri
American Holly
Holly
Yaupon Holly; Schillings

Sword-Leaved Iris; Jodlesong
Siberian Iris; Cabernet
Iris species**
Winter Jasmine
Chinese Juniper; Nick's
Compact. Parsonii
Shore Juniper; Blue Pacific
Parsoni
Creeping Juniper
Walnut*
Shrimp Plant
Crape Myrtle
Crape Myrtle, Tuscarora
Weeping Lantana
Lavender; Munstead
Edelweiss
Chinese Privet; Variegata
Amur Privet
Japanese Privet
Glossy Privet (Waxleaf)
Lily: Jazz
Big Blue Lily; Turf; Silver Mound;
Evergreen Giant
Liriope, Variegated
Liriope, Creeping
Cardinal Flower, Indian Pink
Japanese Honeysuckle
Tatarian Honeysuckle
Burgundy
Loosestrife; Modern pink
Magnolia species**
Ice Plant
Crabapple*
Yaka Jima Silverleder
Heavenly Ban.boo
Narcissus species**
Oleander*
Olive*
Mondo Grass
Osmanthus (False Holly); Gull Tide
Trailing African Daisy
Sourwood

*Do not use on food producing trees, vines, or plants.
**Not for use on container grown plants.

**CRABGRASS PREVENTER WITH
BARRICADE® .20% PLUS
EPA REG. NO. 34704-781**

6/6
*Do not use on food producing trees, vines, or plants
**Not for use on container grown plants.

NOTICE

IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION. ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. THE MANUFACTURER OR SELLER. IN NO CASE SHALL PLATTE, THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER. EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE, THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

Scientific Name

Paeonia suffruticosa
Pennisetum setaceum
Perovskia atriplicifolia
Persea americana
Photinia fraseri
Physotegia Virginiana
Picea spp.

Pieris japonica
Pinus brutia
P. cananensis
P. eliottii
P. halepensis
P. nigra
P. palustris
P. radiata
P. thunbergiana
P. strobus
P. sylvestris
P. taeda
P. virginiana
Pistachio sp.
Pittosporum rhombifolium
P. tobira
Podocarpus macrophyllus
Prunus laurocerasus
Prunus sp.

Pseudotsuga menziesii
Pyracantha coccinea
P. fortuneana
P. koidzumii
Pyrus sp.
Quercus shumardii
Quercus spp.
Raphiolepis indica
Raphiolepis umbellata
Rhododendron
(including Azalea)

Rosa banksiae
Rosmarinus officinalis
Rudbeckia sp.
Santolina virens
Saxifraga sp.
Scabiosa sp.
Sedum album
Seum caudicola
Sedum dasycyphillum
Sedum spurium
Spiraea Bumalda
Syzygium paniculatum
Taxus cuspidata
T. media
Teucrium sp.
Thalictrum dipterocarpum
Tuja occidentalis
Trachelospermum asiaticum
Tsuga canadensis
Tulipa spp.
Veronica sp.

Viburnum japonicum
V. odoratissimum
V. plicatum
V. rigidum
V. suspensum
V. tinus
V. trilobium
V. wrightii
Vinca major
Vinca minor
Vitis sp.
Weigela florida
Yucca aloifolia
Y. filamentosa

Common Name

Tree Peony
Fountain Grass (Dwart)

Avocado*
Fraser's Photinia (Redtip)
Dragonhead, False, Vivid
Spruce species** (Colorado Blue,
Norway, etc.)
Lily-of-the-Valley Bush
Calabrian Pine
Canary Island Pine
Slash Pine
Aleppo Pine
Austrian Pine
Longleaf Pine
Monterey Pine
Japanese Black Pine
Eastern White Pine
Scotch Pine
Loblolly Pine
Virginia Pine
Pistachio*
Queensland Pittosporum
Japanese Pittosporum
Japanese Yew
English Laurel
Almond, Apricot, Nectarine,
Peach, Plum and Prune*
Douglas Fir**
Firethorn (Scarlet)
Firethorn
Firethorn
Bradford Pear sp.
Oak, Shumard's Red
Oak species
Indian Hawthorne
Yedda Hawthorn
'Coral Bells'
'Delaware Valley White'
'Flame Creeper'
'Formosa'
'Girard Crimson'
'George L. Tabor'
'Hino-crimson'
'PJM'
'Roseum Elegans'
'Wakeiebisu'
'White Gumbo'
Lady Bank's Rose
Rosemary
Black-Eyed Susan; Goldstrum
Santolina
Saxifrage; Purple Dome
Pincushion Flower
Stonecrop
Stonecrop; Lidakense
Stonecrop
Stonecrop; Dragon's Blood
Spiraea; Anthony Waterer

Japanese Yew
Yew
Germander
Meadow Rue
American Arborvitae
Star Jasmine
Canada Hemlock
Tulip species
Veronica, Speedwell; Sunny
Border
Japanese Viburnum
Sweet Viburnum
Japanese Snowball
Canary island Viburnum
Arrowood Viburnum
Laurustinus
Cranberry Bush
Leatherleaf Viburnum
Vinca
Periwinkle
Grape*
Old Fashioned Weigela
Spanish Bayonet
Adam's Needle

**FORMULATED FOR
PLATTE CHEMICAL CO.
419 18TH STREET, GREELEY, COLORADO 80631-5852**