



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
34704-792

Date of Issuance:
JAN 28 1997

NOTICE OF PESTICIDE:
 Registration
 Reregistration

Term of Issuance:
Conditional

(under FIFRA, as amended)

Name of Pesticide Product:
Clean Crop
Trifluralin HF

Name and Address of Registrant (include ZIP Code):

Platte Chemical Company
150 South Main Street
Fremont, NE 68025-0667

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes before you release the product for shipment:

- a. Revise the EPA Registration Number to read, "EPA Reg. No. 34704-792".
- b. Revise the oral and inhalation statements of practical treatment as on the Treflan H.F.P (EPA Reg. No. 62719-250) label:

"IF SWALLOWED: Do not induce vomiting. Call a physician or Poison Control Center. If available, administer activated charcoal (6-8 heaping teaspoonfuls) with a large quantity of water. Do not induce vomiting or give anything by mouth to an unconscious person. Immediately transport to a medical care facility and see a physician.

Signature of Approving Official:

Date:
JAN 28 1997

IF INHALED: Remove individual to fresh air. Get medical attention if breathing difficulty occurs. If not breathing, give artificial respiration, preferably cardiopulmonary resuscitation assistance, and get medical attention immediately."

- c. Add the following "Note To Physician" as on the Treflan H.F.P (EPA Reg. No. 62719-250) label:

"This product contains an aromatic hydrocarbon and can be extremely harmful if swallowed. Aspiration of this product may produce a severe pneumonitis. Stomach lavage with a cuffed endotracheal tube in place and immediate administration of activated charcoal, 6 to 8 heaping teaspoonfuls with water, should be considered. Treatment is otherwise symptomatic and supportive."

- d. Delete the directions for use on "Mint". The label for the technical trifluralin source product does not list mint among the use sites for which end-use products may be formulated.
- e. Delete the second paragraph under "CROP RECOMMENDATIONS" regarding plantback restrictions for vegetable crops other than those listed on the label.
- f. Under "SPECIAL PRECAUTIONS" on the first page, delete the heading "In Florida Only" and move the paragraph under the heading so that it appears as the third paragraph under "SPECIAL PRECAUTIONS".
- g. Replace "Trifluralin 4EC" with "Trifluralin HF" in the following locations on the label:

- First paragraph under "GENERAL CHEMIGATION" on page 4.
- In the formula for calculating the amount of Trifluralin HF to be impregnated on a ton of dry bulk fertilizer on page 6.
- In the directions for use on FLAX on page 8.
- In the directions for use on RAPESEED on page 9.
- In the rate table for Trifluralin HF pre-plant followed by Sencor or Lexone as an overlay treatment for weed control in soybeans on page 11.
- In the rate table for the tank mix of Trifluralin HF and Far-Go on page 12.
- In the directions for use on Winter Wheat (Partial control or suppression of annual Brome Species) on page 13. "Trifluralin 4EC" appears four times in this section.

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

On April 30, 1996, the Agency issued a Reregistration Eligibility Decision (RED) for the active ingredient, trifluralin. This product registration relies on acute toxicology and product chemistry data for a trifluralin product

3/116

page 3
EPA Reg. No. 34704-792

(Trifluralin H.F.P.; EPA Reg. No. 62719-250) which is undergoing reregistration. Depending on the outcome of the Agency's review of these data, you may be required to submit additional data and/or revise the product label to support the continued registration of this product.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

RD:STANTON:PM Team 23:Rm. 235:CM-2:305-5218:Disk #4a:34704TOE.REG

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	S. Stanton							
DATE ▶	Jan 24, 1997							

ACCEPTED
with COMMENTS
In EPA Letter Dated
JAN 28 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
34704-792



TRIFLURALIN HF

A selective herbicide for the pre-emergence control
of annual grasses and broadleaf weeds.

ACTIVE INGREDIENT:	
Trifluralin (p,p',p'-trifluoro-2,6-dinitro-N,N-dipropyl- p-toluidine)	43.0%
INERT INGREDIENTS**:	57.0%
TOTAL	100.0%

Contains 4 pounds active ingredient per gallon.

*Contains Petroleum Distillates

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See Below For Additional Precautions And Directions For Use.

EPA REG. NO. 34704

EPA EST. NO. 34704-MS-1

NET CONTENTS: 2 1/2 GALS. (9.46 L)

5902

EXP12P98

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING

Causes Substantial But Temporary Eye Injury • Harmful If Swallowed, Inhaled, Or Absorbed Through The Skin • May Cause Skin Sensitization Reactions In Certain Individuals.

Avoid breathing vapors or spray mist and contact with skin, eyes, or clothing.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

Long-sleeved shirt and long pants, chemical-resistant gloves, such as barrier laminate or viton, shoes plus socks, and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwater.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention if irritation persists.

If on skin: Wash contaminated skin with soap and water. Get medical attention.

If inhaled: Move to clear atmosphere. Restore breathing if necessary. Get medical attention.

If swallowed: Do not induce vomiting and get medical attention immediately. Ingestion is toxic and irritating to the stomach. Vomiting may cause aspiration into the lungs resulting in pulmonary edema which may be fatal. Gastric lavage may be indicated.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat and open flame.

SPECIAL PRECAUTIONS

Applied according to directions and under normal growing conditions, TRIFLURALIN HF will not harm the treated crop. Over application may result in crop injury or a soil residue.

Uneven application or improper soil incorporation can result in erratic weed control or crop injury. Seedling disease, cold weather, deep planting, excessive moisture, high salt concentration or drought may weaken crop seedlings and increase the possibility of damage. Under these conditions, delayed crop development or reduced yields may result.

In the Western United States—Arizona, Colorado, California, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington and Wyoming, and sugar beet growing areas of Minnesota and North Dakota: To avoid crop injury in arid areas, do not plant sugar beets, red beets or spinach for 12 months after a TRIFLURALIN HF spring application or for 14 months after a TRIFLURALIN HF fall application. Plow the land to a depth of 12 inches prior to planting sugar beets to prevent the possibility of crop injury. To avoid crop injury do not plant sorghum (milo), corn or oats for 14 months after a spring application or for 18 months after a fall application. If land has not been irrigated, do not plant any of these crops for 18 months after a spring application or 20 months after a fall application.

In the Western United States—Those portions of Kansas, Nebraska, North Dakota, Oklahoma, South Dakota and Texas where at least 25 inches of irrigation and/or rainfall (total) was used to produce the crop:

Do not plant sorghum or oats for 12 months after a TRIFLURALIN HF application. If less than 25 inches of total water was used to produce the crop, do not plant sorghum or oats for 18 months after application. Cool wet weather conditions during the early stage of growth may increase the possibility of injury to sorghum.

In the Eastern United States:

Moldboard plow before planting sugar beets where a TRIFLURALIN HF spring application was made the previous season. Also note planting restrictions listed in the section on control of rhizome johnsongrass.

In Florida Only:

To avoid crop injury do not plant vegetable crops other than those listed on the label within 5 months following the application of TRIFLURALIN HF.

In North Dakota Only:

Green foxtail resistance to Trifluralin and related herbicides in the spring cereal grain production areas: Certain populations of Green foxtail (Pigeongrass) in North Dakota have been identified as resistant to dinitroaniline type herbicides such as trifluralin. Because CLEAN CROP TRIFLURALIN HF will not control these resistant Green foxtail, alternative types of herbicides that are not dinitroanilines should be used. The grower assumes all risk of nonperformance due to dinitroaniline resistance if CLEAN CROP TRIFLURALIN HF is used to control Green foxtail in affected spring cereal grain crops.

In Texas Only:

Do not use in Pecos or Reeves Counties

WEEDS AND GRASSES CONTROLLED

TRIFLURALIN HF will not control established weeds.

GRASSES CONTROLLED

Annual bluegrass	(<i>Poa annua</i>)
Barnyardgrass	(<i>Echinochloa sp.</i>)
(Watergrass)	
Brachiaria	(<i>Brachiaria sp.</i>)
(Signalgrass)	
Bromegrass	(<i>Bromus tectorum</i>)
(Cheatgrass)	
(Downy brome)	

TRIFLURALIN HF EPA REG. NO. 34704

- Cheat (*Bromus secalinus*)
- (Chess)
- Crabgrass (*Digitaria sp.*)
- (Large crabgrass)
- (Smooth crabgrass)
- Fall panicum (*Panicum dichotomiflorum*)
- (Spreading panicgrass)
- (See Corn, Cotton and Soybean sections for special instructions.)
- Foxtails (*Setaria sp.*)
- (Bottlegrass)
- (Bristlegrass)
- (Giant foxtail)
- (Green foxtail)*
- (Pigeongrass)
- (Robust foxtail)
- (Yellow foxtail)
- Goosegrass* (*Elysiue indica*)
- (Silver crabgrass)
- (Silvergrass)
- (Winegrass)
- (Yardgrass)
- Guinea grass (*Panicum maximum*)
- (See Sugarcane section for special instructions.)
- Johnsongrass (*Sorghum halepense*)
- (Seedling and rhizome)
- (See Soybean section for special instructions on rhizome control.)
- Junglerice (*Echinochloa colonum*)
- Raouigrass (*Hottboellia exaltata*)
- (Itchgrass)
- (See Sugarcane section for special instructions.)
- Sandbur (*Cenchrus incertus*)
- (Burggrass)
- Sprangletop (*Leptochloa filiformis*)
- Stinkgrass (*Eragrostis cilianensis*)
- (Lovegrass)
- Texas panicum (*Panicum texanum*)
- (Buffalograss)
- (Coloradograss)
- (See Corn, Cotton, and Soybean sections for special instructions.)
- Wild Cane (*Sorghum bicolor*)
- (Shattercane)
- (See Soybean section for special instructions.)

*May be locally resistant.

BROADLEAF WEEDS CONTROLLED

- Carpetweed (*Mollugo verticillata*)
- Chickweed (*Stellaria media*)
- Field Bindweed (*Convolvulus arvensis*)
- (See Trees and Vineyard section for special instructions.)
- Florida pusley (*Richardia scabra*)
- (Florida purslane)
- (Mexican clover)
- (Pusley)
- Goosefoot (*Chenopodium hybridum*)
- Henbit (*Lamium amplexicaule*)
- (Fall application only.)
- Knotweed (*Polygonum aviculare*)
- Lambsquarters (*Chenopodium album*)
- Pigweeds (*Amaranthus sp.*)
- (Carelessweed)
- (Prostrate pigweed)
- (Redroot)
- (Rough pigweed)
- (Spiny pigweed)
- Puncturevine (Western U.S. only) (*Tribulus terrestris*)
- (Caltrop)
- Purslane (*Portulaca oleracea*)
- Stinging nettle (*Urtica dioica*)
- (Nettle)

TRIFLURALIN HF will not control certain resistant weeds such as cocklebur, jimsonweed, kochia, nutsedge (nutgrass), ragweed, Russian thistle, velvetleaf or Venice mallow.

Weeds controlled in soybeans by the TRIFLURALIN HF/Sencor* or TRIFLURALIN HF/Lexone* tank-mix in addition to those controlled by TRIFLURALIN HF alone.

- (See Soybean section for special instructions.)
- Jimsonweed (*Datura stramonium*)
- Mallow, Venice (*Hibiscus trionum*)
- (Flower-of-an-hour)
- Mustard, wild (*Brassica kaber*)
- (Charlock)
- (Field mustard)
- Ragweed, common (*Ambrosia artemisiifolia*)
- Sesbania, hemp (*Sesbania exaltata*)
- (Coffeebean)
- (Indigo)
- Smartweed, Annual (*Polygonum pensylvanicum*)
- (Pennsylvania smartweed)
- (Smartweed)

- Prickly sida (*Sida spinosa*)
- (Teaweed)
- (Spiny sida)
- Velvetleaf (*Abutilon theophrasti*)
- (Butterprint)
- (Buttonweed)
- (Cottonweed)
- (Elephant's Ear)
- (Indian mallow)
- (Piemarker)

Cocklebur, morning-glory and giant ragweed: Control of cocklebur, morning-glory and giant ragweed (horseweed) may be erratic, ranging from poor to excellent depending upon soil temperature, time of weed germination, depth of weed seed in the soil and the amount and timing of soil moisture. Control may be improved with timely cultivation. Where cocklebur is a serious problem, an overlay of Sencor or Lexone may be preferable to a tank mix.

Weeds controlled in dry beans and potatoes by the TRIFLURALIN HF/ Eptam* /EPTC tank-mix in addition to those controlled by TRIFLURALIN HF alone. (See Drybean, Potato section for special instructions.)

- Henbit (*Lamium amplexicaule*)
- (Spring applications)
- Nightshade, black (*Solanum nigrum*)
- Nightshade, hairy (*Solanum sarachoides*)
- Nutsedge (*Cyperus sp.*)
- (Nutgrass)
- (Purple nutsedge)
- (Yellow nutsedge)

- Oat, wild (*Avena fatua*)

Weeds controlled in soybeans by the TRIFLURALIN HF/Amiben* tank mix in addition to those controlled by TRIFLURALIN HF alone. (See Soybean section for special instructions.)

- Ragweed, Common (*Ambrosia artemisiifolia*)
- Smartweed, Pennsylvania (*Polygonum pensylvanicum*)
- Velvetleaf (Buttonweed) (*Abutilon theophrasti*)

TRIFLURALIN HF preplant soil incorporated with an Amiben application pre-emergence controls the following additional weeds:

- Coffeeweed (Sesbania) (*Sesbania exaltata*)
- Mustard, Wild (*Brassica kaber*)
- Nightshade, Black (*Solanum nigrum*)
- Prickly sida (Teaweed) (*Sida spinosa*)
- Ragweed, Common (*Ambrosia artemisiifolia*)
- Spurge, Annual (*Euphorbia maculata*)
- Smartweed, Pennsylvania (*Polygonum pensylvanicum*)
- Stinkgrass (*Eragrostis cilianensis*)
- Velvetleaf (Buttonweed) (*Abutilon theophrasti*)

Weeds controlled in cotton by the TRIFLURALIN HF/Caparol* tank mix in addition to those controlled by TRIFLURALIN HF alone. (See Cotton section for special instructions.)

- Smartweed
- Prickly sida (Teaweed)
- Annual morning-glory
- Ragweed
- Groundcherry (Annual)
- Mustard
- Malva
- Wild oat

The tank mix also controls shallow-germinating seedlings of: Cocklebur, Coffeeweed

Weeds controlled in cotton by the TRIFLURALIN HF/Cotoran* tank mix or Cotoran overlaid post plant pre-emergence in addition to those controlled by TRIFLURALIN HF alone, where TRIFLURALIN HF has been applied as a preplant soil incorporated herbicide in cotton. (See Cotton section for special instructions.)

- Ryegrass
- Buttonweed
- Cocklebur
- Goathead
- Groundcherry, Wright
- Jimsonweed
- Morning-glory
- Prickly sida (Teaweed)
- Ragweed
- Sesbania
- Sicklepod
- Smartweed
- Tumbleweed

Weeds controlled in cotton by an overlay treatment of Diuron 80 post plant pre-emergence in fields where TRIFLURALIN HF has been applied as a preplant soil incorporated herbicide in addition to those controlled by TRIFLURALIN HF alone. (See Cotton section for special instructions.)

- Ragweed
- Groundcherry (Annual)
- Dogfennel
- Pennycress
- Morning-glory, Annual
- Shepherd's purse
- Velvetgrass
- Wild lettuce
- Wild mustard

The tank mix of TRIFLURALIN HF plus Far-Go will control wild oat in peas grown in Idaho, Oregon, and Washington in addition to the weeds controlled by TRIFLURALIN HF alone. (See Peas section for special instructions.)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination,

TRIFLURALIN HF

EPA REG. NO. 34704

notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls, chemical-resistant gloves, such as barrier laminate or viton, and shoes plus socks, and protective eyewear.

STORAGE AND DISPOSAL

PROHIBITIONS—Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE—Avoid freezing. Do not store below temperature of (40°F). If frozen, poor weed control may result. Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL—Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. **Plastic:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

TRIFLURALIN HF is a pre-emergence herbicide which is mixed (incorporated) into the soil to provide long-lasting control of a wide range of annual grasses and broadleaf weeds. TRIFLURALIN HF controls weeds as they germinate but will not control established weeds.

SOIL TEXTURE

One key to getting good results with TRIFLURALIN HF is to know your soil texture so that you can apply the correct rate. The amount you apply to your soil will vary with the soil texture. A fine-textured soil requires more TRIFLURALIN HF than a coarse-textured soil.

Soil Texture Guide:

Refer to the following guide to determine your soil texture.

Coarse** Soils	Sand Loamy sand Sandy loam Loam
Medium Soils	Silty clay loam* Silt loam Silt Sandy clay loam*
Fine** Soils	Clay Clay loam Silty clay loam* Silty clay Sandy clay Sandy clay loam*

* Silty clay loam and sandy clay loam soils are transitional soils and may be classified as either medium or fine-textured soils. If silty clay loam or sandy clay loam soils are predominately sand or silt, they are usually classified as medium-textured soils; if predominately clay, they are usually classified as fine-textured soils.

**Soil classification nomenclature for light and heavy-textured soils have been changed to coarse and fine-textured soils at the direction of the Environmental Protection Agency. Throughout this label all previous references to light-textured soils have been changed to coarse-textured soils. All previous references to heavy-textured soils have been changed to fine textured soils. These new descriptions relate more closely to the size of the soil particles in a given classification.

SOIL PREPARATION

Destroy existing weeds before TRIFLURALIN HF application. Chop and thoroughly mix crop residues into the soil to a depth of at least 4 to 6 inches by deep plowing or discing before application. Use machinery that breaks up large clods before application. Crop residue and soil surface must allow for uniform incorporation into the top 2 to 3 inches of soil.

Spray Drift Labeling

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following Aerial Drift Reduction Advisory Information.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzle instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

WIND

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Application should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions), indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

APPLICATION

Add the recommended amount of TRIFLURALIN HF to clean water in the spray tank during the filling operation. Agitate before spraying. Apply in from 5 to 40 gallons of water per acre (broadcast basis), using any properly calibrated low-pressure herbicide sprayer that will apply the spray uniformly. As the amount of water used (spray volume) decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to insure proper calibration and uniform application. Apply TRIFLURALIN HF to the soil surface and incorporate in the same operation, if possible. Do not apply to soils which are wet or in poor condition. Do not apply to soils which are subject to prolonged periods of flooding.

TRIFLURALIN HF EPA REG. NO. 34704

AERIAL APPLICATION

For best results from aerial application of TRIFLURALIN HF, apply to a dry soil surface at a spray volume of from 4 to 10 gallons per acre. Adjust pump pressure, nozzle arrangements, flying speed and flying height to provide a uniform application to the soil surface. Use markers to assure proper application spray widths.

Do not apply TRIFLURALIN HF by aircraft when the wind is blowing at a velocity of 5 mph or greater. This will cause drift of spray particles and result in non-uniform application.

GENERAL CHEMIGATION

Use TRIFLURALIN HF at rates and stages of growth recommended on the label, except as noted below. Apply in 1/2 to 1 acre inch of irrigation water. Mechanical incorporation is not necessary when TRIFLURALIN 4 EC is applied through the irrigation system but, except for established alfalfa, soil preparation must be done according to label instructions.

Treat only the following crops at stage named:

ALFALFA: Apply during dormancy, semi-dormancy or immediately following a cutting. Destroy existing weeds before application.

ASPARAGUS: Apply in the winter or early spring after ferns are removed but before spears emerge.

POTATOES: (Columbia River Basin of Washington and Oregon only): Apply after potato plants have fully emerged on coarse and medium soils.

BEANS: (All types named on label): Preplant except no fall application.

SOYBEANS: Preplant except no fall application.

SUGAR BEETS: Apply when plants are between 2 and 6 inches tall.

CARROTS: Apply before planting.

FIELD CORN: Two-leaf to 30 inches.

Use 1 1/2 to 2 pints per acre. Apply this product only through sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation system(s). Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS Note: Platte Chemical Company does not encourage connecting chemigation systems to public water supplies. The following information is provided for users who have diligently considered all other application and water supply options before electing to make such a connection.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction.

As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either

automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Mix in clean supply tank the recommended amount of this product for acreage to be covered, and needed quantity of water.

This product should not be tank-mixed with other pesticides, surfactants or fertilizers unless prior use has shown the combination noninjurious under your conditions of use.

Follow precautionary statements and directions for all tank-mix products. Meter this product into the irrigation water uniformly during the period of operation.

Do not overlap application. Follow recommended label rates, application timing, and other directions and precautions for crop being treated. Continuous mild agitation of pesticide mixture may be needed to assure a uniform application, particularly if the supply tank requires a number of hours to empty.

INCORPORATION DIRECTIONS

Incorporation Before Planting

For best results TRIFLURALIN HF should be incorporated as soon as possible after application. TRIFLURALIN HF must be incorporated one time within 24 hours after application. A second incorporation is required with most equipment (see Incorporation Equipment section for specific instructions). If TRIFLURALIN HF is applied to a wet, warm soil surface or if the wind velocity is 10 mph or higher variable weed control may result from delaying the first incorporation beyond 24 hours.

Incorporation should place the TRIFLURALIN HF into the top 2 to 3 inches of the final seedbed. Generally, incorporation equipment will place the chemical approximately half as deep as the equipment is run. For example, a disc running 4 inches deep will incorporate TRIFLURALIN HF approximately 2 inches deep.

Incorporation After Planting

(Check crop list for those crops approved for incorporation after planting.)

When incorporating TRIFLURALIN HF after planting or on established row crops use P.T.O.-driven equipment or rolling cultivators. Adjust equipment to till the soil over the seed or throw treated soil toward the crop. Avoid disturbing the seed or mechanically damaging the crop.

Incorporation In Bedded Culture

For effective weed control, TRIFLURALIN HF should be incorporated into the top 2 to 3 inches of the final seedbed.

Knock off beds to planting height before application and incorporation on bedded ground. If TRIFLURALIN HF is applied and incorporated before bedding, do not furrow out deeper than the depth to which TRIFLURALIN HF was incorporated. Furrowing too deep will expose untreated soil and allow weeds to germinate in the bottom of the furrow.

Avoid removal of treated soil from the seedbed before or during the planting operation. This will expose untreated soil and allow weeds to germinate in the drill row.

Incorporation Equipment

Use machinery that mixes TRIFLURALIN HF thoroughly with the soil. Shallow incorporation with implements set to cut less than 2 inches deep may result in erratic weed control. Use of incorporation equipment not listed upon the label may result in poor or erratic weed control and/or crop injury. Except as recommended in the individual crop directions, recommended equipment includes:

Disc set to cut 4 to 6 inches deep and operated in 2 different directions at 4 to 6 mph. A tandem or double-disc operated one time does not provide adequate incorporation.

Field cultivator set to cut 3 to 4 inches deep and operated at 5 mph or more. The field cultivator used alone or in combination with the double-disc will provide effective incorporation providing the following instructions are used:

1. Two passes over the field with a field cultivator with the second pass running at an angle to the first. Do not set cultivator to cut deeper than 4 inches, particularly on the second pass, since untreated soil may be turned up.
2. Field cultivator used for the first pass and the double-disc used for the second pass.
3. Double-disc used for the first pass and the field cultivator used for the second pass.

NOTE: A field cultivator is considered as an implement with 3 to 4 rows of sweeps, spaced at intervals of 7 inches or less and staggered so that no soil is left unturned. Chisel points should not be used.

Rolling cultivator set to cut 2 to 4 inches deep and operated 2 times at 6 to 8 mph. Rolling cultivators are adequate for use on coarse and medium-textured soils only (except when used in sugarcane where the rolling cultivator may be used on fine-textured soils).

Bed conditioner (Do-All) set to cut 2 to 4 inches deep and operated one time at 4 to 6 mph. Bed conditioners are adequate for use on coarse and medium-textured soils only. In bedded culture, one pass is adequate.

Mulch treader and other similar disc-type implements set to cut 3 to 4 inches deep and operated at 5 to 8 mph in two different directions.

P.T.O.-driven equipment (tillers, cultivators, hoes) set to cut 2 to 3 inches deep with rotors spaced to provide a clean sweep of the soil and operated one time. P.T.O.-driven equipment should not be operated at a speed greater than 4 mph.

Other equipment, including the flexible tine-tooth harrow (Flextine, Melroe) is also recommended but only for the special programs for which it is specified in this label.

TRIFLURALIN HF

EPA REG. NO. 34704

CULTIVATION AFTER PLANTING

Soil treated with TRIFLURALIN HF may be shallow-cultivated, rotary-hoed or hand-hoed without reducing the weed control activity of TRIFLURALIN HF. Do not cultivate deeper than the TRIFLURALIN HF treated layer of soil since this may bring untreated soil to the surface and poor weed control may result.

REGIONAL USE MAP



All crop recommendations on this label are given on a regional basis. The dividing line between the Eastern and Western United States is that point where the average rainfall per year is a minimum of 20 to 25 inches. Because the map is based on average conditions, it should be used as a guideline only. Consult your local Agricultural Extension Service and local weather service for information on specific use recommendations and conditions in your area. Note: Sugar beet growing areas of Minnesota and North Dakota are sometimes grouped with Western U.S. on this label.

CROP RECOMMENDATIONS

Important: The following crop recommendations are based on average rainfall conditions. When the annual rainfall amount in your area is radically different than normal, the use recommendations on this label may not be appropriate. For example, below normal rainfall in the Eastern U.S. may result in abnormally long TRIFLURALIN HF carryover. Planting of susceptible rotational crops may then result in crop loss or injury. In the Western U.S., abnormally high rainfall may reduce the period of effective weed control. For all areas, use rates and rotational crops should be determined based on both local factors and crop recommendations on this label. Read the NOTICE OF WARRANTY for manufacturer's disclaimer of liability. If its conditions are not acceptable, do not use this product.

OTHER CROPS

Vegetable crops other than those listed on this label for use with preplant soil incorporated application of Trifluralin HF should not be planted within 5 months after an application of Trifluralin HF.

Eastern United States

GENERAL

These recommendations are given as the broadcast (overall) rates of TRIFLURALIN HF per acre. For band applications, use proportionately less. Apply TRIFLURALIN HF any time after January 1 when the soil can be worked. Also see general and specific fall application directions. TRIFLURALIN HF is not recommended on muck soils. Where a rate range is shown, use the lighter rate for more coarse soils or soils with lower organic matter.

Western United States

GENERAL

These recommendations are given as the broadcast (overall) rates of TRIFLURALIN HF per acre. For band application, use proportionately less. TRIFLURALIN HF is not recommended for peat soils exceeding 20% organic matter or on any muck soils. Do not exceed recommended rates at any time. Where a rate range is shown, use the lighter rate for more coarse soils or soils with lower organic matter.

TRIFLURALIN HF APPLIED ALONE AND IN TANK MIXES WITH FLUID FERTILIZERS

GENERAL

TRIFLURALIN HF alone and TRIFLURALIN HF tank mixes may be mixed with most fluid (liquid) fertilizer materials. TRIFLURALIN HF alone and TRIFLURALIN HF combinations with solution and suspension-type fertilizers has provided weed and grass control as claimed on the respective labels.

All recommendations for TRIFLURALIN HF alone or TRIFLURALIN HF tank-mix combinations regarding rates per acre, approved crops, incorporation, special instructions, warnings and special precautions must be followed.

All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the fertilizer and chemical mixture.

COMPATIBILITY TEST

TRIFLURALIN HF alone and TRIFLURALIN HF combinations and some fluid fertilizer materials may not combine properly. Small quantities should always be tested before full-scale mixing.

- Put 1 pint of fertilizer mixture in a quart jar.
- Add 2 teaspoonfuls of TRIFLURALIN HF and proportional amount of other tank mix products.

- Close jar and shake well.
- Watch mixture for several seconds, check again 30 minutes later.
- If the mix does not separate, or if agitation is only needed to resuspend the mix, the combination may be used. If the mixture separates, gets very thick or syrupy, DO NOT combine for field application.
- Mixing ability may be improved by adding a compatibility agent. Follow the procedure outlined above and add 0.1 teaspoonful of the compatibility agent in Step 2. Complete the other steps to determine if the compatibility agent solves the problem.

If one is needed use a compatibility agent cleared for use on growing crops.

MIXING

If a compatibility agent is needed, add it to the fluid fertilizer before adding the TRIFLURALIN HF alone or TRIFLURALIN HF combination. If compatibility is a problem, mix 2 quarts of water with 1 quart of TRIFLURALIN HF alone before pouring into the fertilizer.

Usually, TRIFLURALIN HF alone may be poured directly into the fluid fertilizer and mixed thoroughly. Wettable powders, dry flowables, liquid flowables or aqueous suspensions should be mixed with the liquid fertilizer before adding TRIFLURALIN HF. Add solution formulations last. Continued agitation is needed until application is complete.

APPLICATION

Spread the fertilizer/chemical mixture normally with a properly calibrated applicator. Be certain the material is applied uniformly to the soil surface.

INCORPORATION

Follow normal TRIFLURALIN HF incorporation procedures.

TRIFLURALIN HF APPLICATION WITH DRY BULK FERTILIZERS

GENERAL

Dry bulk fertilizers may be impregnated or coated with TRIFLURALIN HF. Application of dry bulk fertilizers impregnated with TRIFLURALIN HF has provided weed and grass control equal to the same rates of TRIFLURALIN HF applied in water.

All TRIFLURALIN HF label recommendations regarding rates per acre, approved crops, incorporation, special instructions, cautions and special precautions must be followed.

All individual state regulations relating to dry bulk fertilizer blending, registration, labeling and application are the responsibility of the individual and/or company selling the fertilizer and chemical mixture.

LIMITATIONS

Apply a minimum of 200 pounds per acre of dry fertilizer impregnated with TRIFLURALIN HF at the recommended rates. Any commonly used dry fertilizers can be used for TRIFLURALIN HF impregnation except straight coated ammonium nitrate and straight limestone. These materials will not absorb the herbicide. Blends containing mixtures of these materials can be impregnated.

IMPREGNATION

Use any closed drum, belt, ribbon or other commonly used dry bulk fertilizer blender. The nozzle or nozzles used to spray the TRIFLURALIN HF on to the fertilizer should be placed to provide uniform spray coverage.

RATES

Check the crop section to determine the rate of TRIFLURALIN HF per acre. See the rate table which follows to determine the amount of TRIFLURALIN HF to be impregnated on a ton of dry bulk fertilizer based on the amount of fertilizer which will be applied per acre. (See rate chart.)

APPLICATION

Spread the fertilizer/chemical mixture normally with a properly calibrated applicator. Be certain the material is applied uniformly to the soil surface.

INCORPORATION

Follow normal TRIFLURALIN HF incorporation procedures.

RATE CHART FOR IMPREGNATING FERTILIZER WITH TRIFLURALIN HF

Fertilizer Rate Per Acre	TRIFLURALIN HF added to 2-TON of fertilizer	
	TRIFLURALIN HF Rate	Per Acre
200 pounds	1 pint	1 1/2 pints
	10 pts. or 5 qts. per ton	15 pts. or 7 1/2 qts. per ton
250 pounds	8 pts. or 4 qts. per ton	3 qts. or 1 1/2 gal. per ton
	6 3/4 pts. or 3 1/2 qts. per ton	10 pts. or 5 qts. per ton
300 pounds	5 3/4 pts. or 2 3/4 qts. per ton	9 pts. or 1 1/8 gal. per ton
	5 pts. or 2 1/2 qts. per ton	7 1/2 pts. or 1 gal. per ton
400 pounds	4 1/2 pts. or 2 1/4 qts. per ton	3 1/2 qts. or 7/8 gal. per ton
	4 1/2 pts. or 2 1/4 qts. per ton	7/8 gal. per ton

TRIFLURALIN HF

EPA REG. NO. 34704

RATE CHART FOR IMPREGNATING FERTILIZER WITH TRIFLURALIN HF cont'd.

Fertilizer Rate Per Acre	TRIFLURALIN HF Rate Per Acre		
	2 pints	3 pints	4 pints
200 pounds	10 qts. or 2 1/2 gal. per ton	15 qts. or 3 3/4 gal. per ton	20 qts. or 5 gal. per ton
250 pounds	8 qts. or 2 gal. per ton	12 qts. or 3 gal. per ton	16 qts. or 4 gal. per ton
300 pounds	14 pts. or 1 3/4 gal. per ton	20 pts. or 2 1/2 gal. per ton	27 pts. or 3 1/3 qt. per ton
350 pounds	12 pts. or 1 1/2 gal. per ton	17 pts. or 2 1/4 gal. per ton	23 pts. or 2 7/8 gal. per ton
400 pounds	5 qts. or 1 1/4 gal. per ton	15 pts. or 1 7/8 gal. per ton	10 qts. or 2 1/2 gal. per ton
450 pounds	4 1/2 qts. or 1 1/8 gal. per ton	13 1/2 pts. or 1 3/4 gal. per ton	9 qts. or 2 1/4 gal. per ton

For rates other than those listed above, use the following formula to calculate the amount of TRIFLURALIN HF to be impregnated on a ton of dry bulk fertilizer:

$$\frac{\text{Pints TRIFLURALIN 4EC}}{\text{Per Acre}} \times \frac{1000}{\text{lbs. Fertilizer}} = \frac{\text{Quans TRIFLURALIN 4EC}}{\text{per Ton of Fertilizer}}$$

FALL APPLICATION—General—For areas receiving more than 20 inches average annual rainfall: See directions for specific crops. For any crop on the TRIFLURALIN HF label for which there is no specific directions for fall application and for which preemergence application is recommended, use the rate shown for spring application.

Exceptions: Do not fall apply TRIFLURALIN HF on sugarbeets, potatoes and direct-seeded tomatoes. Do not make fall applications to soils that are wet, prone to prolonged periods of flooding, or where rice was grown the previous season.

Soil Preparation: Ground may be left flat or bedded-up over winter. For bedded ground, knock beds down to desired height before planting, moving some treated soil from beds into furrows. If soil is left over winter, exercise care to not turn up untreated soil during spring bedding operations. Remove established weeds during seedbed preparations as they will not be controlled by TRIFLURALIN HF. If weeds become established in furrows due to uncovering of untreated soil during bedding, destroy such weeds before planting.

Timing: In most states, apply and incorporate TRIFLURALIN HF any time between October 15 and December 31. In the states of MN, MT, ND and SD, apply and incorporate TRIFLURALIN HF between September 1 and December 31.

ALFALFA—Established Alfalfa Only: (Western U.S. Only)
Apply to established alfalfa stands at a broadcast rate per acre of 1 1/2 pints on coarse soils and 2 pints on medium and fine soils. Use incorporation equipment that will insure thorough soil mixing with a minimum of damage to the established alfalfa.

ASPARAGUS—Established—Single or split application for pre-emergence weed control: Follow recommended procedures for soil preparation application and incorporation found near the beginning of the TRIFLURALIN HF label. Volunteer seedling asparagus and field bindweed will be suppressed (reduced in stand and vigor) in addition to weeds otherwise controlled by this product.

Single application—Apply in the winter or early spring after ferns are removed but before spears emerge. Alternatively, apply after harvest in the late spring or early summer before ferning begins.

Split application—Apply at both times described in "Single Application" (before harvest and after harvest), but at reduced rate.

SOIL TEXTURE	BROADCAST TRIFLURALIN HF RATE			
	Split Application		Single Application	
	Before Harvest	After Harvest	Before OR Harvest	After Harvest
Coarse soils	1 pt. + 1 pt.		2 pts. OR 2 pts.	
Medium Soils	1 1/2 pts. + 1 1/2 pts.		3 pts. OR 3 pts.	
Fine soils	2 pts. + 2 pts.		4 pts. OR 4 pts.	

The maximum application per calendar year is 2 pts. per acre on coarse soils, 3 pts. per acre on medium soils and 4 pts. per acre on fine soils.

BEANS—Castor Beans:
Apply and incorporate TRIFLURALIN HF before planting.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2-2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

BEANS—Dry Beans (Kidney, Navy, Pinto, Etc.):
Apply and incorporate TRIFLURALIN HF before planting.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2-2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

BEANS—TRIFLURALIN HF/Eptam/EPTC Tank-Mix for Dry Beans:
The TRIFLURALIN HF/Eptam/EPTC tank-mix effectively controls henbit, black nightshade and nutsedge (nutgrass) in addition to all of the annual grasses and broadleaf weeds listed on the TRIFLURALIN HF label. Follow normal procedures for soil preparation. The TRIFLURALIN HF/Eptam/EPTC tank-mix should be applied from 2 days before planting (up to planting in the Eastern U.S.).

Apply at a broadcast rate of 1 pint of TRIFLURALIN HF and 2 1/2 to 3 1/2 pints of Eptam/EPTC 7E per acre or up to the label recommended rate for each herbicide depending on soil texture and weed problem. Use the higher rate of Eptam/EPTC for nutsedge control. TRIFLURALIN HF at 1 pint per acre, alone or in combination, should not be used on soils containing 5% or more organic matter. Incorporate immediately after application. Follow normal TRIFLURALIN HF procedures for cultivation. **Caution:** Read the Eptam/EPTC label before using. Observe all cautions and limitations on labeling of all products used in mixtures. The combination of TRIFLURALIN HF and Eptam/EPTC should not be used on soybeans, black-eyed peas (beans), lima beans and other flatpodded beans, except Romano. Do not use the foliage from a crop treated with the TRIFLURALIN HF/Eptam tank-mix for feed or for grazing.

BEANS—Fall Application in Dry Beans Grown in Idaho, Oregon and Washington Only:

For dry beans grown in Idaho, Oregon and Washington: Apply and incorporate TRIFLURALIN HF at a broadcast rate per acre of 1 pint on coarse soils; 1 1/4 to 1 1/2 pints on medium soils; and 1 1/2 pints on fine soils.

BEANS—Guar Beans and Mungbeans:
Apply and incorporate TRIFLURALIN HF before planting at a broadcast rate per acre of 1 pint on coarse soils and 1 1/2 pints on medium and fine soils.

BEANS—Lima Beans and Snap Beans:
Apply and incorporate TRIFLURALIN HF before planting at a broadcast rate per acre of 1 pint on coarse and medium soils and 1 1/2 pints on fine soils.

CARROTS:
Apply and incorporate TRIFLURALIN HF before planting.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2-2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

CELERY—Both Direct-seeded and Transplant: (Western U.S. Only)
Apply and incorporate TRIFLURALIN HF before planting or transplanting at a broadcast rate per acre of 1 pint on coarse soils; 1 1/4 to 1 1/2 pints on medium soils; 1 1/2 pints on fine soils; 1 1/2 to 2 pints on soils with 2 to 5% organic matter; and 2 pints on soils with 5.1 to 10% organic matter.

COLE CROPS—Broccoli, Brussels Sprouts, Cabbage and Cauliflower:

For Direct-Seeded cole crops apply and incorporate before planting. (Eastern U.S.—Direct-seeded cole crops have exhibited marginal tolerance to recommended rates of TRIFLURALIN HF. Stunting or reduced stands may occur.)

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 pt.	1 pt.
Fine soils	1 1/2 pts.	1 pt.
Soils with 2 to 5% organic matter	1 1/2 pts.	—
Soils with 2 to 10% organic matter	—	1 1/2 pts.

For Transplant cole crops apply and incorporate TRIFLURALIN HF before transplanting. Do not apply TRIFLURALIN HF after transplanting.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4 to 1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2 pts.	1 1/2 to 2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

CORN (FIELD), GRAIN SORGHUM (MILC)—Over-the-top or directed spray for preemergent weed control: (see "WEEDS AND GRASSES CONTROLLED" section of label).

Field Preparation—Field should be cultivated prior to application of TRIFLURALIN HF to provide loose tilth, remove established weeds and deposit a soil cover at the base of crop plants.

Application—Make application when the crop is well established (minimum 8 inch height). Apply the recommended rate either as an over-the-top spray or as a directed spray. Cover soil surface uniformly with spray using drop nozzles if required. Use the lower rates when light weed pressure is expected and higher rates when heavy weed pressure is expected.

TRIFLURALIN HF

EPA REG. NO. 34704

Apply and incorporate the following rates per acre:

Coarse soils	3/4 to 1 pt.*
Medium soils	1 to 1 1/2 pts.
Fine soils	1 1/2 to 2 pts.

*In corn only, apply 1 to 1 1/2 pts. to control fall panicum and Texas panicum in the states of AL, FL, GA, NC, SC and VA.

Incorporation—A correctly adjusted rolling cultivator or sweep-type cultivator can accomplish incorporation in one pass. Carefully adjust incorporation implement to avoid direct injury to crop. A sweep-type cultivator should have 3 to 5 sweeps per row middle and be operated at 6 to 8 mph. Adjust sweeps so as to avoid exposing untreated soil.

Important: Do not use TRIFLURALIN HF on corn grown for seed. Do not make preplant or crop preemergence applications to corn or sorghum as crop injury or loss may occur.

COTTON—Pre-emergence applications:

Apply and incorporate TRIFLURALIN HF before planting, at planting or immediately after planting using the following broadcast rates per acre:

	Eastern U.S.	Western U.S.
Coarse soils 1 pt. 1 pt.
Medium soils 1 1/2 pts. 1 1/4 to 1 1/2 pts.
Fine soils 2 pts. 1 1/2 pts.
Soils with 2 to 5% organic matter 1 1/2 pts. 1 1/2 to 2 pts.
Soils with 5.1 to 10% organic matter 2-2 1/2 pts. 2 pts.

When incorporating after planting (post-plant), care must be taken not to disturb the seed.

Seedling disease may weaken cotton plants and increase the possibility of damage from TRIFLURALIN HF. To control seedling disease, use a good fungicide program.

COTTON—Post-emergence applications:

Apply TRIFLURALIN HF any time up to layby, but not less than 90 days before harvest. Direct layby applications to the soil between the rows and beneath emerged cotton plants. Use the same rates as for a pre-emergence application.

COTTON—Fall Application:

For cotton grown in Alabama, Arkansas, northern Florida, Georgia, Louisiana, Mississippi, southeastern Missouri, North Carolina, New Mexico, Oklahoma, South Carolina, Tennessee and Texas; Apply and incorporate TRIFLURALIN HF at a broadcast rate per acre of 2 pints on coarse and medium soils and 2 1/2 pints on fine soils. For cotton grown in Arizona, California and Nevada: Apply and incorporate TRIFLURALIN HF at a broadcast rate per acre of 1 1/2 pints on coarse soils; 2 pints on medium soils and 2 1/2 pints on fine soils. For cotton grown in states other than those listed above: Apply and incorporate TRIFLURALIN HF at a broadcast rate per acre of 1 pint on coarse soils; 1 1/2 pints on medium soils; 2 pints on fine soils; 1 1/2 pints on coarse soils with 2 to 5% organic matter; and 2 to 2 1/2 pints on soils with 5.1 to 10% organic matter.

COTTON—Fall panicum and Texas panicum control:

For the control of fall panicum and Texas panicum in the states of Alabama, Florida, Georgia, North Carolina, South Carolina and Virginia, apply and incorporate TRIFLURALIN HF at the broadcast rate of 2 pints per acre on both coarse and medium soils.

Plant cotton after early season adverse weather conditions have passed. Do not plant cotton deeper than 1 inch. Crop injury in the form of delayed growth or reduced yields may occur under adverse cool, wet weather conditions when TRIFLURALIN HF is used according to these special recommendations.

COTTON—Rhizome Johnsongrass control:

All Cotton Producing States except Arizona and California. Commercially acceptable control of rhizome Johnsongrass can be obtained with a double-rate TRIFLURALIN HF program when applied for 2 years in a row.

Soil Preparation—Proper preparation of the soil before application is very important for satisfactory results. Use a chisel plow or similar implement to bring rhizomes to the top of the soil. Then follow with a disc before application to cut the rhizomes into small (2 to 3-inch) pieces. This should also destroy any emerged Johnsongrass.

Application—Choose the one application program that best fits your cultural practices:

Spring Application—Apply TRIFLURALIN HF any time in the spring before planting for 2 years in a row at a broadcast rate per acre of 2 pints on coarse soils; 3 pints on medium soils and 4 pints on fine soils, OR

Fall Application—Apply TRIFLURALIN HF between October 15 and December 31 for 2 years in a row at the same rates as a spring application for the control of rhizome Johnsongrass.

Incorporation—Deep incorporation is essential to good rhizome Johnsongrass control. Incorporate TRIFLURALIN HF thoroughly with a disc set to cut 4 to 6 inches deep and operate in 2 different directions at 4 to 6 mph.

Cultivation—Some Johnsongrass plants will escape. Timely cultivations during the crop season are necessary to obtain commercially acceptable control. Commercially acceptable control will not be obtained with only 1 year of double-rate TRIFLURALIN HF use.

Precautions: Plant cotton after early season adverse, wet-weather conditions have passed. Crop injury in the form of reduced stands and

delayed growth will occur under adverse cool, wet-weather conditions early in the season and may result in delayed maturity and reduced yields when TRIFLURALIN HF is used according to these recommendations. High quality seed accompanied by a good fungicide program to control seedling diseases in addition to other recommended cultural and chemical practices should be used to minimize crop injury from TRIFLURALIN HF.

In the season following either the 1 or 2-year treatments, plant only those crops for which TRIFLURALIN HF has been registered as a preplant treatment or injury may result.

COTTON—More Complete Control of Pigweed and Seedling Johnsongrass in Cotton Grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Southeastern Missouri, North Carolina, South Carolina, Tennessee and southern Virginia:

For more complete control of pigweed and seedling johnsongrass, TRIFLURALIN HF may be applied preplant at a broadcast rate per acre of from 1 to 1 1/2 pints on coarse soils, from 1 1/2 to 2 pints on medium soils and 2 pints on fine soils except in the state of Louisiana where 3 pints per acre are recommended on fine soils.

Precaution: Plant cotton after early season adverse weather conditions have passed. Do not plant cotton deeper than 1 1/2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when TRIFLURALIN HF is used according to these recommendations.

COTTON—More Complete Weed and Grass Control in Certain Counties Along the Texas Gulf Coast:

For more complete control of those weeds and grasses listed in the TRIFLURALIN HF label in the Texas Gulf Coast Counties of Brazoria, Calhoun, Chambers, Fort Bend, Galveston, Harris, Jackson, Jefferson, Liberty, Matagorda, Orange, Victoria, Waller and Wharton, TRIFLURALIN HF may be applied up to 2 weeks before planting at a broadcast rate of 1 1/2 pints on coarse soils, 2 pints on medium soils and 3 pints on fine soils.

Precaution: Plant cotton after early season adverse weather conditions have passed. Do not plant cotton deeper than 1 1/2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when TRIFLURALIN HF is used according to these recommendations.

COTTON—TRIFLURALIN HF/Caparol® tank mix for cotton grown in California, Arizona, New Mexico, and Texas:

The TRIFLURALIN HF/Caparol combination will control certain grasses and broadleaf weeds listed on the TRIFLURALIN HF label. This combination will also control shallow-germinating seedlings of cocklebur and coffeeweed.

NOTE: This combination will not control sunflower, rhizome Johnsongrass, deep-germinating seedlings of cocklebur and sandbur or established perennials such as Bermudagrass. Follow normal TRIFLURALIN HF procedures for soil preparation and incorporation. Apply the tank mix combination to the flat soil surface before disking.

Broadcast Rates Per Acre:

	TRIFLURALIN HF	Caparol 80W	Caparol 4L
Coarse soils* 1 pt. 1 1/2-2 pounds 2.4-3.2 pts.
Medium soils 1 1/2 pts. 2 1/2 pounds 4 pts.
Fine soils 2 pts. 2 1/2 pounds 4 pts.

*Do not use on sands and loamy sands. For band applications use proportionately less. TRIFLURALIN HF is not recommended for use on muck soils. Use less than 2 lbs. Caparol 80W or 3.2 pts. Caparol 4L only in AZ and CA.

Mixing Directions—Carefully follow the procedures on the Caparol 80W label for making a slurry and adding it to a partially filled tank of water or follow the mixing procedure on the Caparol 4L label for adding it to a partially filled tank of water. After the Caparol is thoroughly mixed with the partially filled tank of water, add the TRIFLURALIN HF and continue filling. Agitate during the filling and spraying operation.

Avoid leaving the spray mixture in the tank without constant agitation. If by-pass agitation is used, it should terminate at the bottom of the tank to minimize foaming.

Incorporation Directions—The first incorporation of TRIFLURALIN HF/Caparol should be immediately following application. A second incorporation is required with most equipment. (See incorporation equipment section for further instructions.)

Precautions: Do not apply more than the recommended rate for your soil texture.

The combination of TRIFLURALIN HF/Caparol should not be used under the following conditions because crop injury may result: in the cut areas of newly leveled fields, in areas of excess salt, and where flooding over the beds is likely to occur.

Do not plant cotton in tractor wheel depression or crop injury may result. On mulch-planted cotton, water back only after cotton seedlings get well established.

Crop Rotations—Cabbages, peas, onions and peas may be planted in the fall after a spring application of TRIFLURALIN HF plus Caparol.

Winter barley, winter rye and winter wheat can be planted in the fall also, if they are plowed down and not used for food or feed. Refer to the Caparol label for directions, cautions and precautions.

COTTON—TRIFLURALIN HF/Cotoran tank mix except in Arizona and California:

The TRIFLURALIN HF/Cotoran tank mix effectively controls all the annual grasses and broadleaf weeds listed on the TRIFLURALIN HF label plus many additional annual grasses and broadleaf weeds (see Weeds Controlled section). Follow normal TRIFLURALIN HF procedures for soil preparation. Apply TRIFLURALIN HF/Cotoran tank mix in 15 to 40 gallons of clean water per acre using any properly calibrated low pressure herbicide sprayer that will apply the spray uniformly.

11/6

TRIFLURALIN HF

EPA REG. NO. 34704

Broadcast Rates Per Acre:

	TRIFLURALIN HF	Cotoran 80W
Coarse soils	1 pt.	1 1/4 pounds
Medium soils	1 1/2 pts.	2 pounds
Fine soils	2 pts.	2 1/2 pounds

Mixing Directions—Carefully follow the procedures on the Cotoran 80W label for making a Cotoran slurry and adding it to a partially filled tank of water. After the Cotoran is thoroughly mixed with the partially filled tank of water, add the TRIFLURALIN HF and continue filling. Agitate continuously throughout the filling and application operations. Follow normal TRIFLURALIN HF incorporation procedures. Do not leave spray mixture in tank without constant agitation. If by-pass agitation is used, it should terminate at the bottom of the tank to minimize foaming.

Precautions: Do not use the tank mix in Arizona and California. Do not plant crops other than cotton on the treated land within 6 months after the application of TRIFLURALIN HF plus Cotoran or injury may occur.

West Texas Only: Do not use the tank mix of TRIFLURALIN HF plus Cotoran on sandy, loamy sand or fine sandy loam soils. Do not use on cotton planted in furrows.

Arkansas, Louisiana, and Mississippi Only: Use 1 pound Cotoran 80W in tank mix with TRIFLURALIN HF on sandy loam soils low in organic matter.

New Mexico: Cotton can be planted the next spring. Do not plant treated areas to crops other than cotton on treated land until 1 year after last application. Do not use on sandy or coarse textured soil of less than 1% organic matter.

Do not feed foliage from treated cotton plants or gin trash to livestock.

The tank mix of TRIFLURALIN HF plus Cotoran is not recommended to be applied in liquid fertilizer.

Refer to the Cotoran label for cautions, precautions, and instructions.

Cotoran overlay: Refer to the Cotoran label for cautions, precautions and instructions.

COTTON—Preplant incorporated TRIFLURALIN HF and surface applied, preemergence Diuron 80 for weed control in cotton grown east of the Mississippi River plus Arkansas, Southeastern Missouri, Louisiana, and Eastern Texas:

Preplant soil incorporated applications of TRIFLURALIN HF (see above for rates) may be followed by a surface applied, post-plant, pre-emergence application of Diuron 80. This combination effectively controls all the weeds controlled by TRIFLURALIN HF plus many additional weeds (see weeds controlled section). Apply Diuron 80 at 0.8 to 1.5 pounds per broadcast acre to the soil surface after planting but prior to crop emergence. The higher rates are used on heavier soil types. Do not use Diuron 80 on light (sandy or low organic) soils. Do not use on heavy clay soils above 10 percent organic matter. Consult the Diuron 80 label for additional instructions, cautions and precautions.

CUCURBITS—Cantaloupes, Cucumbers and Watermelons—Post-Plant, emerged in Texas only:

Apply TRIFLURALIN HF as a directed spray to the soil between the rows and beneath plants which are in the 3 to 4 true-leaf stage. Set incorporation machinery to throw treated soil toward plants in the row. Care should be taken that incorporation machinery does not damage the plants.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/4-1 1/2 pts.	1 1/4 to 1 1/2 pts.
Fine soils	1 1/2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2-2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

FLAX (Fall Application Only)

Trifluralin 4EC should be applied and incorporated in the fall for weed control in spring seeded flax. Incorporate once within 24 hours after application. The second incorporation may be completed in the spring prior to planting.

Instructions for Flax:

Incorporation operations or other tillage practices performed in the spring prior to seeding should be relatively shallow to maintain a firm seedbed. The seedbed should be packed prior to seeding. Seeding should be done with a press drill or hoe drill. Seed into moist seedbed and plant no more than 1 1/2 inches deep.

Seeding should not be completed until soil is sufficiently warm. Refer to the "Special Precautions" section of this label for information on growing conditions that can lead to crop injury or yield reduction.

Broadcast Application Rate/Acre:

Coarse Soils	1 pt.
Medium Soils	1 1/2 pts.
Fine Soils	2 pts.

GREENS—Turnip Greens Grown for Processing and All Collard, Kale and Mustard Greens:

Apply and incorporate TRIFLURALIN HF before planting at a broadcast rate per acre of 1 pint on coarse and medium soils and 1 1/2 pints on fine soils.

MUSTARD—Grown For Seed in Minnesota and North Dakota Only: Apply and incorporate TRIFLURALIN HF before planting at a broadcast rate per acre of 1 pint on coarse and medium soils and 1 1/2 pints on fine

soils.

HOPS: (Western U.S. Only.)

Apply and incorporate TRIFLURALIN HF while the crop is dormant at a broadcast rate per acre of 1 pint on coarse soils; 1 1/4 to 1 1/2 pints on medium soils and 1 1/2 pints on fine soils and soils with 2 to 10% organic matter.

MINT—Established Dappled and Spear Mint (Western U.S. Only.)

Apply TRIFLURALIN HF at a rate per acre of 1 pint on coarse soils; 1 1/4 pints on medium soils; and 1 1/2 pints on fine soils. Use incorporation equipment that will insure thorough soil mixing with a minimum of damage to the established, dormant mint.

OKRA:

Apply and incorporate TRIFLURALIN HF before planting.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2-2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

ONIONS (Dry Bulbs Only)

Direct spray Trifluralin HF between established onion rows and apply as a soil incorporated treatment. Use spray shields to avoid injury to foliage or exposed bulbs. Do not apply within 60 days of harvest.

Broadcast Application Rate/Acre:

Coarse soils	3/4-1 pt.
Medium soils	1-1 1/4 pts.

Use the lower rate in rate range in areas receiving less than 20 inches total annual rainfall and irrigation or where light weed pressure is anticipated.

Incorporation

Incorporate with 1 pass of a sweep-type or rolling cultivator 2 to 4 inches deep and operate at 6 to 8 mph. To avoid crop injury, do not cover exposed onion bulbs with treated soil during incorporation. Be careful not to injure crop roots during incorporation.

Precautions: This product will not adversely affect onions when applied according to directions under normal growing condition. The crop may weaken from diseases, improper incorporation depth, excessive moisture, high salt concentration, or drought which may increase the chance of damage from this product.

PEANUTS—Spanish Peanuts Grown in Texas and Oklahoma Only:

Apply and incorporate TRIFLURALIN HF before planting, at planting or immediately after planting at a broadcast rate per acre of 1 pint on coarse soils. When incorporating after planting, care must be taken not to disturb the seed.

PEAS—Dry (Western U.S. Only.)

Apply and incorporate TRIFLURALIN HF before planting at a broadcast rate per acre of 1 pint on coarse and medium soils and 1 1/2 pints on fine soils.

PEAS—English:

Apply and incorporate TRIFLURALIN HF before planting at a broadcast rate per acre of 1 pint on coarse and medium soils and 1 1/2 pints on fine soils.

PEAS—Fall Application in Dry Peas and English Peas Grown in Idaho, Oregon and Washington Only:

For dry peas grown in Idaho, Oregon and Washington: Apply and incorporate TRIFLURALIN HF at a broadcast rate per acre of 1 pint on coarse soils; 1 1/4 to 1 1/2 pints on medium soils; and 1 1/2 pints on fine soils.

PEAS—Southern Peas:

Apply and incorporate TRIFLURALIN HF before planting.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2-2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

PEAS—TRIFLURALIN HF and Far-Go Tank Mix for weed control in

peas grown in Idaho, Oregon, and Washington: The tank mix combination of TRIFLURALIN HF plus Far-Go will provide control of wild oats in addition to other annual grasses and broadleaf weeds controlled by TRIFLURALIN HF.

Application Rates—Broadcast 3/4 pint of TRIFLURALIN HF on coarser textured soils, 1 pint of TRIFLURALIN HF on fine soils. Use 1 1/4 quarts of Far-Go for all soil textures.

Incorporation Directions—Apply the TRIFLURALIN HF plus Far-Go tank mix and incorporate from 3 weeks before seeding up to immediately before seeding. TRIFLURALIN HF and Far-Go must be thoroughly incorporated into the top 2 inches of the soil by 2 incorporations. The first incorporation should be made as soon as possible on the day of application. The second incorporation should be made as soon as possible but before seeding. Incorporate with a disc-type implement set to cut 4 inches deep and operate in 2 different directions at 4 to 8 mph or with a field cultivator set to cut 3 to 4 inches deep and operate at 5 mph or more. Shallow incorporation with implements set to cut less than 2 inches may

TRIFLURALIN HF

EPA REG. NO. 34704

result in erratic weed control.

NOTE: Do not apply to lentils.

Leaf crinkling and delayed maturity of peas may occur, particularly on clay points in the northwest; but this is usually more than offset by a reduction of wild oats. Do not graze livestock on treated crops. Refer to the cautions, precautions and directions on the Far-Go label.

PEPPERS—Transplants only:

Apply and incorporate TRIFLURALIN HF before transplanting. Do not apply after transplanting.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

POTATOES—Not recommended for use in the state of Maine:

Apply TRIFLURALIN HF after planting, up to or immediately following dragoff in the Eastern U.S. or after planting, before emergence on all soil textures, or after the potato plants have fully emerged on coarse and medium soils in the Western U.S. TRIFLURALIN HF is not recommended on muck soils.

Set incorporation equipment so that the bed and furrow will be uniformly covered with a layer of TRIFLURALIN HF. If the layer of TRIFLURALIN HF treated soil is not uniform and the herbicide is concentrated over the bed, potato emergence may be retarded and stem brittleness can occur. Care should be taken so that incorporation machinery does not damage potato seed pieces or elongating sprouts. Cultivation prior to emergence may result in mechanical injury to the elongated potato sprouts.

When applying and incorporating TRIFLURALIN HF after potato plants have fully emerged, do not completely cover the foliage with treated soil. Likewise do not completely cover foliage at subsequent cultivations.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pt.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

POTATOES—Split Application in Idaho, Oregon and Washington Only:

On all soils apply and incorporate 3/4 pint of TRIFLURALIN HF before planting and 3/4 pint after planting when potato plants have fully emerged. Do not apply to soils containing 2% or more organic matter. Follow incorporation directions listed above for application to potatoes after planting.

POTATOES—TRIFLURALIN HF/Eptam/EPTC Tank-Mix for Potatoes Grown in Kansas, Minnesota, Nebraska, North Dakota, Oklahoma, South Dakota and Texas only: (Eastern U.S. Only.)

The TRIFLURALIN HF/Eptam/EPTC tank-mix effectively controls henbit, black nightshade and nutsedge (nutgrass) in addition to all of the annual grasses and broadleaf weeds listed on the TRIFLURALIN HF label. Follow normal TRIFLURALIN HF procedures for soil preparation. The TRIFLURALIN HF/Eptam/EPTC tank-mix may be applied after planting but prior to crop emergence. In areas where potatoes are normally dragged-off, the TRIFLURALIN HF/Eptam/EPTC tank-mix should be applied and incorporated up to or immediately following dragoff at a broadcast rate per acre of 1 pint of TRIFLURALIN HF and 1 3/4 pints of Eptam/EPTC 7E on all soil textures or up to the label recommended rate for each herbicide depending on soil texture and weed problem. See details on Eptam/EPTC label. TRIFLURALIN HF at 1 pint per acre, alone or in combination, should not be used on soils containing 5% or more organic matter. Incorporate immediately after application. Follow normal TRIFLURALIN HF procedures for cultivation.

Caution: Read the Eptam/EPTC label before using. Observe all cautions and limitations on labeling of all products used in mixtures. Do not graze or feed forage to livestock from fields treated with the TRIFLURALIN HF/Eptam/EPTC tank-mix.

POTATOES—TRIFLURALIN HF/Eptam/EPTC Tank-mix: (Western U.S. Only.)

Application After Planting—The TRIFLURALIN HF/Eptam/EPTC tank-mix effectively controls henbit, nightshade and nutsedge (nutgrass) in addition to all of the annual grasses and broadleaf weeds listed on the TRIFLURALIN HF label. Follow normal procedures for soil preparation. The TRIFLURALIN HF/Eptam/EPTC tank-mix may be applied after planting, up to or immediately following dragoff at a broadcast rate per acre of 1 pint of TRIFLURALIN HF and 1 3/4 pints of Eptam/EPTC 7E on all soil textures or up to the label recommended rate for each herbicide depending on soil texture and weed problem. See details on Eptam/EPTC label. TRIFLURALIN HF at 1 pint per acre, alone or in combination, should not be used on soils containing 5% or more organic matter. Incorporate immediately after application. Follow normal TRIFLURALIN HF procedures for cultivation.

Application Before Planting in Washington, Idaho, and Oregon Only—TRIFLURALIN HF/Eptam/EPTC may be also applied before planting at a broadcast rate of 3/4 pint of TRIFLURALIN HF and 3 1/2 pints of Eptam/EPTC 7E on all soil textures and incorporated immediately.

Caution: Do not use this tank-mix both before and after planting in the same season. Read the Eptam/EPTC label before using. Observe all cautions and limitations on labeling of all products used in mixtures. Do not graze or feed forage to livestock from fields treated with the TRIFLURALIN HF/Eptam/EPTC tank-mix.

RADISH

Trifluralin 4EC should be applied as a preplant soil incorporated treatment.

Broadcast Application Rate/Acre:

Coarse soils	1 pt.
Medium soils	1 1/2 pts.
Fine soils	1 1/2 pts.

RAPESEED (CANOLA)

Trifluralin 4EC may be applied and incorporated in the spring before planting or in fall. See "Soil Preparation" in the "Fall Application" section of this label.

Broadcast Application Rate/Acre:

Coarse soils	1 pt.
Medium soils	1 1/2 pts.
Fine soils	2 pts.

Precautions: Do not apply this product to rapeseed (canola) grown in the state of Alaska.

SAFFLOWER:

Apply and incorporate TRIFLURALIN HF before planting.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2-2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.
Soils with 10.1 to 20% organic matter	—	2-3 pts.

SAFFLOWER—FALL APPLICATION:

For safflower grown in Arizona, California, Idaho, Montana, Nevada, Oregon, Utah, Washington and Wyoming: Apply and incorporate TRIFLURALIN HF at a broadcast rate per acre of 1 1/2 pints on coarse soils; 2 pints on medium soils; and 2 1/2 pints on fine soils.

SOYBEANS—Preplant incorporated:

Apply and incorporate TRIFLURALIN HF before planting. Do not plant soybeans deeper than 2 inches. Follow recommended soil preparation, application and incorporation procedures.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2-2 1/2 pts.*	2 pts.

* Except charcoal soils in Arkansas, Louisiana and Mississippi. See below.

SOYBEANS—Soils Containing Charcoal in Arkansas, Louisiana and Mississippi:

Newly cleared land often contains high organic matter (4 to 10%) and charcoal which result from burning debris. This charcoal and/or organic matter tends to tie up TRIFLURALIN HF and reduce its weed control activity. Higher rates of TRIFLURALIN HF are therefore necessary for satisfactory weed control. Increased rates can cause crop injury if charcoal or a high percentage of organic matter is not present to tie up some of the TRIFLURALIN HF. In the actual windrow or burn row, where a high level of charcoal is present, poor weed control may result even with an increased rate of TRIFLURALIN HF.

Apply and incorporate at the following broadcast rates per acre:

Coarse soils	1 1/2 pts.
Medium soils	2 1/4 pts.
Fine soils	3 pts.

SOYBEANS—Fall application:

For soybeans grown in Arkansas, Arkansas, northern Florida, Georgia, Louisiana, Mississippi, southeastern Missouri, North Carolina, Oklahoma, South Carolina, Tennessee or Texas: Apply and incorporate TRIFLURALIN HF at a broadcast rate per acre of 2 pints on coarse and medium soils and 2 1/2 pints on fine soils. For soybeans grown in the Eastern United States other than those states listed above: Apply and incorporate TRIFLURALIN HF at a broadcast rate per acre of 1 pint on coarse soils; 1 1/2 pints on medium soils; 2 pints on fine soils; 1 1/2 pints on coarse soils with 2 to 5% organic matter; and 2 to 2 1/2 pints on soils with 5.1 to 10% organic matter.

SOYBEANS—Fall panicle and Texas panicle control:

For the control of fall panicle and Texas panicle in the states of Alabama, Florida, Georgia, North Carolina, South Carolina and Virginia, apply TRIFLURALIN HF at the broadcast rate of 2 pints per acre on both coarse and medium soils. Plant soybeans after early season adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury in the form of delayed growth or reduced yields may occur under adverse cool, wet weather conditions when TRIFLURALIN HF is used according to these special recommendations.

SOYBEANS—More Complete Control of Pigweed and Seedling Johnsongrass in Soybeans Grown in Alabama, Arkansas, Florida,

12/16

TRIFLURALIN HF EPA REG. NO. 34704

Georgia, Louisiana, Mississippi, southeastern Missouri, North Carolina, South Carolina, Tennessee and southern Virginia:
For more complete control of pigweed and seedling johnsongrass, TRIFLURALIN HF may be applied at a broadcast rate per acre of from 1 to 1 1/2 pints on coarse soils, from 1 1/2 to 2 pints on medium soils and 2 pints on fine soils except in the state of Louisiana where 3 pints per acre are recommended on fine soils.

Precaution: Plant soybeans after early season adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when used according to these recommendations.

SOYBEANS—More Complete Weed and Grass Control in Certain Counties Along the Texas Gulf Coast:

For more complete control of those weeds and grasses listed in the TRIFLURALIN HF label in the Texas Gulf Coast Counties of Brazoria, Calhoun, Chambers, Fort Bend, Galveston, Harris, Jackson, Jefferson, Liberty, Matagorda, Orange, Victoria, Waller and Wharton, TRIFLURALIN HF may be applied up to 2 weeks before planting at a broadcast rate of 1 1/2 pints on coarse soils, 2 pints on medium soils and 3 pints on fine soils.

Precaution: Plant soybeans after early season adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when used according to these recommendations.

SOYBEANS, RED RICE CONTROL—Arkansas, Louisiana, Mississippi and Texas only:

Suppression or partial control of red rice in soybeans can be obtained when TRIFLURALIN HF is applied as directed at double the normal rate the first year (not to exceed 4 pints per acre) and at the normal rate the second year. Follow normal TRIFLURALIN HF directions for soil preparation and soil incorporation.

Application: Year 1

Apply and incorporate TRIFLURALIN HF the first year any time in the spring before planting at the following broadcast rates per acre:

Coarse soils	2 pts.
Medium soils	3 pts.
Fine soils	4 pts.
Coarse soils with 2 to 5% organic matter	3 pts.
Soils with 5.1 to 10% organic matter	4 pts.

Application: Year 2

Apply TRIFLURALIN HF the second year at the following normal label broadcast rates per acre:

Coarse soils	1 pt.
Medium soils	1 1/2 pts.
Fine soils	2 pts.
Coarse soils with 2 to 5% organic matter	1 1/2 pts.
Soils with 5.1 to 10% organic matter	2-2 1/2 pts.

If a combination of high organic matter (4 to 10%) and charcoal are present in the soil, apply TRIFLURALIN HF the second year at the following rates labeled for charcoal soils in Arkansas, Louisiana and Mississippi:

Coarse soils	1 1/2 pts. per acre
Medium soils	2 1/2 pts.
Fine soils	3 pts.

*Newly cleared land often contains high organic matter (4 to 10%) and charcoal which result from burning debris. This charcoal and/or organic matter tends to tie up TRIFLURALIN HF and reduce its weed control activity. Higher rates are therefore necessary for satisfactory weed control. Increased rates can cause crop injury if charcoal or a high percentage of organic matter is not present to tie up some of the TRIFLURALIN HF. In the actual windrow or burn row, where a high level of charcoal is present, poor weed control may result even with an increased rate.

Crop Rotation

The program for red rice control in soybeans is a 2-year program. Use the rates listed for first year application and plant soybeans. The second year use the normal TRIFLURALIN HF rates listed for your soil type and charcoal level and plant only those crops for which TRIFLURALIN HF has been registered as a preplant treatment or injury may result. Do not plant rice the second year. Rice may be planted the third year.

Precautions

Plant soybeans after early adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury may occur under adverse cool, wet weather conditions early in the season when TRIFLURALIN HF is used according to these double-rate recommendations.

SOYBEANS—Rhizome Johnsongrass Control: (Eastern United States and the State of Texas)

Commercially acceptable control of rhizome Johnsongrass can be ob-

tained with a double-rate TRIFLURALIN HF program when applied for 2 years in a row.

Soil Preparation—Proper preparation of the soil before application is very important for satisfactory results. Use a chisel plow or similar implement to bring rhizomes to the top of the soil. Then follow with a disc before application to cut the rhizomes into small (2 to 3-inch) pieces. This should also destroy any emerged Johnsongrass.

Application—Choose the one application program that best fits your cultural practices.

Spring Application—Apply TRIFLURALIN HF any time in the spring before planting for 2 years in a row at a broadcast rate per acre of 2 pints on coarse soils; 3 pints on medium soils; 4 pints on fine soils; 3 pints on coarse soils with 2 to 5% organic matter; and 4 pints on soils with 5.1 to 10% organic matter, OR

Fall Application—Apply TRIFLURALIN HF between October 15 and December 31 for 2 years in a row at the same rates as a spring application for the control of rhizome Johnsongrass, OR

Split Application—Apply as directed under both spring and fall applications for 2 years in a row using the following broadcast rates per acre:

	Spring	and	Fall
Coarse soils	1 pt.		1 pt.
Medium soils	1 1/2 pts.		1 1/2 pts.
Fine soils	2 pts.		2 pts.
Coarse soils with 2 to 5% organic matter	1 1/2 pts.		1 1/2 pts.
Soils with 5.1 to 10% organic matter	2 pts.		2 pts.

Incorporation—Deep incorporation is essential to good rhizome Johnsongrass control. Incorporate TRIFLURALIN HF thoroughly with a disc set to cut 4 to 6 inches deep and operate in 2 different directions at 4 to 6 mph.

Cultivation—Some Johnsongrass plants will escape. Timely cultivations during the crop season are necessary to obtain commercially acceptable control. Commercially acceptable control will not be obtained with only 1 year of double-rate TRIFLURALIN HF use.

Precautions: Plant soybeans after early season adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when used according to these recommendations.

In the season following either the 1 or 2-year treatments, plant only those crops for which TRIFLURALIN HF has been registered as a preplant treatment or injury may result.

SOYBEANS—Wild Cane Control:

Wild Cane (Shattercane) can germinate from greater soil depth than most other weed seeds. Several "flushes" or germinating times are common in one season. Commercially acceptable control of wild cane can be obtained with increased rates of TRIFLURALIN HF.

Land Preparation—Work your land to destroy existing grasses and weeds. Thoroughly mix crop residues into the soil to a depth of 4 to 6 inches.

Application—Apply TRIFLURALIN HF before planting at a broadcast rate per acre of 1 pint on coarse soils; 2 pints on medium soils; and 2 1/2 pints on fine soils.

Incorporation—Deep incorporation is essential to good wild cane control. Incorporate (mix) TRIFLURALIN HF thoroughly with a disc only set to cut 4 to 6 inches deep and operate in 2 different directions at 4 to 6 mph.

Cultivation—Cultivations during the crop season will also contribute to control.

Precaution: Plant soybeans after early season adverse weather conditions have passed. Do not plant soybeans deeper than 2 inches. Crop injury in the form of delayed growth may occur under adverse cool, wet weather conditions early in the season when used according to these recommendations.

SOYBEANS—TRIFLURALIN HF/Sencor or TRIFLURALIN HF/Lexone Tank-Mix—Preplant Incorporated:

The TRIFLURALIN HF/Sencor or TRIFLURALIN HF/Lexone tank-mix effectively controls additional weeds than controlled by TRIFLURALIN HF alone (see Weeds Controlled section). Follow normal procedures for soil preparation. The TRIFLURALIN HF/Sencor or TRIFLURALIN HF/Lexone tank-mix should be applied from 2 weeks before planting up to planting in 10 to 40 gallons of water with any low-pressure herbicide sprayer equipped with hydraulic tips and screens no finer than 50 mesh in nozzle and in-line strainers.

Broadcast Rates Per acre:			
	TRIFLURALIN HF	Sencor DF	Lexone DF or
Coarse soils	1 pt.	1/2 pound	
Medium soils	1 1/2 pts.	3/4 pound	
Fine soils	2 pts.	1 pound	
	TRIFLURALIN HF	Lexone 4L or	Sencor 4L
Coarse soils	1 pt.	1/2 pt.	
Medium soils	1 1/2 pts.	3/4 pt.	
Fine soils	2 pts.	1 pt.	

Important: Refer to Sencor and Lexone labels for soil type/rate details. Do not plant any crop other than soybeans within 4 months after treatment. Follow normal TRIFLURALIN HF procedures for incorporation and cultivation.

NOTE: In those areas of the Mid-South where cocklebur is a serious problem, an overlay of Sencor or Lexone may be preferred to the

TRIFLURALIN HF

EPA REG. NO. 34704

TRIFLURALIN HF/Sencor or TRIFLURALIN HF/Lexone tank-mix.

Special Precaution: Applied according to directions and under normal growing conditions, the TRIFLURALIN HF/Sencor or TRIFLURALIN HF/Lexone tank-mix will not harm the treated crop. Over-application may result in crop injury or soil residue. Uneven application or improper soil incorporation of the TRIFLURALIN HF/Sencor or TRIFLURALIN HF/Lexone tank-mix can result in erratic weed control or crop injury. Seeding disease, cold weather, deep planting, excessive moisture, soil pH over 7.5, high salt concentration, or drought may weaken crop seedlings and increase possibility of damage from the TRIFLURALIN HF/Sencor or TRIFLURALIN HF/Lexone tank-mix. Under these conditions, delayed crop development or reduced yields may result. **Warning:** Observe all warnings and limitations on labeling of all products used in mixtures. Sencor may be harmful if swallowed or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing of dust or spray mist. Wash clothing thoroughly with soap and hot water before reuse. Do not contaminate feed or food. Keep out of reach of children.

Do not use the foliage from soybeans treated with the TRIFLURALIN HF/Sencor or TRIFLURALIN HF/Lexone tank-mix for feed or forage. Do not contaminate any body of water nor apply to any area not specified on this label. Do not allow sprays to drift onto adjacent desirable plants.

SOYBEANS—TRIFLURALIN HF pre-plant followed by Sencor or Lexone as an overlay treatment for weed control in soybeans: TRIFLURALIN HF effectively controls certain annual grasses and broad-leaf weeds. See Sencor or Lexone label for additional weeds controlled. Apply TRIFLURALIN HF as a preplant incorporated herbicide. As a separate operation, make a single application of Sencor or Lexone as either a band or broadcast spray during planting or as a separate operation after planting, but before the soybeans emerge. Do not spray Sencor or Lexone over the top of emerged soybeans or injury may result.

Use Directions—Follow directions on the TRIFLURALIN HF, Sencor or Lexone labels for specific instructions regarding each chemical.

Special Precautions—Do not use Lexone or Sencor on Tracy, Semmes, Altona, Vansoy or Coker 102 soybeans as these varieties are sensitive to Lexone or Sencor and injury to the crop may result. See current Lexone or Sencor label for complete information on sensitive varieties.

Do not use treated vines for feed or forage.

Seed must be planted at least 1 1/2 inches below the soil surface but not more than 2 inches before a Sencor or Lexone application.

Do not replant areas treated with Sencor or Lexone to any crop other than soybeans within 4 months after treatment.

Injury to soybeans may occur if Lexone or Sencor is used on soils having a calcareous surface or pH of 7.5 or higher, or if used in conjunction with soil applied organic phosphate pesticides.

Important: Read the TRIFLURALIN HF, Sencor or Lexone labels carefully before using. Note all warnings, precautions and special precautions.

Broadcast Rates Per Acre*	TRIFLURALIN 4EC—Preplant	Lexone or Sencor DF or 4L—Postplant/Preemergence			
		Lexone DF	Sencor DF	Lexone 4L	Sencor 4L
Soil Texture					
Coarse	1 pt.	1/2 lb.	1/2-2/3 lb.	3/4 pt.	3/4-1 pt.
Medium	1 1/2 pts.	1/3-2/3 lb.	1/2-1 lb.	1/2-1 1/4 pts.	3/4-1 1/2 pts.
Fine	2 pts.	1/2-2/3 lb.	2/3-1 1/8 lbs.	3/4-1 1/4 pts.	1-1 3/4 pts.
Mississippi Delta Only	Rate according to soil texture	2/3-1 lb.	1-1 1/3 lbs.	1-1 1/2 pts.	1 1/2-2 pts.

*See Lexone or Sencor use label for complete directions and limitations, including exact rates according to soil organic matter and other factors.

SOYBEANS—TRIFLURALIN HF/Amiben: Amiben may be applied in a band over the soybean row at planting time in fields where TRIFLURALIN HF has been applied as a preplant soil incorporated herbicide. (See Weed Control section for weeds controlled by this treatment.) Or Amiben may be applied several days prior to planting as a broadcast tank-mix with TRIFLURALIN HF. The tank mixture should be used as a spring preplant soil incorporated treatment. The tank-mix improves broadleaf weed, control of species such as smartweed, velvetleaf, and ragweed. For broadcast treatments, incorporate chemicals immediately and thoroughly to an approximate depth of 2 inches with a disc, field cultivator or similar tool set to cut a depth of 4 to 8 inches. Apply Amiben at a rate of 1 gallon (2.0 pounds acid equivalent) per broadcast acre. Apply TRIFLURALIN HF at a rate of 1 1/2 pints for medium soils and 2 pints for fine soils. Do not use on muck or charcoal soils. Read and observe all directions and cautions on the Amiben label.

SUGAR BEETS:

Apply TRIFLURALIN HF as a broadcast, overtop spray when plants are between 2 and 6 inches tall. Exposed beet roots should be covered with soil before application to reduce the possibility of girdling. Set incorporation machinery to throw treated soil toward the plants in the row. Care should be taken that incorporation machinery does not damage the sugar beet taproot.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	1 1/2 pts.	1 1/2 pts.

SUGAR BEETS—Incorporation with a Tine-Tooth Harrow in the States of California, Colorado, Idaho, Kansas, Montana, Nebraska, Oregon, Texas, Utah, Washington and Wyoming Only:

A properly operated tine-tooth harrow (Flexline or Mekroe) can provide adequate incorporation of TRIFLURALIN HF for effective weed control in sugar beets. Operate the tine-tooth harrow 2 times over the field in opposite directions at a speed of 3 to 6 mph and set the harrow to cut 1 to 2 inches deep. Care should be taken to insure that the tine-tooth harrow does not damage the sugar beet taproot.

SUGARCANE—Plant Cane Only

Apply and incorporate TRIFLURALIN HF twice a year at a broadcast rate per acre of 2 to 4 pints for all soil textures. Make the first application in the fall on firmly packed beds immediately after the seed pieces are planted. Make the second application in the spring before or shortly after the cane emerges. Loosen rain-packed beds 2 to 3 inches deep before the spring application. Care should be taken so that incorporation machinery does not damage the seed pieces or emerging shoots.

SUGARCANE—Applications up to Layby for Plant Cane or Ratoon Cane Grown in Louisiana or Texas only:

Apply and incorporate TRIFLURALIN HF at a broadcast rate per acre of 2 to 4 pints for all soil textures. Make the TRIFLURALIN HF application in the spring from before or shortly after the cane emerges up to layby. Make the TRIFLURALIN HF application after the beds have been shaved or false shaved. Loosen rain-packed beds 2 to 3 inches deep before application. Care should be taken so that incorporation machinery does not damage seed pieces or emerging shoots. A rolling cultivator or bed chopper may be used to incorporate TRIFLURALIN HF layby applications in sugarcane on all soil textures. Follow normal incorporation directions for the rolling cultivator. Set bed chopper to cut 3 to 4 inches deep and operate 2 times at 4 to 6 mph.

SUGARCANE—Raouigrass Control in Louisiana only:

Apply and incorporate TRIFLURALIN HF on either plant or ratoon cane at a broadcast rate per acre of 4 pints for all soil textures. Make the TRIFLURALIN HF application in the spring from before or shortly after the cane emerges up to layby. Make the TRIFLURALIN HF application after the beds have been shaved or false shaved. Loosen rain-packed beds 2 to 3 inches deep before application. Care should be taken so that incorporation machinery does not damage seed pieces or emerging shoots. A rolling cultivator or bed chopper may be used to incorporate TRIFLURALIN HF layby applications in sugarcane on all soil textures. Follow normal incorporation directions for the rolling cultivator. Set bed chopper to cut 3 to 4 inches deep and operate 2 times at 4 to 6 mph.

SUGARCANE—For control of most annual grasses, including guineagrass (Post-Plant in Hawaii Only):

Surface apply TRIFLURALIN HF after planting (for plant cane) or after harvesting (for ratoon cane), at a broadcast rate per acre of 6 to 8 pints for all soil texture. Apply TRIFLURALIN HF from shortly before or after cane emergence until layby. In ratoon cane, excess crop residue should be removed before application. If large amounts of crop residues are present, TRIFLURALIN HF will not be effective. Apply just before anticipated rainfall or irrigate immediately after application.

SUNFLOWER:

Apply and incorporate TRIFLURALIN HF before planting.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2-2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

TOMATOES:

For Direct-seeded tomatoes apply TRIFLURALIN HF at blocking or thinning. Apply as a directed spray to the soil between the rows and beneath the plants and incorporate.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pt.
Medium soils	1 1/2 pts.	1 1/4-1 1/2 pts.
Fine soils	2 pts.	1 1/2 pts.
Soils with 2 to 5% organic matter	1 1/2 pts.	1 1/2-2 pts.
Soils with 5.1 to 10% organic matter	2 pts.	2 pts.

For Transplant tomatoes apply and incorporate TRIFLURALIN HF before transplanting. Do not apply after transplanting.

	Eastern U.S.	Western U.S.
Coarse soils	1 pt.	1 pts.

TRIFLURALIN HF

EPA REG. NO. 34704

Medium soils 1 1/2 pts. 1 1/4-1 1/2 pts.
Fine soils 2 pts. 1 1/2 pts.
Soils with 2 to 5% organic matter 1 1/2 pts. 1 1/2-2 pts.
Soils with 5.1 to 10% organic matter 2 pts. 2 pts.

TREES AND VINEYARDS: (Eastern U.S. only)

For *New Plantings of Vineyards, Citrus and Pecan Trees* apply and incorporate TRIFLURALIN HF before planting at a broadcast rate per acre of 1 pint on coarse soils; 1 1/2 pints on medium soils; 2 pints on fine soils; 1 1/2 pints on fine soils with 2 to 5% organic matter; and 2 pints on soils with 5.1 to 10% organic matter.

For *Non-Bearing Established Plantings of Citrus and Pecan Trees and Bearing Plantings of Grapefruit, Lemon, Orange, Pecan, Tangelo, Tangerine Trees* apply TRIFLURALIN HF at a broadcast rate per acre of 2 to 4 pints for all soil textures. In these established plantings, apply as a directed spray to the soil around the trees and use incorporation methods not injurious to the trees.

NOTE: If crops are planted between the trees, label directions for those specific crops apply to the area which is interplanted. For continued weed control in citrus trees, apply TRIFLURALIN HF 2 times a year at an interval of approximately 4 to 6 months.

TREES AND VINEYARDS: (Western U.S. only)

For *New Plantings of Almond, Apricot, Citrus, Nectarine, Peach, Pecan and Walnut trees* apply and incorporate TRIFLURALIN HF before planting at a broadcast rate per acre of 1 pint on coarse soils; 1 1/4 to 1 1/2 pints on medium soils; 1 1/2 pints on fine soils; 1 1/2 to 2 pints on soils with 2 to 5% organic matter; and 2 pints on soils with 5.1 to 10% organic matter.

For *New Plantings of Vineyards* apply and incorporate TRIFLURALIN HF before planting at a broadcast rate per acre of 1 to 1 1/2 pints on coarse soils; 1 1/2 to 3 pints on medium soils and 3 to 4 pints on fine soils or soils with 2 to 10% organic matter. Do not use more than 2 pints per acre on heat-treated vines.

For *Post-Plant Applications on Bearing or Non-Bearing Established Plantings of Vineyards, Almond, Apricot, Grapefruit, Lemon, Nectarine, Orange, Peach, Pecan, Plum, Prune, Tangelo, Tangerine and Walnut Trees* apply TRIFLURALIN HF at a broadcast rate per acre of 2 to 4 pints for all soil textures. In these established plantings, apply as a directed spray to the soil around the trees or vines and use incorporation methods not injurious to the trees or vines. Do not apply to vineyards within 60 days of harvest.

NOTE: If crops are planted between the trees or vines, label directions for those specific crops apply to the area which is interplanted. For continued weed control in citrus trees, apply TRIFLURALIN HF 2 times a year at an interval of approximately 4 to 6 months.

TREES AND VINEYARDS—Rhizome Johnsongrass Control: (Western U.S. Only)

Commercially acceptable control of rhizome Johnsongrass can be obtained with post-plant applications in Bearing and Non-Bearing established plantings of Vineyards, Almond, Apricot, Grapefruit, Lemon, Nectarine, Orange, Peach, Pecan, Tangelo, Tangerines and Walnut trees with a TRIFLURALIN HF program when applied for 2 years in a row.

Soil Preparation—Work the soil thoroughly to bring the rhizomes nearer the surface.

Application—Apply TRIFLURALIN HF at a broadcast rate per acre of 4 pints on all soil textures each year for 2 years in a row. Do not apply to vineyards within 60 days of harvest.

Incorporation—Incorporate TRIFLURALIN HF thoroughly with a disc set to cut 4 to 6 inches deep and operate 2 times at 4 to 6 mph.

Cultivation—Some Johnsongrass plants will escape. Timely cultivations are necessary to obtain commercially acceptable control. Commercially acceptable control will not be obtained with only 1 year of TRIFLURALIN HF use.

Precautions: Do not use the 2-quart rate on new plantings as injury may result. Do not interplant orchards or vineyards with other crops. If TRIFLURALIN HF-treated vineyards and orchards are diverted to other crop uses, plant only those crops for which TRIFLURALIN HF has been registered as a pre-plant treatment.

TREES AND VINEYARDS—Field Bindweed Control in Vineyards, Almond, Apricot, Grapefruit, Lemon, Nectarine, Orange, Peach, Pecan, Tangelo, Tangerine and Walnut Trees in California only: For the control of field bindweed in the state of California, apply TRIFLURALIN HF at a broadcast rate of 4 pints per acre on all soil textures. TRIFLURALIN HF must be applied in the spring with a specially designed spray blade which applies a thin concentrated layer at a soil depth of 4 to 6 inches. The layer of TRIFLURALIN HF prevents bindweed shoots from emerging.

Land Preparation—Destroy all weeds and grasses with soil tillage before applying TRIFLURALIN HF. This tillage is necessary to prevent trash from interfering with the operation of the spray blade.

Equipment—This operation requires a spray blade capable of running 4

to 6 inches below the surface of the soil. The spray blade should be equipped with nozzles located under the blade and directed so that the TRIFLURALIN HF spray will be trapped under the soil which is flowing over the blade as it is pulled through the soil. Use a sufficient number of nozzles with spacing to completely and uniformly apply TRIFLURALIN HF underground in a thin horizontal layer.

Application—Apply TRIFLURALIN HF in 40 to 80 gallons of water per acre. Operate the spray blade at a depth of 4 to 6 inches.

Precaution: Some soils develop cracks as they dry after rainfall or irrigation. Field bindweed may emerge if the cracks extend through the TRIFLURALIN HF layer. Prevent or eliminate cracks by shallow disking or other tillage. Avoid deep tillage which disturbs the subsurface layer. Cultivation or tillage also aids the control of germinating seeds.

WHEAT (SPRING), DURHAM AND BARLEY—Postplant incorporated application for control of foxtail (pigeongrass): Plant 2 to 3 inches deep in a seedbed of good tillth. Make TRIFLURALIN HF application after seeding but before crop emergence. Incorporate 1 to 1 1/2 inches deep by use of flex-tine or diamond harrows operated twice in different directions at a minimum speed of 5 mph. Combine herbicide application and first incorporation when possible. Both incorporations must be completed in 24 hours following application.

Apply and incorporate at the following broadcast rates per acre:

Coarse soils 1 pint
Medium soils 1 pint
Fine Soils 1 1/2 pints

WHEAT (SPRING), DURHAM, BARLEY—FALL APPLICATION—Preplant incorporated application for control of foxtail (pigeongrass): This fall application of TRIFLURALIN HF is for crops to be planted the following spring. Fields should not have excessive trash and may have been fallowed or pre-tilled according to local cropping practices. Initial incorporation must be made within 24 hours of application. A second incorporation must be made prior to planting to uniformly distribute treated soil and eliminate emerged weeds.

Apply and incorporate at the following broadcast rates per acre:

Coarse soils 1 pint
Medium soils 1 pint
Fine soils 1 1/2 pints

Fall Incorporation Pass—Use any of the following implements:

1. Chisel plow (3 rows of up to 18-inch sweeps on 12-inch centers or less): Sweeps must be staggered so as to turn all soil. Operate at depth of 4 to 5 inches and speed of 4 to 6 mph.
2. Tandem disc: Operate at depth of 3 to 4 inches and speed of 4 to 6 mph.
3. Field cultivator (3 or 4 rows of sweeps with C- or S-shaped shanks spaced 7 inches or less apart): Sweeps must be staggered so as to turn all soil. Operate at depth of 3 to 4 inches and speed of 5 mph minimum.

Spring Incorporation Pass—The disc or field cultivator may be used, but the chisel plow is not recommended. The spring pass implement must operate at a more shallow depth than the fall pass implement.

Planting Depth—Seed should be placed at approximately 2 inches deep. Note: The application described may result in a reduction of stand, however, a slight stand reduction usually does not affect yield.

BARLEY, SPRING SEEDED-SPRING APPLICATION PREPLANT INCORPORATED FOR FOXTAIL (PIGEONGRASS) CONTROL (For use in Minnesota, North Dakota and South Dakota)

Apply Trifluralin HF as a preplant incorporated treatment prior to planting spring seeded barley. This product may be applied to ground that has a manageable level of crop residue or has been fallowed or pre-tilled. The first incorporation is required within 24 hours after application. The second incorporation required prior to planting to destroy emerged weeds and to insure even distribution of this product in the soil surface. Broadcast Application Rate/Acre: Apply at a rate of 1.0 pint per acre for all textures regardless of organic matter content.

Incorporation: Recommended incorporation tools include the chisel plow (first incorporation pass only), tandem disc and field cultivator. Refer to "Incorporation Directions" section of this label for details on operation of incorporation equipment.

Planting Directions: Barley should be seeded approximately 2 inches deep.

Precautions:

- Carefully read and follow precautionary information before applying Trifluralin HF.
- While use of this weed control practice may result in stand reduction, slight stand reductions do not normally affect yield.

WHEAT (SPRING), DURHAM AND BARLEY—TRIFLURALIN HF/FAR-GO TANK MIX—Postplant incorporated application for control of foxtail (pigeongrass) and wild oat:

Plant 2 to 3 inches deep in a seedbed of good tillth. Make herbicide application after seeding, but prior to crop emergence. Incorporate 1 to 1 1/2 inches deep by use of flex-tine or diamond harrows operated twice in different directions at a minimum speed of 5 mph. Combine herbicide application and first incorporation when possible. If not possible, incorporate immediately after application.

Apply and incorporate at the following broadcast rates per acre:

SOIL TEXTURE	TRIFLURALIN 4EC	Far-Go	
	Barley, Durham, Spring Wheat	Durham, Spring Wheat	Barley
Coarse soils	1 pt.	2 1/2 pts.	2 pts.
Medium soils	1 pt.	2 1/2 pts.	2 pts.
Fine soils	1 1/2 pts.	2 1/2 pts.	2 pts.

Important: Do not overapply as crop injury may result. Read Far-Go label carefully before using.

11/16

TRIFLURALIN HF

EPA REG. NO. 34704

WHEAT (WINTER)—TRIFLURALIN HF for preplant pre-emergence control of cheatgrass and other weeds in winter wheat grown in Washington, Oregon, Idaho, and Montana:
When applied as directed, TRIFLURALIN HF will provide effective pre-emergence control of cheatgrass and a number of other annual grasses and broadleaf weeds controlled by TRIFLURALIN HF in winter wheat grown in Washington, Oregon, Idaho, and Montana. The growth, development and yield of winter wheat will not be adversely affected, provided the seed is placed below the zone of soil treated with TRIFLURALIN HF.

Broadcast Rates Per Acre:

Apply TRIFLURALIN HF any time during a period from 3 weeks up to immediately prior to planting. Broadcast TRIFLURALIN HF at the following rates per acre according to soil texture.

Soil Texture	TRIFLURALIN HF
Coarse	1 1/2 pts.
Medium	1 1/2 pts.
Fine	2 pts.

Incorporation Directions—Shallowly incorporate TRIFLURALIN HF into the soil with a flexible tine-tooth harrow (Flexdine, Melroe) set to cut 1 to 2 inches deep. Operate the equipment in 2 different directions at a speed 3 to 6 miles per hour. The first incorporation must be within 24 hours after application. The second incorporation may be done at any time but before planting. Do not till the soil with a disc after the TRIFLURALIN HF has been applied and incorporated with a flexible tine harrow.

Seeding Directions—Use only a deep furrow or semi-deep furrow drill that will be sure to place the seed below the zone of soil into which TRIFLURALIN HF has been incorporated.

Wheat planted in direct contact with TRIFLURALIN HF treated soil may suffer crop injury in the form of delay in emergence and development.

WHEAT (WINTER)—Fallow-soil application of TRIFLURALIN HF for weed control in winter wheat grown in Washington and Oregon:
Uniformly applied TRIFLURALIN HF at the recommended rate and shallowly incorporated into fallow soil as much as four months ahead of planting time, will effectively control cheatgrass and certain annual grasses and broadleaf weeds in winter wheat grown in Washington and Oregon. The growth, development, or yield of winter wheat will not be adversely affected, provided the seed is placed below the zone of soil treated with TRIFLURALIN HF with deep or semi-deep furrow-drill.

Broadcast directions and application rates per acre:

Soil Texture	TRIFLURALIN HF
Coarse	1 1/2 pts.
Medium	1 1/2 pts.
Fine	2 pts.

Apply any time from May to September prior to the fall planting of winter wheat.

Incorporation—Shallowly incorporate TRIFLURALIN HF into the soil with a flexible tine-tooth harrow (also called Flexdine or Melroe) set to cut 1 to 2 inches deep and operated at 3 to 6 mph. Thorough incorporation requires two passes of the equipment in different directions over the field. The first pass must be made within 24 hours after application. The second pass may be delayed for several weeks but should be made before seeding. Do not till the soil with a disc after TRIFLURALIN HF has been applied with a flexible tine harrow.

Precaution: Use only deep furrow or semi-deep furrow drills. Place seed below the zone of soil into which TRIFLURALIN HF has been incorporated. Do not plant wheat directly into the zone of soil treated with TRIFLURALIN HF as injury to the crop or a delay in its emergence and development may occur.

WINTER WHEAT — Partial Control or Suppression of Annual Brome Species (Cheatgrass, Downey Brome, Japanese Brome, Hairy Chess) and Jointed Goatgrass in Colorado, Kansas, Nebraska and Wyoming.

Apply Trifluralin 4EC as a preplant incorporated treatment anytime from three (3) weeks before planting up to immediately prior to planting.

Broadcast rates per acre:

Soil Texture	Trifluralin 4EC
Coarse	1-1 1/2 pts.
Medium	1-1 1/2 pts.
Fine	1 1/2 pts.

Use higher rate range where heavy weed populations are anticipated or where medium to high crop residues are present.

Incorporation and Planting Directions

Incorporated with tillage equipment (flexible tine-tooth harrow or springtooth harrow such as Flex-Tyne® or Crust-Buster®) that mixes the soil no more than 1-2 inches deep.

The grain drill (double disc drill or hoe drill) can serve as the incorporation method. Do not use discs, under cutters or heavy field cultivators for incorporating. Incorporate once within 24 hours after application. Use a drill that will place the seed below soil which has been incorporated with Trifluralin 4EC.

One pass incorporation is adequate. Where the grain drill is used as the incorporation tool, mounting a springtooth harrow in front of the drill to

aid incorporation can enhance performance. THE WHEAT MUST BE SEEDED BELOW THE TRIFLURALIN HF TREATED SOIL OR CROP INJURY MAY RESULT. The wheat seed should be placed at least 1 1/2 inches deep.

Precautions:

- Crop injury in the form of delayed emergence and development result from planting wheat in direct contact with treated soil.
- Do not use under cutters, field cultivators, chisel plows or disc as incorporation tools. Any implement that incorporates Trifluralin 4EC deeper than the seeded wheat will contribute to crop injury.
- Use of seeding equipment that does not place the seed below the treated soil layer will result in crop injury.
- Use of Trifluralin in accordance with the label may result in some crop stand reduction but does not normally adversely affect yield.
- Heavy rainfall prior to wheat emergence can cause soil compaction and soil crusting resulting in delayed emergence, stand reduction, stunting and yield loss.

NOTICE OF WARRANTY

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

- SENCOR is a Reg. T.M. of Parent Co. of Farbenfabriken Bayer GmbH, Leverkusen.
- KARMEX and LEXONE are Reg. T.M.'s of E.I. duPont de Nemours and Co., Inc.
- EPTAM is a Reg. T.M. of the ICI Group.
- AMIBEN is a Reg. T.M. of Rhone-Poulenc.
- COTORAN and CAPAROL are Reg. T.M.'s of Ciba-Geigy Corp.
- FAR-GO is a Reg. T.M. of Monsanto Co.

FORMULATED FOR
PLATTE CHEMICAL CO.
150 SO. MAIN STREET FREMONT, NEBRASKA 68025-5697