box: "The requirements in this box apply to uses of this product are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or

Date:

greenhouses."

Cynthia Giles-Parker Product Manager (22) Fungicide Branch

Registration Division

EPA Form 8570-6

continued on page 2
Signature of Approving Official:

page 2 EPA Reg. No. 34704-802

4. Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

ACCOST 1G

ACTIVE INGREDIENT:

1-(4-Chlorophenoxy)-3,3-dimethyl-1(1 H-1,2,4-triazol-1-yl)-2-butanone 1%
INERT INGREDIENTS: 99%
100%

Turf and Sod Production Fungicide Contains Bayleton®

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUCION AL USUARIO: Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

> EPA REG. NO. 34704- INE EPA EST. NO. _____ NET WEIGHT:____

BAYLETON is a Registered TM of Bayer A.G.

EXP 04/97

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor, or dust. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and Other Handlers Must Wear:

Long-Sleeved Shirt and Long Pants

Waterproof Gloves

Shoes plus Socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash-thoroughly and put on clean clothing.

STATEMENTS OF PRACTICAL TREATMENT

If swallowed: Administer water freely and induce vomiting by giving one dose (½ oz or 15 mL) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose, If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Get medical attention immediately.

If on skin: Wash skin immediately with soap and warm water, if imitation occurs, get medical attention.

If In eyes: Wash with plenty of water for at least 15 minutes. Get medical attention.

ACCEPTED
with COMMENTS
In EPA Letter Dated
JAN 26 1998

Under the Federal Insecticide,

If Inhaled: Remove to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. Apply this product only as specified on this label. Do not make applications when weather conditions lavor drift from target area. This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labelling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

Coveralis Waterproof GLoves Shoes plus Socks

NON AGRICULTURAL USE REQUIREMENTS

Restricted Entry Interval: When used on commercial sod (sod farms), follow PPE and reentry instructions in the Agricultural Use Requirements section of this label. For use on other turf areas, do not allow public use of treated areas during application or until dust has settled.

HOW TO USE

This product is absorbed rapidly and works systemically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices. Apply when the turf is fully established and actively growing. Applications should be applied at recommended intervals to maintain disease control. Do not use on crops grown for food or forage.

34704-802

ACCUSIO 1G **EPA REG. NO. 34704-**

RECOMMENDED APPLICATIONS

TURF GRASS DISEASE CONTROL					
DISENSE	Lb. of ACCOST 1 G		REMARKS		
	Per 1000				
Calles Cash	Preventive	Curative			
Dollar Spot	1.5	}	PREVENTIVE PLATE: Apply		
(Scierotinia			recommended rate on 30 day		
homoeocarpa)		-	intervals. Protective activity of		
			this fungicide may extend for as		
		}	long as 60 days dependent		
			upon environmental conditions.		
	0.75		PREVENTATIVE RATE: Apply		
	j '		recommended rate on 14 day		
	i		intervals. Protective activity of		
	1		this fungicide may extend for as		
	<u>'</u>	}	long as 30 days dependent		
			upon environmental conditions.		
	-	3	CURATIVE RATE: To control		
		_	existing infections, apply the		
			curative rate. Re-application		
	Į I		should be no closer than 14		
	į		days. When infections have		
			been controlled, refer to the		
	!	i i	preventive rate schedule for		
	i	}			
Copper Spot	'		rates and timing. PREVENTIVE RATE: Apply at		
	1.5	3			
(Gloeocercospora	1	\ \ \	15 to 30 day intervals. When		
sorghi)	Į.		environmental conditions favor		
Powdery Mildew	!		light to moderate disease devel-		
(Epysiphe	-	[[opment, use a longer interval.		
graminis)			Protective activity of this fungi-		
Corticium Red	,	1	cide can be greater than 30		
Thread]))	days depending on environmen-		
(Laetisara			tal conditions.		
fuciformis)	;	(l			
Rusts	1		CURATIVE RATE: To control		
(Puccinia spp.)		l i	existing infections, apply the		
Brown Patch/	,	l i	curative rate. Re-application		
Rhizoctonia Blight	1	1	should be no closer than 14		
(Ahizoctonia	,	l	days. When infections have		
solani)	!		been controlled, refer to the		
(Suppression)		ì	preventive rate schedule for		
(Deppicasion)	1		rates and timing.		
Anthracnosa	3		PREVENTIVE RATE: Apply at		
(Colletotrichum	3	1	30 day intervals and repeat as		
1	1	!]			
graminicola)	1	[[necessary for seasonal control.		
			Dependent upon environmental		
	-		conditions, residual control may		
		ļ <u>-</u>	be extended to 45 days.		
	1.5	- :	PREVENTIVE RATE: Apply at		
	í		21 day intervals and repeat as		
		{	necessary for seasonal control.		
	i i		Dependent upon environmental		
	1	1	conditions, residual control may		
		L	be extended to 30 days.		
	•	3	CURATIVE RATE: To control		
	j		existing infections, apply the		
	1	1	curative rate. Re-application		
	}	1	should be no closer than 14		
	i	1	days. When infections have		
	1	<u> </u>	been controlled, refer to the		
	1	1	preventive rate schedule for		
	}	i	rates and timing.		
			I I I I I I I I I I I I I I I I I I I		

DISEASE	Lb. of ACCOST 1G Per 1000 Sq. Ft.* Preventive Rates Only	REMARKS
Stripe Smut (Ustilago striiformis)	3	Make a single application in spring as grass growth begins.
Fusarium Bilight (Fusarium culmorum) (Fusarium poae) Summer Patch (Magnaporthe poae) (Except California)	3	Apply first application in mid- June or 30 days prior to time this blight normally becomes evident. Applications should be made on 30 day intervals.

DISEASE	Lb. of ACCOST 1G Per 1000 Sq. Ft."	REMARKS
Fusarium Blight (Fusarium culmorum) (Fusarium poae) Summer Patch (Magnaporthe poae) (California only)	3 to 6	Apply first application in mid- June or 30 days prior to time this blight normally becomes evident. Applications should be made at the 3 to rate on 15 day intervals or at the 5 to rate on 30 day intervals.
Zoysia patch. Large patch of zoysia (Fihizoctonia solani)	3 to 6	Make first application in early fall (mid-Septemper to mid-October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
Gray Snow Mold/ Typhula Blight (Typhula incarnata) (Except California)	6	Apply in fall 30 days pnor to turf dormancy. If turf breaks dormancy during winter months a second application should be made. Do not apply over snow cover, or when turf is dormant.
Pink Snow Mokd/ Fusarium Patch (Microdochium nivalis) Southern Blight (Sclerotium rolfsii) (Except California)	3 to 6	Apply before conditions favorable for infection occur. Re-application should be made as needed at a 60 to 90 day interval. Do not apply over snow cover, or when turf is dormant. Use higher rate in areas with a history of severe disease damage.

*Apply the specified amount of ACCOST 1G with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq ft. Make application when turf is dry and following the mowing operation. Do not mow until a minimum of .25 inch of water has been applied to treated area. Do not use clippings for animal feed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Household: Do not reuse empty container. Securely wrap original container in several layers of newspaper and put in trash collection.

Pesticide Disposal (Except Household): Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal (Except Household): Do not use container in connection with food, feed, or drinking water. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities by burning. If burned, stay out of smoke. Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked stor-

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Material that cannot be used as directed should be disposed of as directed above. In spill or leak incidents, keep unauthorized people away. You may contact CHEMTREC at 800-424-9300 for decontamination procedures or any other assistance that may be necessary.

NOTICE

PLATTE CHEMICAL CO. WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTION TIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT, PLANT INJURY, INEFFEC-TIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE.
IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT
DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, ALL SUCH
RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

PLATTE CHEMICAL CO.

150 SO. MAIN STREET

FREMONT, NEBRASKA 68025-5697