

34704-805

04/02/2001

1/6



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
34704-805

Date of
April 2, 2001

NOTICE OF PESTICIDE:
 Registration, Reregistration

(under FIFRA, as amended)

Term of Issuance:
Conditional

Name of Pesticide Product:
BioCover MLT

Name and Address of Registrant (include ZIP Code):

Platte Chemical Company
419 18th Street
P.O. Box 667
Greeley, CO 80632-0667

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 34704-805".
3. Submit three copies of the final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "accepted with comments" is enclosed for your records.

Signature of Approving Official:
Akiva Abramovitch, Ph.D. *Akiva Abramovitch*
Chemist
Insecticide-Rodenticide Branch
Registration Division (7505C)

Date: April 2, 2001

APR 2 - 2001

ACCEPTED with COMMENTS In EPA Letter # 34704-805

BioCover MLT

for the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 34704-805

APR 2 2001

ACTIVE INGREDIENT:	BY WT.
Petroleum Oil	98.00%
INERT INGREDIENTS:	2.00%
TOTAL	
	100.00%
Unsulfonated Residue	99.00%
Aromatic Composition by ASTM D2140	0.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See Below For Additional Precautionary Statements.

FIRST AID

Have the product container label with you when calling a poison control center or doctor or going for treatment.

If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to by a poison control center or doctor.
If in eyes:	<ul style="list-style-type: none"> Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Petroleum distillate poses aspiration pneumonia hazard. Only minor irritation should be expected from any type of exposure other than pulmonary aspiration. Ingestion may result in some gastroenteritis and mild diarrhea. Since systemic toxicity is not expected, the added risk of aspiration from attempts to induce vomiting or lavage are deemed not warranted.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:
1-800-228-5635, EXT. 136, OR CALL COLLECT, 612-851-8180, EXT. 136.

EPA REG. NO. 34704-INL
EPA EST. NO. _____

NET CONTENTS _____ GAL (_____ L)

IHT 10Y00

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION

Harmful if swallowed or absorbed through skin or inhaled. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. This product is a potential skin sensitizer. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment: Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves, such as barrier laminate, nitrile rubber, butyl rubber, neoprene or viton, and shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that

meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:
 Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Apply this product only as specified on the label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over long-sleeved shirt and long pants, shoes plus socks and chemical-resistant gloves, such as barrier laminate, nitrile rubber, butyl rubber, neoprene or viton.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal. In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call collect: CHEMTREC (800) 424-9300. To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dust. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers. **Plastic:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

APPLICATIONS

Total coverage of all parts of tree is essential for maximum insect control. Apply as directed to thoroughly drench all limbs, twigs, and leaves without significant runoff.

BioCover™ MLT
EPA REG. NO. 34704-INL

This product may be used as a carrier for appropriate insecticides and fungicides which are registered for use on deciduous fruits and nuts and which can be applied to give a uniform spray. Consult your local Agricultural Extension Service.

DO NOT COMBINE WITH SULFUR SPRAYS AND DO NOT APPLY spray oil within 30 days before or after an application of sulfur except in the Northwest (Oregon and Washington) where it may be used in combination on Pears as a Post Harvest (after all fruit is off the tree), Dormant and Delayed Dormant (before the scales slip, or at or prior to bud swell) treatment and on Apples prior to or up to the Delayed Dormant (½ inch green) period.

Do not apply to Captan within 90 days after or 60 days before this oil. If in doubt, spray a portion of one tree previously treated with Sulfur or Captan using 1 quart to 25 gallons water. Burn will show in several days if sufficient time has not elapsed. Avoid application of oil sprays just prior to or during hot weather (95°F), rainy weather or drying winds. Apply when temperature is above 32°F.

DO NOT USE THIS MATERIAL IF IT DOES NOT EMULSIFY.

DO NOT MIX with insecticides or fungicides incompatible with oil. Consult your State Agricultural Extension Service for specific compatibilities.

DO NOT APPLY to trees weakened by disease, drought, drying winds, or high nitrogen applications.

DO NOT USE this product when temperatures are above 95°F.

Use only in spray machines with powerful positive agitation. Keep agitator running at all times during spray operations. This product is formulated as a "fast break" spray oil and will not stay in solution unless agitated constantly.

Do not apply this product through any type of irrigation system.

See specific instructions for concentrate and aerial application.

ORCHARD CONCENTRATE APPLICATION: Apply according to rates given in the table for each crop. All quantities of spray solution are for mature trees and vines with normal foliage cover. Quantities should be reduced proportionally for smaller immature trees and vines. Use only on crops listed on this label. Special concentrate spray equipment is necessary for this application.

NOTE: Use efficient equipment of the proper type. Avoid spraying during or immediately prior to hot or freezing weather (over 95°F. or under 32°F.), hot or winds, rain or other unsuitable conditions; avoid overspraying or double spraying. Plants should be sprayed only when in vigorous condition and when the moisture condition is suitable.

Fill spray tank with ½ the required amount of water. Start agitation. Add total amount of spray oil. Continue operation of agitator while adding balance of water and while spraying.

APPLES AND PEARS:

PESTS	APPLICATION RATE & TIMING	COMMENTS
ON APPLES AND PEARS: European Fruit Lecanium Scale Olive Scale San Jose Scale European Red Mite Eggs	DORMANT OR DELAYED DORMANT: Use 1 to 2 gals. per 100 gals. of water at 400 to 800 gals. per acre. (Concentrate Spray: Use 6 to 8 gals. per acre in a minimum of 20 gals. water per acre.)	
Two Spotted Spider Mite Pacific Red Mite European Red Mite	FOLIAR SPRAY: Use 1 to 1½ gal. per 100 gals. water 400 to 800 gals. per acre. (Concentrate Spray: Do not use more than 4 gals. per acre in a minimum of 20 gals. water per acre.)	
ON APPLES ONLY: Rosy Apple Aphid Eggs Apple Aphid Eggs	DORMANT OR DELAYED DORMANT: Use 1 to 2 gals. per 100 gals. of water at 400 to 800 gals. per acre. (Concentrate Spray: Use 6 to 8 gals. per acre in a minimum of 20 gals. water per acre.)	
Fruit Tree Leafroller Apple Red Bug Scurfy Scale	GREEN TIP TO DELAYED DORMANT: Use 1 to 3 gals. per 100 gals. of water at 400 to 800 gals. per acre. (Concentrate Spray: Use 6 to 12 gals. per acre in a minimum of 20 gals. water per acre.)	
European Red Mite		Delayed dormant preferred
San Jose Scale Forbes Scale	DORMANT, GREEN TIP TO DELAYED DORMANT: Use 1 to 2 gals. per 100 gals. of water at 400 to 800 gals. per acre. (Concentrate Spray: Use 6 to 8 gals. per acre in a minimum of 20 gals. water per acre.)	
ON PEARS ONLY: Pear Psyllid	DORMANT: Use 1 to 2 gals. per 100 gals. of water at 400 to 800 gals. per acre. (Concentrate Spray: Use 6 to 8 gals. per acre in a minimum of 20 gals. water per acre.)	Apply before eggs are laid.
Fruit Tree Leafroller Pear Leaf Blister Mite	DORMANT: Use 1 to 3 gals. per 100 gals. of water at 400 to 800 gals. per acre. (Concentrate Spray: Use 6 to 12 gals. per acre in a minimum of 20 gals. water per acre.)	

For Aerial Application: Dormant and Delayed Dormant Application Only: Apply 3 to 5 gals. in a minimum of 20 gals. of water per acre. Apply by air only when soil conditions do not permit regular ground application.

BioCover™ MLT
EPA REG. NO. 34704-INL

WALNUTS:

PEST	APPLICATION RATE & TIMING	COMMENTS
Black Scale Calico Scale European Fruit Lecanium Scale Frosted Scale Olive Scale San Jose Scale European Red Mite Eggs Walnut Aphid Eggs	SUMMER STAGES: Use 4 to 6 gals. per acre in 200 to 500 gallons of water per acre.	Use only on trees that have not suffered from lack of moisture at any time during the growing season. Do not apply to orchards where soil moisture is low. Trees must be fully dormant. Do not apply after husk split.
Spider Mites Soft Scales	SUMMER FOLIAR (when leaves fully expanded): Use 4 to 6 gals. per acre in 200 to 500 gals. water per acre.	

GRAPES:

PEST	APPLICATION RATE & TIMING	COMMENTS
Mealybug	DORMANT: Use 2 to 3 gals. plus the appropriate label rate of insecticide registered for use on grapes in 200 to 300 gals. of water per acre.	Do not apply over 300 gals. of water per acre.

OLIVES:

PEST	APPLICATION RATE & TIMING	COMMENTS
Olive (Parlatoria) Scale Oleander Scale Black Scale	POSTBLOOM THROUGH AUGUST AND POST HARVEST: Use 1 to 1½ gals. per 100 gals. of water at 400 to 1500 gals. per acre.	

ALMONDS, APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS AND PRUNES: (Directions for use on Plums and Prunes in California's Sacramento Valley can be found in the table following this one):

PEST	APPLICATION RATE & TIMING	COMMENTS
Black Scale European Fruit Lecanium Scale San Jose Scale Olive Scale Frosted Scale Aphid Eggs European Red Mite Eggs Brown Mite Eggs Fruitree Leafroller Eggs	DORMANT AND DELAYED DORMANT: Use 1 to 2 gals. per 100 gals. of water at 300 to 600 gals. per acre. (Concentrate Spray: Use 6 to 8 gals. per acre in a minimum of 20 gals. of water per acre.)	Preferred times of application for European Fruit Lecanium Scale and European Red Mite Eggs is Green Tip to Delayed Dormant, and for San Jose Scale is Dormant, Green Tip to Delayed Dormant.
Terrapin Scale	DELAYED DORMANT: Use 2 to 3 gals. per 100 gals. of water at 300 to 600 gals. per acre. (Concentrate Spray: Use 8 to 12 gals. per acre in a minimum of 20 gals. of water per acre.)	
Cottony Peach Scale	DORMANT: Use 2 to 3 gals. per 100 gals. of water at 300 to 600 gallons per acre. (Concentrate Spray: Use 8 to 12 gals. per acre in a minimum of 20 gals. of water per acre.)	
Two-spotted Spider Mite Pacific Mite European Red Mite Brown Mite	FOLIAR SPRAY: Use ½ to 1½ gals. per 100 gals. of water at 300 to 600 gals. per acre. (Concentrate Spray: Do not use more than 4 gals. per acre in a minimum of 20 gals. water per acre.)	Certain varieties of plums and prunes may be injured by oil sprays, check for tolerance prior to treatment.
White Peach Scale on Peaches: Only for use in Georgia, Florida, South Carolina, North Carolina, and Alabama.	DORMANT: Use 1½ to 2 gals. per 100 gals. of water at 300 to 600 gals. per acre. (Concentrate Spray: Use 6 to 8 gals. per acre in a minimum of 20 gals. of water per acre.)	Apply before buds begin to swell in the spring. Make 2 applications 7 to 14 days apart, according to local recommendations, during the dormant period.

For Aerial Application: Dormant and Delayed Dormant Application Only: Apply 3 to 5 gals. in a minimum of 20 gals. of water per acre. Apply by air only when soil conditions do not permit regular ground application.

BioCover™ MLT
EPA REG. NO. 34704-INL

PLUMS AND PRUNES: (Sacramento Valley of California Only)

PEST	APPLICATION RATE & TIMING	COMMENTS
Black Scale European Fruit Lecanium Scale San Jose scale Olive Scale Frosted Scale Mealy Plum Aphid Eggs Leaf Curl Plum Aphid Eggs Brown Mite Eggs European Red Mite Eggs	PRE-BLOOM WHEN FRUIT BUDS SHOW GREEN TIPS: Use 1½ to 2 gals. per 100 gals. of water at 300 to 600 gals. per acre. (Concentrate Spray: Use 6 to 8 gals. per acre in a minimum of 20 gals. water per acre.)	Application at this time is preferred.
Two-spotted Spider Pacific Mite European Red Mite Brown Mite	FOLIAR SPRAY: Use ½ to 1 gals. per 100 gals. water at 300 to 600 gals. per acre. (Concentrate Spray: Use 4 to 6 gals. per acre in a minimum of 20 gals. water per acre.)	Oil sprays may injure certain varieties of plums and prunes, check for tolerance prior to treatment.

For Aerial Application: Dormant and Delayed Dormant Application Only: Apply 3 to 5 gals. in a minimum of 20 gals. of water per acre. Apply by air only when soil conditions do not permit regular ground application.

CITRUS: (California Only) - Lemons, Grapefruit, and Oranges (Navel and Valencia), Limes, Tangerines and Tangelos.

PEST	APPLICATION RATE & TIMING	COMMENTS
Mites -Citrus Bud -Citrus Flat -Citrus Red	Central California: Apply ½ to 1½ gals. of oil per 100 gals. of water as thorough coverage spray (TC). Alternately, use a low volume spray (LV) to apply 1 to 10 gals. of oil per acre.	For Lemons, Grapefruit, Oranges (Navels and Valencias), Limes, Tangelos and Tangerines apply July through September.
Scales -Black -Brown Soft -California Red -Citricola -Purple -Yellow	Apply up to 10 gals. of oil per acre on bearing trees up to 10 feet in height. Add 1 gal. of oil per acre for each foot of tree height over 10 feet. Apply as a low volume spray (LV) 1 to 10 gals. of oil per acre. Apply as a low volume spray (LV) 1 to 10 gals. of oil per acre.	For Lemons, Grapefruit, Oranges (Navels and Valencias), Limes, Tangelos and Tangerines. For Lemons and Limes, make applications in all months except December through February; for Grapefruit make applications July through October; for Oranges make applications on Navels July through September and on Valencias, Tangelos and Tangerines July through October.
Mites -Citrus Bud -Citrus Flat -Citrus Red	Southern California: Apply ½ to 1½ gals. of oil per 100 gals. water as a thorough coverage spray (TC). Alternately, use a low volume spray (LV) to apply 1 to 10 gals. of oil per acre.	For Coastal Lemons and Limes, apply May and June and/or September through December. For Interior Lemons and Limes, apply in April and May and/or September through November.
Scales -Black -Brown Soft -California Red -Citricola -Purple -Yellow	Apply up to 10 gals. of oil per acre on bearing trees up to 10 feet in height. Add 1 gal. of oil per acre for each foot of tree height over 10 feet. Apply as a low volume spray (LV) 1 to 10 gals of oil per acre. Apply as a low volume spray (LV) 1 to 10 gals. of oil per acre.	For Lemons, Grapefruit, Oranges (Navels and Valencias), Limes, Tangelos and Tangerines. For Lemons and Limes, make applications in all months except December through February; for Grapefruit make applications July through October; for Oranges make applications on Navels July through September and on Valencias, Tangelos and Tangerines July through October.
Mites -Citrus Bud -Citrus Flat -Citrus Red	Oxnard Plains Area: Apply ½ to 1½ gals. of oil per 100 gals. of water as a thorough coverage spray (TC). Alternately, use a low volume spray (LV) to apply 1 to 10 gals. of oil per acre. Apply up to 10 gals. of oil per acre on bearing trees up to 10 feet in height. Add 1 gal. of oil per acre for each foot of tree height over 10 feet. Apply as a low volume spray (LV) 1 to 10 gals. of oil per acre.	For Lemons, Grapefruit, Oranges (Navels and Valencias), Limes, Tangelos and Tangerines. Make applications as needed throughout the year.
Scales -Black -Brown Soft -California Red -Citricola -Purple -Yellow		

Thorough coverage spray (TC): Applications achieving a uniform film wetting of all portions of the tree at 250 to 4500 gallons of spray solution per acre, the total volume is dependent of the size of the trees.

Low-volume spray (LV): Applications with droplet depositions achieved on all interior and exterior parts of the tree using 10 to 100 gallons of spray solution per acre.

Do not apply prior to or during a period of excessively high temperatures.

Do not apply oil spray when trees show stress.

Consult with your local University of California Cooperative Extension Specialist for proper timing and spray program.

6/6

BioCover™ MLT
EPA REG. NO. 34704-INL

CITRUS: (Florida Only) -Grapefruit, Lemons, Limes, Oranges, Tangelos and Tangerines.

PEST	APPLICATION RATE & TIMING	COMMENTS
Citrus Rust Mite	5 to 10 gals. of oil per acre. Apply as a dilute spray.	Does not control citrus Snow Scale.
Spider Mites	5 to 10 gals. of oil per acre. Apply as a dilute spray.	
Scale Insects -Black Soft -Brown Soft -Chaff -Glover -Purple -Red -Yellow	5 gals. of oil per acre + 6 pints Ethion 4EC. Apply as a dilute spray.	Summer spray only. Observe all applicable precautions on Ethion label. Do not apply to Navel oranges.
Scale Insects -Black Soft -Brown Soft -Chaff -Glover -Purple -Red -Yellow	5 to 10 gals. of oil per acre. Apply as a dilute spray.	
Whitefly	5 gals. of oil per acre. Apply as a dilute spray.	
Greasy Spot (Grapefruit)	5 gals. of oil per acre. Apply as a dilute spray.	Tank mix with sufficient copper fungicide to provide 4 lbs. per acre of metallic copper.
Greasy Spot (all other varieties)	5 to 10 gals. of oil per acre. Apply as a dilute spray.	If disease pressure is severe, tank mix 4 lbs. (metallic) copper per acre. Do not exceed 5 gals. oil if copper used.
Loosening sooty mold	5 to 10 gals. of oil per acre. Apply as a dilute spray.	Use higher rate when disease pressure is heavy.

Dilute spray: Apply the spray oil application rate in up to 4500 gallons of water per acre, or 30 to 35 gallons per tree, to over 45 gallons per tree for large trees. Best results can be obtained when the oil is added to a partially filled spray tank under proper agitation and all foliage is thoroughly covered.

Precautions in the use of oil include: Do not apply oil spray when trees are wilting. Do not apply oil and sulfur within 3 weeks of each other in Florida. Oil spray applied after October 1 may increase susceptibility of trees to cold damage and may reduce the fruit crop the following year. Oil sprays applied in the fall may inhibit solids formation in the juice and retard coloring of fruit and should not be applied within 60 days of anticipated harvest. Sensitive foliage may be injured.

Consult the Florida Citrus Spray and Dust Schedule, or your Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local.

AVOCADOS - HAAS VARIETY ONLY:

PESTS	APPLICATION RATE & TIMING	COMMENTS
Avocado Brown Mite	Use ½ to 1 gal. per 100 gals. of water with thorough coverage in 400 to 3000 gal. per acre. (Concentrate Spray: Use 4 to 15 gals. per acre in a minimum of 50 to 100 gals. per acre.)	Apply through late fall. Consult with your local University of California Cooperative Extension Service for proper timing and application parameters.

BANANAS:

PESTS	APPLICATION RATE & TIMING	COMMENTS
Black Leaf Streak (<i>Mycosphaerella fijensis</i>)	Use ½ to 1½ gals. in 20 to 100 gals. of water per acre. (Mist Blower: Use 1½ gals. in 6 gals. of water per acre.)	Begin application when disease first appears and continue every 10 to 15 days during rainy periods or as is needed. Do not apply when daytime temperatures exceed 80° F as injury may occur. Phytotoxicity may be affected by the quantity of oil used, vigor of trees and temperature at the time of application. Bag fruits or avoid direct spraying on unbagged fruits.

NOTICE

IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE, THE MANUFACTURER OR SELLER. IN NO CASE SHALL PLATTE, THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE, THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

FORMULATED FOR
PLATTE CHEMICAL CO.
419 18TH STREET, GREELEY, COLORADO 80631-5852