

MAR 21 2002


Ms. Glenda Haage
Registration Manager
PLATTE CHEMICAL COMPANY
419 18th Street
Greeley, CO 80632-0667

SUBJECT: Label Amendment
Weed & Feed for St. Augustine Grass, EPA Reg. No. 34704-822
Your Application Dated February 19, 2002

The above mentioned amendment submitted in connection with registration under Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable.

A stamped copy is enclosed for your records. Please submit one(1) final printed copy for the above mentioned label before releasing the product for shipment. The amended labeling supersedes all previously accepted ones.

Sincerely,


James Tompkins
Product Manager, (25)
Herbicide Branch
Registration Division (7505C)

Enclosure

CONCURRENCES

SYMBOL ▶	7505C							
SURNAME ▶	MINOR, E.							
DATE ▶	Mar 20, 2002							

ACCEPTED
with COMMENT
In EPA Letter Dated

MAR 21 2002

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

Weed & Feed

For St. Augustine Grass

34704-822

- WEED CONTROL PLUS TURF FERTILIZER
- FOR EXCLUSIVE USE ON ST. AUGUSTINE, CENTIPEDE, AND ZOYSIA GRASSES ONLY
- FOR PRE-EMERGENT AND POST-EMERGENT LAWN WEED CONTROL

ACTIVE INGREDIENT:

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine 0.92%

INERT INGREDIENTS: 99.08%
TOTAL 100.00%

This product contains 18.35 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

Refer to Fertilizer Tag for Analysis

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

EPA REG. NO. 34704-822

EPA EST. NO. 61933-FL-10 & 11

NET WEIGHT 50 LBS. (22.68 KG)

IHT

EXP 01/02

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid inhalation of dust. Wash thoroughly after handling and before eating or smoking. Keep away from feed or food products.

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-228-5635, EXT. 136, OR CALL COLLECT, 612-851-8180, EXT. 136.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Use only as directed on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in original labeled bag in a cool, dry, locked area designated for turf herbicides, away from growing plants; out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

HOUSEHOLD: Do not reuse empty bag. Securely wrap original container in several layers of newspaper and discard in trash.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

CONTROLS THESE WEEDS

GROWING IN ST. AUGUSTINE - CENTIPEDE- ZOYSIA GRASSES

Annual Bluegrass (Poa Annual)	Henbit
Annual Sowthistle	Knotweed
Annual Clover	Lespedeza
Bahiagrass	Mat Lippia (Matchweed)
Betony	Moneywort
Black Medic (yellow trefoil)	Morningglory
Blue Toadflax	Narrowleaf Vetch
Bull Paspalum	Oxalis (Common and Yellow Woodsorrel)
Buckhorn Plantain	Parsley-Piert
Bur Clover	Pennywort (dollarweed)
Carolina Geranium (Cranesbill)	Red Sorrel
Chickweed (Common and Mouse Ear)	Rescuegrass
Common Dandelion	Sandspur
Common Purslane	Sedges*
Crabgrass	Shepherdspurse
Cudweed	Spurge
Cutleaf Evening Primrose	Spurweed
Dallisgrass	Stinging Nettle
Dichondra	Swinecrest
Dollar Weed	Vasseygrass
Florida Betony (Rattlesnake Weed)	Venus Looking Glass
Florida Pusley	Virginia Creeper
Fragrant Cudweed (Rabbit Tobacco)	Wild Mustard
	Wiregrass

*Repeat applications are required for control of nutsedge grasses.

This product will control both emerged weeds and weeds from seed. Some

**WEED & FEED
FOR ST. AUGUSTINE GRASS
EPA REG. NO. 34704-822**

moisture is naturally required to activate the herbicide and the fertilizer. For best results rainfall or watering is necessary within 7 to 10 days after treatment. See "AMOUNT TO USE".

WHEN TO APPLY

For best results apply as a pre-emergent application before weeds come up. Established weeds are more difficult to control and may require an application both in the fall and spring. On newly sprigged St. Augustine, Centipede or Zoysiagrass lawns, apply immediately after planting to kill weeds before they compete with new lawn growth. Apply when grass blades are dry. On hot days (above 80 degrees) water lawn to rinse particles off the grass.

FOR SPRING GERMINATING SEEDS

Apply in January, February or March to prevent germination of Crabgrass, Bull Paspalum, Sedges (from seed) Spotted Spurge and other weeds as listed. For control of broadleaf weeds this product may be applied up through June 15th.

FOR WINTER WEEDS

A fall or early winter application will give best control of weed seeds such as Henbit, Chickweed, Rabbit Tobacco, Florida Betony, and Evening Primrose that germinates in the late fall or winter months.

AMOUNT TO USE

Apply at the rate of 50 pounds per 10,000 sq. ft.

Water thoroughly after application. Use herbicides properly. Do not overdose. Read Directions for Use carefully.

IMPORTANT PRECAUTIONS

Do not apply this product or over the root zone of ornamental trees and shrubs, bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to newly seeded St. Augustine, Centipede or Zoysia lawns until after the second mowing. Avoid applications on garden plants, flowers and shrubs. Delay spring plantings for one month after application. Clean spreader after use.

Do not plant sprigs for one month, or sow grass seed for six months following application of this product. May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80° F or above, water sufficiently to rinse particles off the grass blades.

SPREADER SETTINGS - 50 lbs./10,000 Sq. Ft.

Spyker Pro	xx
Scotts R7X	xx
Lesco B'Cast	xx
Cyclone Pro	xx
Prize CB-85	xx
Prize CBT-85	xx

These spreader settings are for use with new equipment. Your spreader may need calibrating since proper application is based on the spreader conditions with regard to age and use, your walking speed and the unevenness of the terrain. Poor weed control will result from uneven or improper application.

NOTICE

IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE, THE MANUFACTURER OR SELLER. IN NO CASE SHALL PLATTE, THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE, THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

**FORMULATED FOR
PLATTE CHEMICAL CO.
419 18TH STREET, GREELEY, COLORADO 80631-5852**