



U.S. ENVIRONMENTAL PROTECTION  
AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

EPA Reg.  
Number:

34704-834

Date of Issuance:

21 MAR 2008

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:

**Ronstar 0.67% on Fertilizer**

Name and Address of Registrant (include ZIP Code):

**Loveland Products, Inc.**  
**P.O. Box 1286**  
**Greeley, CO 80632-1286**

**Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.**

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:

1. Per the acute toxicity review, the signal words currently on the label "WARNING-AVISO" must be revised to read "CAUTION".
2. Per the acute toxicity review, no precautionary statements are required for this product. However, you may use toxicity category III precautionary statements, if you wish.
3. Add the bolded word to the User Safety Recommendation text in the statement "User should remove clothing/PPE immediately if pesticide gets inside."
4. Per the RED, the early entry PPE must be revised to read, "Coveralls, Shoes plus socks, Chemical-resistant gloves made of any waterproof material."
5. Per the RED, the following text must be added to the Non-Agricultural Use Requirement box, "Do not enter or allow others to enter the treated areas until dusts have settled. If watered-in immediately after application: Do not enter or allow entry to the treated area (except for those involved in the watering) until the watering-in is complete and the surface is dry."

Signature of Approving Official:

Joanne I. Miller  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

*Joanne I. Miller*

Date:

21 MAR 2008

6. To the Warranty section add "to the extent consistent with applicable law" in front of "the exclusive remedy" and change "to the extent allowed by applicable law" to "to the extent consistent with applicable law".

Submit one copy of the revised final printed label for the record before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505P)

Enclosure



# Ronstar® 0.67% on Fertilizer

For sale to and use by professional applicators only.  
Not for sale to or use by homeowners/consumers.

A Pre-emergent Weed Control Product plus a Quality Fertilizer for use on varieties of Perennial Bluegrass, Bentgrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustine grass, Seashore Paspalum, Tall Fescue and Zoysia. Also For Use Around Ornamental Shrubs, Vines, Trees & Ground Covers.

ACTIVE INGREDIENT:	
Oxadiazon*	0.67%
OTHER INGREDIENTS:	99.33%
<b>TOTAL</b>	<b>100.00%</b>

Refer to Fertilizer Tag for Analysis

\*CAS [19666-30-9] Contains RONSTAR® Herbicide. A registered trademark of Aventis, Inc.

- Not for use on home lawns.
- For use on golf courses, nurseries, parks, sod farms, athletic fields, and other recreational and ornamental turfgrass.

## KEEP OUT OF REACH OF CHILDREN WARNING—AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA REG. NO. 34704-834

EPA EST. NO. \_\_\_\_\_

NEW CONTENTS 50 POUNDS (22.68 KG)

IHT

EXP 12/05

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**WARNING:** Causes substantial but temporary eye injury. Wear protective eyewear (goggles, face shield or safety glasses). Causes skin irritation. Do not get in eyes, on skin or clothing. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid breathing dust, contact with skin, eyes or clothing. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Loaders, applicators, and other handlers must wear: Long-sleeved shirt and long pants, and Shoes plus socks. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

21 MAR 2008

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

34704-834

### FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:  
1-800-301-7976.

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment.

### USER SAFETY RECOMMENDATIONS

Users should:  
Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.  
Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.  
Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system. Apply only as specified on this label.

A maximum application rate of 895 pounds of product per acre per year (equivalent to 6 pounds ai/A/year), except in areas where there is heavy weed infestation. In areas of heavy weed infestation, the maximum application rate is 1194 pounds of product per acre per year (equivalent to 8 pounds ai/A/year).

A maximum single application rate of 597 pounds of product per acre (equivalent to 4 pounds ai/A)

Not for use on home lawns

# Ronstar® 0.67% on Fertilizer

EPA REG. NO. 34704-834

For best results, water-in the product as soon as practical after application.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on sod farms, and nurseries, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Shoes plus socks, and chemical-resistant gloves made of any waterproof material.

Controls Crabgrass, Goosegrass, Poa Annua, Annual Sedge and other weeds as listed, on Golf Courses, Commercial Lawns, and other Athletic and/or Recreational Ornamental Turf.

## WHERE, HOW MUCH AND WHEN TO USE ON TURFGRASS

This product may be used at full rate or as a "split application" in established perennial Bluegrass, Bentgrass (see restrictions), Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue, and Zoysiagrass turf, at apartment/condominium complexes, corporate and public parks, golf courses, sod farms, athletic fields, playgrounds, cemeteries and other ornamental and recreational turfgrass areas. Not for use on home lawns. The turf should be free of leaves and thatch and in condition to allow granules to reach the soil surface. Know the square footage of the area to be treated and uniformly apply using a granule spreader. Do not apply to turf when it is wet. When using at full rate do not overlap spreader applications, especially on St. Augustinegrass since discoloration "streaking" may temporarily occur from the "doubled rate".

See section on this label entitled PRECAUTIONS IN USING ON TURFGRASS for the maximum rate for St. Augustinegrass and Bentgrass of 3 lbs. a.i. per acre which is equivalent to 10.2 lbs. of this product per 1,000 sq. ft. After applications, if rain is not expected within 24 hours, sprinkle irrigate the treated turf with up to a ¼ inch of water to get maximum weed control and quicker response from the fertilizer nutrients. This product is a pre-emergence herbicide and must be applied prior to weed seed germination (see map of U.S. on this label). It will not control growing weeds.

\*\*See section on this label entitled FOR SPRING APPLICATION ON GOLF COURSES for instructions to apply the 3 lb. or 4 lb. A.I. rate as a SPLIT APPLICATION.

## Pre-emergence Weed Control

Weeds Controlled	Lbs. a.i. / Acre Lbs. product/acre	Lbs. of Product per 1,000 sq. ft.	Directions - Refer to Map of U.S.
Crabgrass ( <i>Digitaria</i> , spp.)	2 to 4 lbs. a.i./A 300 to 600 lbs. product/acre	6.8 to 13.6 lbs.	Apply in late winter or early spring prior to weed seed germination. See map. In areas 4 & 5 applications made after April or May, may not be fully effective.
Goosegrass ( <i>Eleusine indica</i> ) Field Sandbur ( <i>Cenchrus incertus</i> ) Annual Sedge ( <i>Cyperus compressus</i> )			
Annual Bluegrass ( <i>Poa annua</i> spp.) (See Note Below)	2 to 4 lbs. a.i./A 300 to 600 lbs./A	6.8 to 13.6 lbs.	Apply in late summer to early fall prior to weed seed germination.
Virginia Buttonweed ( <i>Diodia virginiana</i> ) Pigweed ( <i>Amaranthus</i> spp.) Florida Pusley ( <i>Richardia scabra</i> ) Oxalis ( <i>Oxalis</i> spp.) Stinging Nettle ( <i>Urtica dioica</i> ) Carpet Grass ( <i>Axonopus officinis</i> )	4 lbs. a.i./A 600 lbs. product/acre	13.6 lbs.	Apply in early spring prior to weed seed germination.

## Fall Application for Spring Weed Control in Cool Season Turf

Weeds Controlled	Lbs. a.i. / Acre Lbs. product/acre	Lbs. of Product per 1,000 sq. ft.	Directions - Refer to Map of U.S.
Crabgrass, Goosegrass and other annual weeds as listed.	3 to 4 lbs. a.i./A 450 to 600 lbs. product/acre	10.2 to 13.6 lbs./1,000 sq. ft.	For use in Bluegrass, Bentgrass, perennial Ryegrass and Tall Fescue. Make application in the fall after turf has become dormant. For late fall applications, seeding into treated areas should be delayed until 4 months after greenup.

## Annual Bluegrass Control in Bermudagrass Fairways Overseeded with Perennial Ryegrass only.

Weeds Controlled	Lbs. a.i. / Acre Lbs. product/acre	Lbs. of Product per 1,000 sq. ft.	Directions - Refer to Map of U.S.
Annual Bluegrass ( <i>Poa annua</i> spp.)	2 lbs. a.i./A 300 lbs/acre	6.0 lbs./1,000 sq. ft.	Apply in late summer to early fall prior to <i>Poa annua</i> spp. seed germination. Any <i>Poa annua</i> spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where <i>Poa trivialis</i> is present in the overseed mixture. Apply the full 2 lb. a.i. rate in one application.

NOTE: In many areas annual bluegrass exists as at least two biotypes or subspecies of *Poa Annua*. The true annual biotype, "*Poa Annua* spp. *Annua*" which germinates from seed every year is controlled while the biotype, "*Poa Annua* spp. *raptans*" which behaves like a perennial and survives through the summer, as an established plant, cannot be controlled. Consult the extension service or university weed specialist in your area for more information concerning your locale.

## Annual Weed Control during establishment from "Sprigs" of Bermudagrass, Zoysiagrass, and Seashore Paspalum.

Weeds Controlled	Lbs. a.i. / Acre Lbs. product/acre	Lbs. of Product per 1,000 sq. ft.	Directions - Refer to Map of U.S.
Crabgrass and Goosegrass, Annual Sedge and other annual weeds.	2 to 3 lbs. a.i./A 300 to 450 lbs. per acre	6.8 to 10.2 lbs. per 1000 sq. ft.	For Bermudagrass make application immediately prior to, or immediately after sprigging. For Seashore Paspalum and Zoysiagrass make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable Seashore Paspalum injury. Disturbance of the soil surface after application may disrupt the herbicide barrier and reduce effectiveness. For best results sprinkler irrigate the "sprigged" area with up to ¼ inch water after application.

## CRABGRASS WEED GERMINATION DATES



## NON-AGRICULTURAL USES REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or nurseries. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and impervious gloves when handling. Launder exposed clothing before reuse.

## FOR SPRING APPLICATION ON GOLF COURSES

The 3 lb. and 4 lb. a.i. per acre rates may be applied as a split application. Weed control and better turf management can be programmed by splitting the a.i. per acre rate of active ingredient into two separate applications. For example to accomplish this in region #5 (See map of U.S. on this label); first apply 1.5 lbs. a.i. (for required 3 lbs. a.i./acre rate) or first apply 2 lbs. a.i. acre (for a 4 lb. a.i./acre rate) as the 1st application 1 to 2 weeks before estimated weed seed germination or in late December or early January; then follow with the second "½ rate" in 60 days or the first of March. The timing for these applications should be closely programmed to the geographic region (see map of U.S. on this label) and adjusted to an "early" or "late" spring. Do not split the 2 lb. a.i./acre rate. The 3 lb. a.i. / acre rate should not be split if applied to new turf areas first under turf management where weed pressure is expected to be heavy.

## Bermuda & Paspalum Directions - Refer to U.S. Map on this label.

For Bermudagrass make application immediately prior to or after after sprigging. For Seashore Paspalum apply 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable paspalum injury. Disturbance of the soil surface after application may disrupt the herbicide barrier and reduce effectiveness. For best results sprinkler irrigate the "sprigged" area with up to ¼ inch water after application.

**Ronstar® 0.67% on Fertilizer**  
EPA REG. NO. 34704-834

**PRECAUTIONS IN USING ON TURFGRASS  
and also PRECAUTIONS WHEN SEEDING**

For St. Augustinegrass and Bentgrass turf, DO NOT exceed the rate of 3 lbs. of active ingredient per acre which is equal to 10.2 lbs. of this product per 1,000 sq. ft. DO NOT apply to Bentgrass turf mowed less than 3/8 inch. This product may cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass but this discoloration is temporary and normally outgrown within 2 to 3 weeks. Avoid contact with flowers and shrubs except as recommended elsewhere for ornamental plants on this label. DO NOT apply to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. DO NOT apply to Dichondra or Centipedegrass. DO NOT use on golf course putting greens or tees because of the varying cultural practices used.

**WEEDS CONTROLLED**

Annual Bluegrass	Goosegrass*
Annual Sedge	Jungle Ricegrass**
Barnyardgrass*	Lambsquarter
Bittercress	Liverwort**
Carpetweed	Oxalis
Carpetgrass	Panama Paspalum**
Common Groundsel	Pennsylvania Smartweed
Common Purslane	Redroot Pigweed
Crabgrass	Sensitive plant
Evening Primrose*	Shepardspurse
Fireweed	Sow thistle
Field Sandbur	Speedwell-Spotted Catsear
Foxtail; Green, Bristly	Spurge; Garden, Petty, Prostrate**
Fall Panicum	Stinging Nettle
Florida Pusley	Swinecress*
Galinsoga*	Virginia Buttonweed
Golden Ragwort	Yellow Woodsorrel

\* For the weeds with 1 asterisk above, use 3 to 4 lbs. a.i. per acre (equal to 450 lbs. to 600 lbs. of this product per acre or 10.2 lbs. to 13.6 lbs. per 1,000 sq. ft.).

\*\*For the difficult-to-control weeds with 2 asterisks above, use the 4 lbs.a.i. per acre (equal to 600 lbs. of this product/acre or 13.6 lbs. per 1,000 sq. ft.)

**NOTE:** For newly seeded lawns do not apply this product until after the second cutting. For best turf management, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied.

**SPREADER SETTINGS**

Spreader settings are calculated to apply the recommended rates of active ingredient per acre based on this product bulk density of 50 to 60 lbs. per cubic foot. Settings will vary if your turfgrass requires this EPA registered herbicide/fertilizer combination product with a lighter bulk density. To check your spreader settings refer to the spreader manual or manufacturer. Test apply enough product to cover 500 sq. ft. based on the amount of product a.i. you wish to apply per acre. Select a suitable rate from the Pre-emergence Weed Control chart on this label. Adjust and record the setting if required. Apply as directed on this label.

Spreader Settings	for 2 lbs. a.i./Acre	for 3 lbs. a.i./Acre	for 4 lbs. i./Acre
Cyclone B3-Pro	#2.75	#3.25	#3.9
Scotts Pro R-8A	#H	#1+	#J+

Plan on an approximately 3 mph spreader speed to obtain best coverage. Know the exact square footage before application. Incorrect application will result in poor weed control.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

For a Wide Variety of listed Woody \* Ornamental Shrubs, Vines, Trees, and Ground Covers in Landscape Plantings and for Container and Field Grown Nursery Plants.

For use only on plants that will not bear edible fruit within one year of application.

5 8 6

**ESTABLISHED ORNAMENTALS and GROUND COVERS**  
**COMMON NAMES**

Abutilon	Jasmine, Orange
Acacia	Jasmine, Primrose
Agapanthus	Jasmine, Star
Allamanda, Common	Jasmine, Showy
Andromeda	Juniper
Aralia	Justica
Arborvitae	Kwanzan Cherry
Ash, Green	Lamb's Ear
Azalea	Lantana, Trailing
Baby's Breath	Lapalapa
Bald Cypress	Larch, European
Bamboo, Feathery	Laurel, English
Banyan, Weeping	Leopards - Bane
Barberry	Leucothoe, Coast
Beefsteak Plant	Leucothoe, Drooping
Bearberry	Lily of the Nile
Beech, European	Lilac
Birch, Paper	Littleleaf Linden
Blanket Flower	Loquat*
Blazing Star	Magnolia
Boxwood, Japanese	Mahonia, Grape Holly
Boxwood, Common	Maple
Bougainvillea	Metallic Plant
Brassia, Dwarf	Mexican Orange
Buddleia	Mock Orange
Bugleflower	Money Tree
Callistemon	Murraea
Camellia, Common, Jap	Myoporum
Camellia, Sasanque	Myrtle
Candytuft Carissa	Nandina
Carnation	Oak
Ceanothus	Ochana
Cedar, Red	Ohai
Cedar, Atlas	Oleander
Cistus	Orthosiphon
Citrus, Ornamental*	Osmanthus
Grapefruit	Pachysandra
Lemon	Palm, Bamboo
Lime	Palm, Fishtail
Mandarin	Palm, Licuala
Tangelo	Palm, Macarthur
Tangerine	Palm, Mabnee
Temple	Palm, Manila
Cuban Shaddock	Palm, Parlor
Valencia	Palm, Rattan
Washington Navel	Palm, Scaling Wax
Cleyera	Palm, Sugar
Coreopsis	Paperbark Tree
Coromandel	Peony
Cotoneaster	Periwinkle, Common
Crabapple	Periwinkle, Large
Crapemyrtle	Photinia, Frasers
Chrysanthemum	Poinciana
Cypress	Pine
Cypress, Leyland	Pittosporum, Japanese
Dahlia	Plum, Natal
Daphne, Fragrant	Podocarpus, Yew
Deutzia, Slender	Pomegranite
Dieffenbachia	Poplar
Dogwood, Common	Potentilla
Dogwood, Redosier	Privet
Dogwood, Kousa	Protea species
Douglas Fir	Pyracantha
Elm	Raphiolepis
Escallonia	Redwood, Dawn
Eucalyptus	Rhododendron
Euryops, Gray Leaved	Rose spp.
Euonymus, Spreading	Russian Olive
Euonymus, Wintercreeper	Sargent Cherry
Falsecypress, Hinoki	Scotch Broom
Falsecypress, Japanese	Sedum
Flowering Quince	Serviceberry, Saskatoon*
Forsythia	Spruce
Gardenia	Statice
Gazania Trailing	St. Johnswort, Golden
Ginko	St. Johnswort, Goldencup
Gladiolas	Strawberry Tree
Golden Raintree	Sumac, Staghorn
Heath, Spring	Sweet William
Heather, Scotch	Syzygium
Heliotropium	Trailing African Daisy
Hemlock, Canada	Viburnum
Hibiscus	Wedelia
Honeysuckle	Wisteria, chinese
Holly	Witchhazel, common
Ice Plant	Woadwaxen, common
Ilima	Woolly Thyme
Ivy, Alergian	Xylosma
Ivy, English	Yew

6 8 6

**Ronstar® 0.67% Fertilizer**  
**EPA REG. NO. 34704-834**

**WHERE, HOW MUCH, AND WHEN  
 TO USE AROUND ORNAMENTALS**

Can be applied anytime of the year prior to weed seed germination to both newly transplanted and established actively growing or dormant ornamental plants where both weed control and fertilization is desired. Rainfall or overhead irrigation ¼ inch water will improve weed control actively. Do not mix into soil. Disturbing or cultivating the soil after treatment may reduce weed control. Do not apply granules to wet foliage or under conditions in which granules will collect on leaves, if so, sprinkler-irrigate the plants to move granules off foliage and down to soil level. Do not apply in greenhouses as plant injury may likely result. Use the higher application rates for areas that have a history of heavy weed pressure. For best weed control around ornamentals first remove any existing weed growth and if possible rake, cultivate and smooth the soil surface by breaking up clumps. Broadcast the granules evenly over the prepared soil by applying 2 lbs. to 4 lbs. active ingredient per acre, equal to 300 to 600 lbs. of this product/acre or 6.8 to 13.6 lbs. per 1,000 sq. ft.

**NEWLY TRANSPLANTED  
 ORNAMENTALS AND GROUND COVERS**  
**COMMON NAMES**

Blanket Flower	Leopards-Bane
Brassaia, Dwarf	Metallic plant
Camellia	Oak
Carnation	Periwinkle, Common
Cedar, Red	Periwinkle, Large
Coromandel	Pine
Dogwood, Flowering	Poplar
Elm	Privet
Forsythia	Protea Species
Gazania, Trailing	Rhododendron
Hackberry	Russian Olive
Ivy, Algerian	Spruce
Ivy, English	Trailing African Daisy
Hemlock, Canada	Wedelia
Holly	Wisteria, Chinese
Ice Plant	Yew
Juniper	

**FOREST NURSERY APPLICATION**

<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>
Pine, Loblolly	<i>Pinus, taeda</i>
Pine, Eastern White	<i>Pinus, Strobus</i>
Pine, Slash	<i>Pinus, Canbaea</i>

**SENSITIVE PLANTS**

Under certain growing conditions, the plant varieties listed below may be sensitive to this product. Growers should evaluate usage on a small scale prior to large scale applications.

<b>PLANT SPECIES</b>	<b>COMMON NAMES</b>
Andromeda ( <i>Pieris japonica</i> )	<b>VARIETIES</b> Snowdrift, Temple Bell
Azalea	Pink Champagne, Hershey, ( <i>Rhododendron</i> spp.)
Cotoneaster	Leuchtentuer, Memoire John Hearrens, Hino Crimson, Everest, Girard Eichholz, Himalayan, Shangri-la ( <i>Cotoneaster</i> spp.)
Crape myrtle ( <i>Lagerstroemias indica</i> )	Petite Red Imperial, and Seminole Bush
Heath, Spring ( <i>Erica carnea</i> )	Mediterranean Pink
Hemlock, Canada ( <i>Tsuga canaadensis</i> )	Pendula
Hibiscus ( <i>Hibiscus Syriacus</i> )	Red Heart
Rhododendron	Daphnoides, Nova Zembia, ( <i>Rhododendron</i> spp.)
Spruce, Norway ( <i>Picea abies</i> )	Catawba Album Pumila
Spruce, White ( <i>Picea glauca</i> )	Densata
Viburnum	Davidii ( <i>Viburnum</i> spp.)

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**  
 BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS PRODUCT IS SOLD AS IS TO THE EXTENT ALLOWED BY APPLICABLE LAW. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND, TO THE EXTENT REQUIRED BY APPLICABLE LAW, WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, 7251 WEST 4TH STREET, GREELEY, CO 80634.

TO THE EXTENT ALLOWED BY APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT, TO THE EXTENT ALLOWED BY APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

FORMULATED FOR



P.O. BOX 1286, GREELEY, COLORADO 80632-1286