

34704-849

1/12/2009

10A13



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. Scott Baker
Registration Manager
Loveland Products, Inc,
P.O. Box 1286
Greeley, Colorado 80632-1286

JAN 12 2009


RE: Notification of Alternate Brand Name: "Leaf Life Organics Brand Glacial Spray Fluid"
EPA Registration Number: 34704-849
Date of Submission: November 14, 2008

Dear Ms. Baker:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated November 14, 2008, for the product Glacial Spray Fluid. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,


Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

20913

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 34704-849	2. EPA Product Manager PM# 67	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Glacial Spray Fluid	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	
5. Name and Address of Applicant (Include ZIP Code) Loveland Products, Inc. P.O. Box 1286 Greeley, Colorado 80632-1286 <input type="checkbox"/> Check if this is a new address	NOTIFICATION JAN 12 2009	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification to add an alternate brand name under PR Notice 98-10. This application is to add the alternate brand name Leaf Life Organics Brand Glacial Spray Fluid.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Scott Baker	Title Registration Manager	Telephone No. (Include Area Code) 970-347-1468
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Registration Manager	
4. Typed Name Scott Baker	5. Date November 14, 2008	



November 14, 2008

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900,
2777 S. Crystal Drive
Arlington, VA 22202

RE: Glacial Spray Fluid (EPA Reg. No. 34704-849)


Loveland Products, Inc., is submitting this notification under PR Notice 98-10 to add an alternate brand name. The alternate name is Leaf Life Organics Brand Glacial Spray Fluid. Please find the following enclosed.

- 1. Application for Registration.
2. 2 copies of the label with the alternate name.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA

If you have any additional questions or concerns, please contact me by telephone 970-347-1468 or scott.baker@uap.com by email.

Sincerely,
[Signature]
Scott Baker
Registration Manager
Enclosures



40913

NOTIFICATION

JAN 12 2009

LEAF LIFE ORGANICS



FOR ORGANIC PRODUCTION

A White Mineral Horticultural Spray Fluid For Insect, Mite and Disease Control.

ACTIVE INGREDIENT:	By Weight
Petroleum Oil	98.4%
INERT INGREDIENTS:	1.6%
	TOTAL 100.0%
Unsulphonated Residue	99.00% (minimum)
Aromatic Composition by ASTM D2140	0.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Inside For Additional Precautionary Statements.


**EPA REG. NO. 34704-849
EPA EST. NO. 51263-FL-001
NET CONTENTS 2½ GALS. (9.46 L)**

IHT 081706 V3 10P07

FORMULATED FOR



P.O. BOX 1286
GREELEY, COLORADO 80632-1286



FIRST AID

Have the product container label with you when calling a poison control center or doctor or going for treatment.

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not give any liquid to a person. • Do not give anything to an unconscious person. • Do not induce vomiting unless told to by a poison control center or doctor.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Petroleum distillate poses aspiration pneumonia hazard. Only minor irritation should be expected from any type of exposure other than pulmonary aspiration. Ingestion may result in some gastroenteritis and mild diarrhea. Viscosity is such that chemical (hydrocarbon) pneumonia is not a hazard; however, aspiration or lipid pneumonia of a less severe variety could result from pulmonary aspiration. Since systemic toxicity is not expected, the added risk of aspiration from attempts to induce vomiting or lavage are deemed not warranted.

**FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:
1-800-301-7976.**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber or viton and shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no

such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Apply this product only as specified on the label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves, such as barrier laminate, nitrile rubber, neoprene rubber or viton and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets out of the treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call collect: CHEMTREC -- 1-(800)-424-9300.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers. **Plastic:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

MIXING DIRECTIONS

Use only in equipment with sufficient agitation to keep spray thoroughly mixed. Be sure tank is clean. With agitator running, start filling tank with water; add GLACIAL SPRAY FLUID when tank is about 1/2 full. Pump this mixture through the overflow and back into tank for about two minutes. If this mixture turns white, it indicates good emulsification. Fill the tank with water, adding other desired materials as tank is filling. If wettable powders are used, add and thoroughly mix with water before adding GLACIAL SPRAY FLUID. Addition of an emulsifiable formulation should be done after the GLACIAL SPRAY FLUID and water has been thoroughly mixed. Observe all precautions and limitations on labeling of all products used in mixtures. Keep agitator running at all times. If an injector is used, first add 25 to 50 gals. of water to tank with engine running to provide good pressure. Put injector suction tube into the measured amount of GLACIAL SPRAY FLUID, opening injector valve to allow the GLACIAL SPRAY FLUID to be sucked into the tank, then fill the tank with water.

Do not use if this material does not emulsify.

APPLICATION INFORMATION

Apply when pests are present and as recommended below. Do not use in undiluted form. Sensitive foliage may be injured.

Evergreens: Glacial Spray Fluid may be used on evergreens, taking into consideration the use precautions noted elsewhere on this label. In order for the spray solution to be effective the target pest and plant should be completely covered. Thorough coverage is essential. Best coverage is achieved with ground equipment using proper spray pressure, gallonage per acre, nozzles, nozzle spacing and tractor speed.

For Vegetables/Plant Virus: Spray at no less than 400 psi using ceramic hollow cone nozzles (ALBUZ® ATR (lilac); HCA (green) color or their equivalent). Adjust the angles of the spray nozzles to achieve complete coverage of the foliage including the lower leaf surfaces. Do not exceed the recommended maximum rates or apply more than recommended.

Preharvest interval - GLACIAL SPRAY FLUID may be used up to the day of harvest.

To ensure the best coverage, dilute applications (greater than 150 gallons of spray solution per acre) are recommended. Concentrate applications (usually from 20 to 150 gallons of spray solution) may reduce the coverage and thus the effectiveness. Application volumes depend on the crop type, crop size and target pest. Adjust the spray volume to obtain the best coverage without runoff. Extreme care should be taken when using concentrate sprays as the potential for enhanced crop phytotoxicity is increased. A concentrate application can provide satisfactory results as long as the spray unit is properly engineered, calibrated and operated.

AERIAL APPLICATIONS - Dormant and Delayed Dormant Application: Only: Apply 3 to 5 gals. in a minimum of 20 gals. of water per acre. Use only as an emergency application when soil conditions or site does not permit regular ground application.

Note: Use efficient equipment of the proper type. Avoid spraying during or immediately prior to hot or freezing weather (over 95°F or under 32°F), hot dry winds, rain or other unsuitable conditions; avoid over spraying or double spraying. Plants should be sprayed only when in vigorous condition and when their moisture condition is suitable. Before using, make certain spray tank is free of sulfur residues. Do not apply sulfur within 14 days except in the Northwest (Oregon and Washington) where it may be used in combination on Pears as a Post Harvest (after old fruit is off the tree), Dormant and Delayed Dormant (before the scales slip, or at or prior to bud swell) treatment and on Apples prior to or up to the Delayed Dormant (1/2 inch green) period. In areas west of the Continental Divide do not apply to Captan treated trees within 90 days after or 60 days before this oil. In areas east of the Continental Divide do not apply to Captan treated trees within 10 to 14 days before and after this oil. If in doubt, spray a portion of one tree previously treated with Sulfur or Captan using 1 quart to 25 gallons water. Burn will show in several days if sufficient time has not elapsed.

This product has not been tested on all species or varieties. Before treating a large area, treat a small area and observe prior to full scale application.

Do not use chlorothalonil or dimethoate in a spray program with GLACIAL SPRAY FLUID on grapes, ornamentals and strawberries. Do not use dicofol in a spray program with GLACIAL SPRAY FLUID on ornamentals and strawberries. Do not use propargite with an oil spray or within 30 days before or after an oil spray.

A period of 14 days must elapse before any of the following chemicals are used before or after an application of GLACIAL SPRAY FLUID: chlorothalonil, dicloran, dicofol, permethrin.

VEGETABLE CROPS

Crop	Pest	Application Rate (gallons of GLACIAL SPRAY FLUID) per 100 gallons of water	Comments
Asparagus Beans Beets Cabbage Cauliflower and other Cole crops Celery Corn Cucurbits Eggplant Lettuce Melon Pepper Potato Pumpkin Radish Squash Sweet Potato Tomato Herbs and Spices Curly Leaf Basil Lemon Balm Mexican Oregano Mint	Aphids Mites Beetle Larvae Leafminers Thrips Leafhopper Whitefly	0.75-1	For virus control on cucurbits use 0.75 gallons per 100 gals. Spray once or twice weekly depending on the rapid growth of the crop and / or vector pressure. When using to control insect transmitted diseases, use higher pressure 400 psi and ceramic hollow cone nozzles.
Oriental Vegetables Acerola, Atemoy, Balsam pear (Bitter melon), Carambola, Chinese broccoli (Gai Lon), Chinese cabbage (Gai Choy), Chinese spinach, Chinese waxgourd, Cilantro, Citron melon, Dasheen, Ginger, Ginseng, Japanese artichoke, Japanese radish (Daikon), Rambutan	Aphids Mites Beetle Larvae Leafminers Thrips Leafhopper Whitefly	0.75-1	For virus control on cucurbits use 0.75 gallons per 100 gals. Spray once or twice weekly depending on the rapid growth of the crop and / or vector pressure. When using to control insect transmitted diseases, use higher pressure 400 psi and ceramic hollow cone nozzles.

FRUIT AND NUT CROPS			
Crop	Pest	Application Rate (gallons of GLACIAL SPRAY FLUID) per 100 gallons of water	Comments
Almond Apricot Cherry	Mites (eggs) San Jose Scale	2	Dormant or Delayed Dormant
	Fruit Tree Leafroller (eggs)	3	
Pistachio	Mites Scales	1-1.5	Summer (foliar or cover) or postharvest. Do not spray oil sensitive varieties. Do not apply to trees lacking moisture.
	Aphid (eggs) Apple Red Bug Mite (eggs) Scales	2	
Apple	Fruit Tree Leafroller (eggs) Mites	3 1-2	Dormant to delayed dormant to 1/2 inch green. Summer (foliar or cover). Do not spray oil sensitive varieties.
	Avocado Brown Mite Thrip Praceia Mite	0.75-1	
Avocado Hass variety only	Black Leaf Streak (<i>Mycosphaerella fijiensis</i>) Black Sigatoka (<i>Mycosphaerella fijiensis</i> var. <i>difformis</i>) Yellow Sigatoka (<i>Mycosphaerella musicola</i>)	0.5-1.5 (Mist Blower: 1 1/2 gals. in a minimum of 6 gals. water per acre.) 1-1.5	Apply through late fall. Consult with your local University of California Cooperative Extension Service for proper timing and application parameters. Begin application when disease first appears and continue every 10 to 15 days during rainy periods or as needed. Do not apply when daytime temperatures exceed 80°F as injury may occur. Phytotoxicity may be affected by the quantity of oil used, vigor of trees and temperature at the time of application. Bag fruits or avoid direct spraying on unbagged fruits. This application is also effective in loosening Sooty Mold Fungus and in preventing its formation by the control of Aphids, Mealybugs, Scales and Whitefly.
Blueberries* Bushberries* Caneberries	Mites Powdery Mildew Rust	0.75-1.5	Using ground equipment, spray for optimum coverage of leaf surfaces. For fungal diseases use at least 200 psi spray pressure.
Figs	Fig Scale Mites	1-3 1-2	Dormant or Delayed Dormant Foliar Spray
	Mealybug Scale		
Grapes	Mealybugs Mites Leafhopper* Whiteflies*	1-2	Using ground equipment, spray for optimum coverage of leaf surfaces. Repeat sprays every 10-14 days. For powdery mildew - Make first application pre-bloom and continue every ten days to three weeks depending on level of disease pressure. Use higher rate and/or shorter spray interval when disease conditions are severe. Oil will remove the bloom on grapes. Table grapes should not be sprayed within 60 days of harvest. On grapes, do not tank mix oil and copper more than once/season. Do not use copper and oil together with fruit present.
	Powdery Mildew		

*Asterisk denotes crop, pest Not approved in California.

FRUIT AND NUT CROPS cont'd.:

Crop	Pest	Application Rate (gallons of GLACIAL SPRAY FLUID) per 100 gallons of water	Comments
Kiwi	Latania Scale Greedy Scale Oleander Scale San Jose Scale	Use 4 to 6 gallons of oil in a minimum of 100 gallons of water per acre. Use higher rate when scale populations are high.	Do not apply after budbreak. Do not use on weak or stressed vines, or where soil moisture is inadequate.
Mango	Mites Powdery Mildew	Use 1 to 1.5 gallons of oil in 100 gallons of water per acre.	Apply every 2 to 3 weeks as necessary depending on level of pest pressure.
Olive	Mites Scales	1.5	Prebloom to Postbloom Buckshot.
Papaya	Mites Powdery Mildew Papaya Ringspot Virus	Use 0.75 to 1.5 gallons in 100 gallons of water per acre.	For Powdery Mildew/Mites: Spray every 10 to 14 days depending on the level of pest pressure. For Virus: Initiate spray when seeds are germinated. Spray weekly using 400 psi spray pressure and ceramic hollow cone nozzles.
Peach Nectarine	Aphid (eggs) Mite (eggs) Fruit Tree Leafroller (eggs) San Jose Scale Mites Scales	2-3 1-2	Dormant or Delayed Dormant Use higher rates for leafroller and scales. Summer (foliar or cover) or postharvest.
Pear	Mite (eggs) Pear Psylla Scales Fruit Tree Leafroller (eggs) Mites Pear Psylla	2 3 1-2 1.5-2	Dormant or Delayed Dormant (up to and including petal fall). Summer (foliar or cover) or postharvest. Do not spray oil sensitive varieties.
Pecan	Aphid (eggs) Mite (eggs) Scales	3	Dormant Do not apply to sensitive varieties or after husk split.
Pineapple	Mealybugs Bud Moth Chinese Rose Beetle	Spray - Use 1 to 2 gallons of oil in 100 gallons of water per acre Dip - Use 1 to 2 gallons of oil in 100 gallons of water, dip for 30 seconds.	Allow fruit to dry for 1 hour before being stored.
Plum Prune	Aphid (eggs) Mite (eggs) Scales Mites Scales	1.5-2 1-1.5	½ inch Summer (foliar to cover) or postharvest. Do not apply to certain fresh market fruits after bloom starts to form as the oil spray will remove the waxy bloom. Do not spray oil sensitive varieties.
Strawberries	Leafminers* Mites Powdery Mildew	0.75	Spray at no less than 400 psi using ceramic spray nozzles (ALBUZ® ATR, lilac color or their equivalent).
Walnut	Mite (eggs) Scales	1-2	Delayed Dormant Do not apply to sensitive varieties or after husk split.

*Asterisk denotes crop, pest Not approved in California.

CITRUS: (California Only) - Grapefruit, Lemons, Limes, Oranges (Navel and Valencia), Tangelos, and Tangerines and other citrus hybrids

PESTS	APPLICATION RATE & TIMING	COMMENTS
Mites -Citrus Red -Citrus Rust	Apply ½ to 1½ gals. of oil per 100 gals. of water as thorough coverage spray (TC). Alternately, use a low volume spray (LV) to apply 1 to 10 gals. of oil per acre.	For Lemons, Limes, Grapefruit and Oranges (Navels and Valencias).
Scale Insects -Black -Brown Soft -Citicola -Purple -Red -Yellow	Apply up to 10 gals. of oil per acre on bearing trees up to 10 feet in height. Add 1 gal. of oil per acre for each foot of height over 10 feet. Apply as a low volume spray (LV) 1 to 10 gals. of oil per acre.	

Thorough coverage spray (TC): Applications achieving a uniform film wetting of all portions of the tree at 250 to 4500 gallons of spray solution per acre, the total volume is dependent of the size of the trees.

Low-volume spray (LV): Applications with droplet depositions achieved on all interior and exterior parts of the tree using 10 to 100 gallons of spray solution per acre.

Do not make an oil spray application within 2 weeks prior to or after an application of sulfur. On Citrus do not make an oil application within 60 days prior to or after an application of sulfur. Do not apply prior to or during period of excessively high summer temperatures. Do not apply oil spray when trees show stress. Consult with your local University of California Cooperative Extension Specialist for proper timing and spray program.

CITRUS: (Florida and Texas) -Grapefruit, Lemons, Limes, Oranges, Tangelos and Tangerines and other citrus hybrids

PESTS	APPLICATION RATE-GALS. GLACIAL SPRAY FLUID	COMMENTS
Citrus Rust Mite	5 to 10 gals. per acre. Apply as a dilute spray.	Does not control Citrus Snow Scale.
Spider Mites	5 to 10 gals. per acre. Apply as a dilute spray.	Summer spray only. Observe all applicable precautions on Ethion label. Do not apply to Navel oranges.
Scale Insects -Black Soft -Brown Soft -Chaff -Glover -Purple -Red -Yellow	5 gals. per acre + 6 pints Ethion 4EC. Apply as a dilute spray.	
Scale Insects -Black Soft -Brown Soft -Chaff -Glover -Purple -Red -Yellow	5 to 10 gals. per acre. Apply as a dilute spray.	
Whitefly	5 gals. per acre. Apply as a dilute spray.	
Greasy Spot (Grapefruit)	5 gals. per acre. Apply as a dilute spray.	Tank mix with sufficient copper fungicide to provide 4 lbs. per acre of metallic copper.
Greasy Spot (all other varieties)	5 to 10 gals. per acre. Apply as a dilute spray.	If disease pressure is severe, tank mix 4 lbs. (metallic) copper per acre. Do not exceed 5 gals. oil if copper used.
Loosening sooty mold	5 to 10 gals. per acre. Apply as a dilute spray.	Use higher rate when disease pressure is heavy.

Dilute spray: Apply the GLACIAL SPRAY FLUID application rate in up to 4500 gallons of water per acre, or 30 to 35 gallons per tree, to over 45 gallons per tree for large trees.

Best results can be obtained when the GLACIAL SPRAY FLUID is added to a partially filled spray tank under proper agitation and all foliage is thoroughly covered.

Precautions in the use of GLACIAL SPRAY FLUID include: Do not apply GLACIAL SPRAY FLUID when trees are wilting. Do not apply GLACIAL SPRAY FLUID and sulfur within 3 weeks of each other in Florida or Texas. GLACIAL SPRAY FLUID applied after October 1 may increase susceptibility of trees to cold damage and may reduce the fruit crop the following year. GLACIAL SPRAY FLUID applied in the fall may inhibit solids formation in the juice and retard coloring of fruit and should not be applied within 60 days of anticipated harvest. Sensitive foliage may be injured.

Florida growers consult the Florida Citrus Spray and Dust Schedule, or your Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your locale. Texas growers consult with your Agricultural Experiment Station or State Extension Specialist for complete details on the spraying program suited to your locale.

COFFEE

PESTS	APPLICATION RATE-GALS. GLACIAL SPRAY FLUID	COMMENTS
Green Scale	Use 3 to 6 gallons in 100 gallons of water per acre. Use only water as a diluent.	Apply when insects appear and treat as necessary at 30 to 60 day intervals. Do not exceed 18 gallons of oil per acre in a growing season. Do not apply within 30 days of harvest when berries are handpicked. Oil residues make picking difficult.

ORNAMENTALS

Crop	Pest	Application Rate (ounces per gallon of water)	Comments
Foliage Ornamentals & Bedding Plants	Aphids Adelgids	1-2.5 oz. per gallon	Summer (Foliar or Cover): Do not spray plants during flowering.
Shade Trees** and Shrubs Including Conifers, Deciduous Broadleaf Evergreens & Woody Ornamentals	Aphids Adelgids Eriophyid Mites Gall Mites Honey Locust Plant Bug Lace Bug Leaf Beetle (larvae) Leafminer Leafrollers Mealybugs Psyllids (immature) Red Mites Sawfly (larvae) Scales (immature) Spider Mites Webworms Whiteflies	2.5-4 oz. per gallon	Winter Dormant Period: Summer (Foliar or Cover): CAUTION: Spray no more than 4 times during the growing season; Use a 2 week minimum application interval. Do not spray when there is obvious moisture deficit in leaves, or the plant is under stress. For fall dormant applications, reduce rate to 2 gallons oil per 100 gallons water and limit use only to: American Red Oak, Japanese Black Pine, Dogwood, Weeping Cherry, Cornelian Cherry, Crabapple, Norway Maple, Purple Plum. NOTE: May cause discoloration of Blue Spruce. Do not use on conifers or ferns not tolerant to oil sprays. Use enough spray solution to completely penetrate the leaf canopy and cover both top and bottom of all of the leaves and stems until wet but without significant runoff. Do not spray during flowering.
Ornamental Trees** Shrubs along city Streets, other rights-of-way including Conifers, Deciduous and Broadleaf Evergreens		1-2.5 oz. per gallon	
Flowering and Foliage Plants including Roses and other Flowering Shrubs	Same as above	Winter Dormant Period: 2.5-4 oz. per gallon	Do not spray during flowering.
Christmas Trees **	Scale (soft & hard) (immature) Spider Mites	2-3 0.75-2	Winter Summer

Use a lower concentration for more sensitive plants.

** GLACIAL SPRAY FLUID removes the glaucous (blue) bloom from such evergreens as Colorado Blue Spruce and Koster Spruce. Always use lower dosage or test spray oil sensitive plants such as Cryptomeria, Smoke Tree, Chamaecypris, Juniper, Japanese Holly and Spruce. Tendency toward sensitivity: Red Cedar and Douglas Fir.

GREENHOUSE AND OUTDOOR

Frequency of Application: For the greenhouse pests listed, use once a week initially, then as the pest is controlled, decrease the frequency to every 2-3 weeks as needed.
Application safety during bloom period should be determined for each individual species of plant to be treated by conducting a small test.

Crop	Pest	Application Rate (gallons of GLACIAL SPRAY FLUID) per 100 gallons of water	Comments
Ageratum Crown of Thorns Dieffenbachia Ferns Ficus Lisianthus Orchid Palms Petunia Poinsettia Schefflera Sunflower	Aphids Fungus Gnats Leafminers Mealybugs Scales (soft & hard) Spider Mites Thrips Whitefly	0.5-1	Due to varietal differences in response to a treatment of GLACIAL SPRAY FLUID at the recommended rate, we recommend conducting a small test on 1 or 2 plants of the specific variety to be treated.
Leaf Polish for hardy plants		1	
Aglaonema Aster Azaleas Begonias Browallia Camellias Carnation Celosia Chrysanthemum Coleus Cosmos Dusty Miller Easter Lilies Gardenias Gerbera Helichrysum Hibiscus Foliage Hydrangea Jade Plant Marigold New Guinea Impatiens Nicotonia Philedendron Portulaca Reiger Begonias Snapdragon Zinnias	Aphids Fungus Gnats Leafminers Mealybugs Scales (soft & hard) Spider Mites Thrips Whitefly	1-2	Due to varietal differences in response to a treatment of GLACIAL SPRAY FLUID at the recommended rate, we recommend conducting a small test on 1 or 2 plants of the specific variety to be treated.

Some plants are more sensitive to oil treatments if any of the following conditions are present when application is made: High humidity (74%), an overcast or cloudy day, inadequate air flow, a small greenhouse where the sunlight is magnified causing a fast build-up of temperature and light intensity.

120613

FIELD CROPS

Crop	Pest	Application Rate (gallons of GLACIAL SPRAY FLUID) per 100 gallons of water	Comments
Corn (sweet & field) Sugar Beets	Aphids Mites Leafminers Corn Earworm Rootworm Armyworm Whitefly	Corn: 1-2 Sugar Beets: 2	
Hops*	Mites Powdery Mildew	1-2	For powdery mildew: Initiate sprays at early leaf stage. Continue sprays every 10-14 days. Mite control will be effective under the same spray interval as powdery mildew sprays. Discontinue sprays at burr development.
Tobacco	Aphids Mites Beetle Larvae Leafminers Thrips Leafhopper Whitefly	0.75-1	

*Asterisk denotes crop, pest Not approved in California.

This product has not been tested on all species or varieties. Before treating a large area, treat a small area and observe prior to full scale application.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS PRODUCT IS SOLD AS IS TO THE EXTENT ALLOWED BY APPLICABLE LAW. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND, TO THE EXTENT REQUIRED BY APPLICABLE LAW, WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, 7251 WEST 4TH STREET, GREELEY, CO 80634. TO THE EXTENT ALLOWED BY APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT ALLOWED BY APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Albuz® is a registered trademark of Saint Gobain Ceramiques Avancees Desmarquest Corporation.
Glacial® is a registered trademark of Loveland Products, Inc.

NOTIFICATION

JAN 1 9 2009