



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

September 24, 2025

SENT BY EMAIL

Leah Strayer
leah@pyxisrc.com
LOVELAND PRODUCTS, INC.

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - Updating first aid statement and other typographical updates.
Product Name: BOLL BUSTER
Admin Number: 34704-856
EPA Receipt Date: 08/13/2025
Action Case Number: 00664389

Dear Leah Strayer:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Thomas Harty via email at harty.thomas@epa.gov.

Sincerely,

Thomas Harty, RiskManager
FB, RD
Office of Pesticide Programs

BOLL BUSTER

For Commercial Use Or Agricultural Use Only.
Not For Residential Use

NOTIFICATION

34704-856

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

09/24/2025

ACTIVE INGREDIENT:

Ethephon (2-Chloroethyl) phosphonic acid*55.4%

OTHER INGREDIENTS:44.6%

TOTAL100.0%

*Boll Buster® contains 6.0 pounds ethephon per gallon.

KEEP OUT OF REACH OF CHILDREN

DANGER—PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

{Note to Reviewer: If space limitations on the front panel do not permit addition of the First Aid Statement in English and/or Spanish (per PRIA V, PR Notice 2001-1 40 CFR 156.68 (d), including guidance from Docket ID: EPA-HQ-OPP-2016-0545, the First Aid in English and/or Spanish will be added inside label booklet on page 2. A referral statement such as the following will be used.}

See [Below] [Inside] [This] [Label] [Booklet] [for] [additional] [First Aid][/] [Spanish First Aid][:] [and] [additional] [Precautionary Statements][:] [and] [Directions for Use] [(including) [Mitigations for Endangered Species)][,][and] [Storage and Disposal] [Instructions]) [:] [and] [Other] [Required Spanish] [Labeling] [and] [Other] [Use Information] [.]

FIRST AID

If In eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Immediately call a poison control center or doctor for treatment advice.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Have person sip a glass of water if able to swallow.• Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565/800-424-9300.

NOTE TO PHYSICIAN: Treat symptomatically. Consideration should be given to the possibility that overexposure to materials other than Boll Buster may have occurred. No specific antidote is available. Probable mucosal damage may contraindicate the use of gastric lavage.

FOR A MEDICAL EMERGENCY OR HELP WITH ANY SPILL, LEAK, FIRE OR EXPOSURE INVOLVING THIS MATERIAL, CALL DAY OR NIGHT CHEMTREC 1-800-424-9300.

EPA REG. NO. 34704-856

[Label Date Code]

EPA EST. NO. 34704-XX-XXX

NET CONTENTS XX GAL. (XX L)

~~FORMULATED~~ MANUFACTURED FOR

LOVELAND PRODUCTS, INC.®

P.O. BOX 1286 GREELEY, ~~[~~COLORADO~~]~~ [CO] 80632-1286 [USA]

[Note to reviewer: Text in brackets is print optional]

[NOTIF V1 08425]

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
DANGER**

Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear coveralls over short sleeved shirt, chemical resistant gloves, chemical resistant footwear plus socks, goggles and chemical resistant headgear for overhead applications. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirts and short pants,
- Chemical resistant gloves made of any waterproof material (such as Nitrile, Butyl, Neoprene and/or Barrier Laminate),
- Chemical-resistant footwear plus socks,
- Protective eyewear,
- Chemical-resistant headgear for overhead exposure and chemical-resistant apron when cleaning equipment or mixing and loading.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with Boll Buster's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash body thoroughly and put on clean clothing.
- Remove PPE immediately after handling Boll Buster. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate. Do not contaminate water used for irrigation or domestic purposes.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use Boll Buster in a manner inconsistent with its labeling. Do not apply Boll Buster in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations. Read entire label before using Boll Buster.

AGRICULTURAL USE REQUIREMENTS

Use Boll Buster only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry intervals. The requirements in this box only apply to uses of Boll Buster that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. The REI is 72 hours in areas where average rainfall is less than 25 inches per year.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil or water is:

- Coveralls over long-sleeved shirt and long pants,
- Chemical resistant gloves such as any waterproof material, such as polyvinyl chloride, nitrile rubber, or butyl rubber,
- Chemical resistant footwear plus socks, and
- Protective eyewear.
- For overhead exposure, chemical-resistant headgear is also required.

Notify workers of the application by warning them orally and posting warning signs at entrances to treated areas.

SPRAY DRIFT

Aerial Applications:

- Do not release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a fine to medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a fine to medium or coarser droplet size (ASABE S572.1-).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size- Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift.
- Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size- Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT- Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

BOOM-LESS GROUND APPLICATIONS:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift

RELEASE HEIGHT- Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

PRODUCT INFORMATION

Cotton: A foliar spray of Boll Buster will accelerate opening of mature unopened cotton bolls and enhance defoliation which can result in earlier harvest with an increased recoverable yield. Boll Buster treatment on cotton allows increased efficiency from a once-over harvest.

Tobacco (Flue-Cured): A foliar spray of Boll Buster promotes early, uniform "yellowing" of mature tobacco. Boll Buster reduces curing time, allowing more efficient use of curing barn space, and increased control over harvest schedules.

Spray Preparation

Add 1/2 to 3/4 of the required amount of water to the spray tank. Start agitation. Add the required amount of Boll Buster and the remaining amount of water. Prepare only as much spray solution as can be used on the day of mixing. Do not allow spray solution to stand overnight. Do not spill the concentrated product on spray equipment, or any airplane parts.

ANY SPILLS SHOULD BE RINSED IMMEDIATELY WITH PLENTY OF WATER.

Use of a nurse tank is highly recommended for avoiding possible spills of concentrated formulation on spray equipment or any airplane parts.

Tank Mixtures ~~with~~ With Defoliants And Insecticides

Boll Buster is compatible with DFT 6 EC and Malathion. Boll Buster may be applied in sequence or as a tank mixture (DO NOT TANK MIX WITH DESICCANTS IF COTTON IS TO BE SPINDLE HARVESTED). Follow all applicable use precautions and rate per acre recommendations on labels of products applied as tank mixtures or in sequence with Boll Buster. In some cases, slight reduction in boll opening response has been observed when tank mixes with defoliants were used. Good agitation in the spray tank is essential and a tank mixture should not be allowed to stand without agitation for more than 5 to 10 minutes. Read and observe all appropriate label use directions and precautions for the defoliants and insecticides used.

NOTE: UNDER CERTAIN CONDITIONS, TANK MIXTURES OF BOLL BUSTER WITH DESICCANTS CONTAINING SODIUM CHLORATE COULD RESULT IN THE FORMATION OF HYPOCHLOROUS ACIDS, WHICH ON HEATING WILL EMIT TOXIC CHLORIDE FUMES.

Equipment Cleaning

Because of the acidic nature of Boll Buster, prolonged exposure to spray deposit will damage acrylic plastics, certain paints, and metals.

Rinse thoroughly with a detergent and water all exposed acrylic plastic-type materials (e.g., aircraft windshields), and painted surfaces within an hour after exposure to spray deposits.

At the end of each day, rinse thoroughly with a detergent and water all the metal parts of the aircraft and the associated spray equipment exposed to the spray deposits.

USE RESTRICTIONS

- Do not apply Boll Buster through any type of irrigation system.
- Do not plant another crop within 30 days after treatment.

USE PRECAUTIONS

- Avoid spray drift to nearby crops as Boll Buster will cause modifications in plant growth. Plant injury or reduced yields will result.
- Mix only the amount of spray you expect to use each day. Do not allow mixed solution to stand overnight.

COTTON

USE	EXPECTED CONDITIONS	Boll Buster RATE	1 GALLON Boll Buster TREATS	MINIMUM SPRAY VOLUMES (GALS/A)*			APPLICATION TIMING
		Pts/A	Lbs. AI	Acres	Ground	Air****	
Boll Buster Boll Opener**	Hot and dry 80 °F or higher Dry and 75 to 80 °F Cool but above 65 °F or Rank cotton	1.33 2.0 2.66	1.0 1.5 2.0	6 4 3	10.0	2.0	Apply when the number of mature unopened bolls is sufficient to produce the desired crop. See below for test of boll maturity. Treatment uniformly opens bolls 7 to 14 days earlier.

Tank mix of Boll Buster and DFT 6 EC Cotton Defoliant***	High soil moisture or High fertility level or Rank cotton	0.33	0.25	24	10.0	5.0	Apply 4 to 7 days prior to Boll Buster opening application. To be used as a sequential treatment with, not in place of Boll Buster boll opening treatment.
Pre-Conditioner for Defoliant	Hot, dry, above 80 °F Cool, above 65 °F or Rank cotton	0.66 1.33	0.5 1.0	12 6	10.0	2.0	Apply 4 to 7 days prior to defoliant. Enhances top crop defoliation, reducing deterioration of bottom crop and allows for earlier harvest.

*For best performance, by ground or air application, choose equipment and spray volumes that will ensure uniform coverage of foliage and bolls.

****Pretreatment With Defoliants Prior to Boll Buster Boll Opening Treatment:** If the cotton is overly **rank** or laying down in the **middles** and good spray coverage of the bolls with Boll Buster is difficult, a pretreatment with defoliants will be useful to improve boll coverage with Boll Buster. Use dosage rates of Boll Buster specified for boll opening. Read and observe all appropriate label use directions and precautions for the defoliant used.

***Can use up to full label rate for each product.

****In California and Arizona, use a volume of no less than 5.0 gallons per acre for aerial applications.

Boll Maturity

A boll is mature when it is too hard to be dented when squeezed between thumb and fingers, too hard to be sliced with a sharp knife, and when the seed coat becomes light brown in color.

Use Restrictions

- Do not exceed a maximum of 2.0 pounds ethephon active ingredient per acre per year through combined or repeated uses of any ethephon products.
- Do not plant another crop within 30 days after treatment. Small grains planted earlier than 1 month or intercropped with the cotton crop to which Boll Buster will be applied may only be used as cover crops and may not be harvested for food or feed. Boll Buster may cause yellowing and growth inhibition of treated small grains.

Use Precautions

- Boll Opening: Do not tank mix Boll Buster with a desiccant if the cotton is to be spindle harvested.
- Pre-Condition for Defoliation: Do not tank mix Boll Buster with desiccants unless plant desiccation is required. Do not use a defoliant before there is sufficient mature unopened bolls to produce the desired yield (see Product Information section on how to test for boll maturity).
- DO NOT APPLY ~~Boll Buster~~**BOLL BUSTER** IF RAIN IS EXPECTED WITHIN 6 HOURS. Rainfall within 6 hours of application may reduce product performance.

When To Harvest Cotton

Do not harvest cotton sooner than 7 days after a treatment with Boll Buster.

Observe the treated crop and harvest when optimum boll opening has been reached. Too early harvest might reduce the full advantage of the treatment and too late a harvest may result in reduced quality and loss of lint which will drop from the plant.

TOBACCO (FLUE-CURED ONLY)

CROP SITUATION	BOLL BUSTER PTS/A	MINIMUM SPRAY VOLUMES GALS/A	SPECIFIC DIRECTIONS
Directed Spray Application	1.33	50.0	Use drop nozzles. Choose TG or OC spray tips designed to apply 50.0 to 60.0 gpa at 35 to 40 psi and at tractor speed of 2 to 3 mph. Use 2 nozzles per row; one on each side of the row dropped low enough to direct the spray to the leaves to be ripened and harvested. Thorough spray coverage is essential. With a directed spray, be sure to harvest all leaves with 20% or more yellowing.
Over-The-Top Application	1.33 to 2.66	40.0	Treat only when leaves remaining on the stalk are mature. To ensure remaining leaves are mature, test spray several tobacco plants as described under the section "Proper Time of Treatment". Use the lower rate in a normally mature crop when experience indicates that minimum ripening inducement is required. Use the higher rate when the crop is heavy and has a tendency to be more rank or when temperatures are lower than normal. Always test spray to determine if the tobacco is mature enough to respond to treatment with Boll Buster. Apply over-the-top Boll Buster spray as a fine mist using three nozzles (one nozzle tip over the center of the plant, and one on each side) so all leaves are covered thoroughly, similar to the application pattern of systemic sucker control agents. Use a spray pressure of 40 to 60 psi.

Use Restrictions

- Do not use Boll Buster with additives other than referenced on this label.
- Do not plant another crop within 30 days after treatment.
- Do not exceed a maximum of 2.0 pounds ethephon active ingredient per acre per year through repeated uses of any ethephon products.

Use Precautions

- Application of Boll Buster to immature leaves can result in unsatisfactory coloring, weight loss and reduced leaf quality.
- Allowing the crop to over ripen in the field after using Boll Buster may cause some reduction in yield and quality.
- Treatment before an anticipated major storm could prevent harvest and result in crop loss.
- Avoid application of Boll Buster IF RAIN IS EXPECTED WITHIN 6 HOURS. Rainfall within 6 hours of application may reduce product performance.

Application Timing

Successful results with Boll Buster call for treatment when leaves are mature, not overly rank green when sprayed. To easily determine the proper treatment timing and the number of leaves per stalk ready for harvest, test spray several plants in more than one location in each field and observe the response. Mature leaves will begin to yellow in 24 to 72 hours. Test leaves that fail to yellow in 72 hours are not mature and are not ready for Boll Buster treatment. Wait a few days to permit further natural maturing, then make another test spray or "maturity" check.

To avoid quality loss and or possible leaf drop, harvest any yellowed leaves prior to application. Use lower rates under most conditions. Limit use of higher rates to cool (below 65 °F at the time of treatment), slow ripening conditions.

When you have confirmed the desired number of leaves per plant that will color, you can determine the number of acres to treat in order to fill the barn.

Prepare your test spray by mixing one teaspoon of Boll Buster in one quart of water. Spray each test plant with about 1.0 ounce of this mixture, covering all leaves with a fine mist. Boll Buster will not color immature sprayed leaves.

When To Harvest

All mature, sprayed leaves will begin to color within 24 to 72 hours after application of Boll Buster. The yellowing process is weather dependent; cool weather will delay, while hot, sunny weather can speed up the process. Harvest treated tobacco when leaves have reached the desired color intensity.

Harvest can commence 48 hours after application of Boll Buster. To determine harvest timing and avoid quality loss or leaf drop, closely monitor treated crop and weather conditions.

CURING BOLL BUSTER TREATED TOBACCO

Curing procedures are as much an art as a science and each cure must be judged on the basis of tobacco condition, interval between treatment and harvest, weather and type of curing facility before prescription temperature and ventilation schedules can be established. To obtain maximum quality, care must be taken to observe and control the curing process closely, especially during the late "coloring" and early "drying" stages of the leaf.

Boll Buster treated tobacco will have started the coloring process when harvested, reducing the time required in the coloring phase of curing. Treated tobacco should be dried faster. If tobacco leaves are green or contain some green when harvested, it may be necessary to color them for a few hours. If the leaves are completely yellow, temperature and ventilation must be adjusted in a manner to dry the tobacco as fast as possible without scalding. Once the leaf is dried (3/4 dry), you should follow normal procedures for curing. Since Boll Buster treated leaves cure faster, treated and untreated leaves should not be cured together in the same barn.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal.

If container is broken or contents have spilled, follow all precautions indicated above and clean up immediately. Before cleaning up, put on full-length trousers, long-sleeved shirt, protective gloves, and goggles or face shield. Soak up spill with absorbent media such as sand, earth or other suitable material and dispose of waste at an approved waste disposal facility.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: [Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.agrecycling.org. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.]

[For packages up to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

[For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete

revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

[For packages greater than 56 gallons: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. **For refillable containers:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.]

[For refillable containers:

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.]

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY

FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Boll Buster is a registered trademark of Loveland Products, Inc.

{OPTIONAL GRAPHICS}

{Note to Reviewer: The below graphic(s) may appear anywhere on the label. Further, the below graphic(s) may appear in various colors or black/white; not to impact legibility.}



{Option Marketing Claims, Logos and Graphics}

{Note to Reviewer: These logos and graphics or their respective components may be used throughout this label. The optional claims next to the logo is [PGR]. The logo may be used with one or more of the optional claims, or it may be used by itself.}

