

34704-861

10/9/2009

Jacket  
1041



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

EPA Reg. Number:

34704-861

Date of Issuance:

OCT 9 2009

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:

Rifle Herbicide

Name and Address of Registrant (include ZIP Code):

Loveland Product Inc.  
7251 W 4<sup>th</sup> Street  
P.O. Box 1286  
Greeley, CO 80632-1286

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.

Signature of Approving Official:

Joanne Miller  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

Date:

OCT 9 2009

2) Per the acute toxicity review, the Hazards to Humans and Domestic Animals must be revised to read:

**“WARNING**

Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Avoid contact with skin.”

3) The text “protective eyewear” appearing in the handler PPE section should be revised to read “goggles or faceshield.”

4) Per the RED label table, the text in **bold type** below must be added to the following User Safety Requirements:

“...If no such instructions for washables **exist**, use detergent and hot water...”

5) The text in bold type must be added to the User Safety Recommendation text currently on the label:

“User should remove clothing/**PPE** immediately if pesticide gets inside.”

6) Per the RED, the text “Do not contaminate water when cleaning equipment or disposing of equipment washwater” currently appearing in the Environmental Hazard section of the label should be revised to read “Do not contaminate water when disposing of equipment washwater or rinsate.”

7) The double notification text (“Notify workers of application by warning them orally and by posting warning signs at entrances to treated areas”) appearing in the Agricultural Use Requirements box is no longer needed and should be deleted from this section of the label.

8) Per the revised RED label table, the following revisions are needed to the directions for use and any conflicting text must be deleted:

-For all uses (except where the label restricts the use to one application per year), the rate restriction text should be revised to read:

“Maximum single application rate is 1.0 lbs ae per acre.  
Maximum annual application rate is 2.0 lbs ae per acre per year.”

**-Sugarcane, Pasture, Hay, Rangeland and General Farmstead, Conservation Reserve Program, Preplant Directions (Post Harvest/Fallow/Crop Stubble/Set-A-Side), Cropping Restrictions (Pages 35-37 of label):**

Rates in excess of 1 lb ae dicamba per acre per application (including spot treatment) are

Page 3  
EPA Reg. 34704-861

unacceptable and must be deleted from the label.

-PHI text must read:

"The PHI for sorghum grain is 30 days.

The PHI for sorghum forage is 0 days.

The PHI for sorghum fodder is 30 days.

The PHI for wheat grain is 7 days.

The PHI for barley is 7 days.

The PHI for oat grain is 7 days.

The PHI for sugarcane is 87 days.

The PHI for grass forage is days.

The PHI for grass hay is 7 days."

9) Add "or rinseate" after "washwaters" to the Environmental Hazards section.

10) Change "General Information" to "Product Information" on page 5. Delete "General" from "General Weed List" on page 8 and throughout the label. Delete "General" from "General Farmstead". On page 35, change "the following recommendations" to "the following directions" under cropping restrictions. Change Restictions to Restrictions.

A stamped copy of the label is enclosed for your records. You must submit one copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of the product constitutes acceptance of these conditions. This label supersedes all other previously accepted labels. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Stamped 6-11-09  
4041

EDITOR's NOTE: EXP Label for the Dicamba RED 6-8-09



**ACCEPTED  
with COMMENTS**  
In EPA Letter Dated:  
OCT - 9 2009  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

34704-861

For Weed Control in Corn, Sorghum, Small Grains, Pasture, Hay, Rangeland, General Farmstead (Non-Cropland), Fallow, Cotton, Sugarcane, Asparagus, Turf, and Grass Seed Crops

ACTIVE INGREDIENT:

Dimethylamine salt of dicamba* .....	48.2%
OTHER INGREDIENTS: .....	51.8%
TOTAL .....	100.0%

\*This product contains 40.0% 3,6-dichloro-o-anisic acid (dicamba) or 4 pounds per gallon.

**KEEP OUT OF REACH OF CHILDREN**

**WARNING-AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF SWALLOWED:	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-866-944-8565 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Probable mucosal damage may contraindicate the use of gastric lavage.	

See inside booklet for additional PRECAUTIONARY STATEMENTS.

EPA Reg. No. 34704-861

EPA Est. No: 0-12750-MO-001

NET CONTENTS: \_\_\_\_\_ Gals.

EXP 060809 jt

5041

MANUFACTURED FOR:  
Loveland Products, Inc.  
P.O. Box 1286  
Greeley, CO 80632-1286

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### WARNING

Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wear protective eyewear (goggles or face shield). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are butyl rubber, natural rubber, neoprene rubber or nitrile rubber. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

#### All mixers, loaders, and applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Chemical-resistant gloves (except for applicators using ground boom equipment, pilots and flaggers)
- Shoes plus socks, and
- Protective eyewear.

#### See engineering controls for additional requirements and exceptions.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6).**

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Apply this product only as directed.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- 1. Coveralls worn over short-sleeved shirt and short pants.**
- 2. Chemical resistant footwear plus socks**
- 3. Chemical resistant gloves made of any waterproof material.**
- 4. Chemical resistant headgear for overhead exposure.**
- 5. Protective eyewear**

**Notify workers of application by warning them orally and by posting warning signs at entrances to treated area.**

**NON-AGRICULTURAL USE REQUIREMENTS**

70941

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. This product may not be mixed, loaded, or used within 50 feet of all wells including abandoned wells, drainage wells, and sinkholes.

**PESTICIDE STORAGE:** Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

**PESTICIDE DISPOSAL:** Triple rinse pesticide from containers and use rinsates in the pesticide application. Wastes which cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Non-refillable containers (1, 2.5, 30 & 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**(non-refillable <5 gallons):** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**(non-refillable >5 gallons):** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle inside of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable container (250 gallon & bulk):** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinseate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.

**GENERAL INFORMATION**

The following directions apply to all uses of RIFLE. Additional precautions and restrictions will be found in each specific use section. Do not treat irrigation ditches or water used for crop irrigation or domestic uses. Do not apply this product through any type of irrigation system.

**MIXING AND APPLICATION**

Unless otherwise specified under the individual use headings of this label, the following directions apply to all crop and noncrop uses of RIFLE. Refer to individual use sections for additional precautions, restrictions, application rates and timings.

RIFLE is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (see COMPATIBILITY TEST below) should be made prior to tank mixing.

Ground or aerial application equipment which will give good spray coverage of weed foliage should be used. However, do not use aerial application equipment if spray particles can be carried by wind into areas where sensitive crops or plants are growing or when temperature inversions exist.

Apply 3 to 50 gallons of diluted spray per treated acre when using ground application equipment, or 1 to 10 gallons of diluted spray per treated acre (2 to 20 gallons of diluted spray per acre for preharvest uses) in a water-based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, RIFLE should not be applied during periods of gusty wind or when wind is in excess of 15 mph.

Avoid disturbing (e.g., cultivating or mowing) treated areas for at least 7 days following application.

**GROUND AND SURFACE WATER PROTECTION**

1) Point source contamination – To prevent point source contamination, do not mix or load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below. Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment. Care must be taken when using this product to prevent: a) back-siphoning into wells, b) spills or c) improper disposal of excess pesticide, spray mixtures or rinsates. Check valves or antisiphoning devices must be used on all mixing equipment.



2) Movement by surface runoff or through soil – Do not apply under conditions which favor runoff. Do not apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for ground water contamination. Ground water contamination may occur in areas where soils are permeable or coarse and ground water is near the surface. Do not apply to soils classified as sand with less than 3% organic matter and where ground water depth is shallow. To minimize the possibility of ground water contamination, carefully follow application rate recommendations as affected by soil type in the general information section of this label.

3) Movement by water erosion of treated soil – Do not apply or incorporate this product through any type of irrigation equipment nor by flood or furrow irrigation. Ensure treated areas have received at least one-half inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

**SENSITIVE CROP PRECAUTIONS**

RIFLE may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes, and other broadleaf plants when contacting their roots, stems or foliage. These plants are most sensitive to RIFLE during their development or growing stage. Follow the precautions listed below when using RIFLE.

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of RIFLE with the roots of desirable plants such as trees and shrubs.
- Avoid making applications when spray particles may be carried by air currents to areas where sensitive plants are growing, or when temperature inversions exist. Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive plants. Leave an adequate buffer zone between area to be treated and sensitive plants. Coarse sprays are less likely to drift out of the target area than fine sprays.
- Use coarse sprays to avoid potential herbicide drift. Select nozzles which are designed to produce minimal amounts of fine spray particles. Examples of nozzles designed to produce coarse sprays via ground applications are Delavan Raindrops, Spraying Systems XR flat fans or large capacity flood nozzles such as D10, TK10, or greater capacity tips. Keep the spray pressure at or below 20 psi and the spray volume at or above 20 gpa, unless otherwise required by the manufacturer of drift-reducing nozzles. Consult with your spray nozzle supplier concerning the choice of drift-reducing nozzles.
- Agriculturally approved drift-reducing additives may be used.
- Do not apply RIFLE adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85°F as drift is more likely to occur.
- To avoid injury to desirable plants, equipment used to apply RIFLE should be thoroughly cleaned (see PROCEDURE FOR CLEANING SPRAY EQUIPMENT) before reusing to apply any other chemicals.

All crop uses of RIFLE are intended for a normal growing interval between planting and harvest. No crop rotation restrictions exist if normal harvest of treated crop has occurred. If this interval is shortened, such as in cover crops that will be plowed under, do not follow up with the planting of a sensitive crop.

Crops growing under stress conditions such as drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied. Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations. Tank mix recommendations are for use only in states where the tank mix product and application site are registered.

**BAND TREATMENTS**

RIFLE may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per treated acre} = \text{Band rate per treated acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast volume per treated acre} = \text{Band volume per treated acre}$$

### COMPATIBILITY TEST

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

Amount of Herbicide to Add to One Pint of Spray Carrier  
(Assuming Volume is 25 Gallons per Acre)

HERBICIDE FORMULATIONS	RATE PER ACRE	LEVEL TEASPOONS
Dry	1 lb.	1 1/2
Liquid	1 pt.	1/2

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested spray mix is compatible. Usually, incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Rerun the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of fluid fertilizer).

### PROCEDURE FOR CLEANING SPRAY EQUIPMENT

The steps listed below are suggested for thorough cleaning of spray equipment following applications of RIFLE or tank mixes of RIFLE plus 2,4-D amine:

1. Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
2. Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Remove the nozzles and screens and flush the system with two full tanks of water.
5. The steps listed below are suggested for thorough cleaning of spray equipment used to apply RIFLE as a tank mix with wettable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible formulations. RIFLE tank mixes with water-dispersible formulations require the use of a water/detergent rinse:
6. Complete step 1.
7. Fill tank with water while adding 2 lbs. of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
8. Flush the detergent solution out of the spray tank through the boom.

9. Repeat step 1, and follow with steps 2,3, and 4.

**GENERAL WEED LIST**

This is a general list of weeds which may be treated with RIFLE in accordance with this label as recommended under the RATES AND TIMING sections of the individual use headings. Proper usage of this product will give control of growth suppression of any annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species including:

**ANNUALS**

Amaranth, Spiny (Spiny Pigweed)  
 Aster, Slender  
 Bedstraw

Beggarweed, Florida  
 Broomweed, Common  
 Buckwheat, Wild  
 Buffalobur  
 Burclover, California  
 Burcucumber  
 Buttercup Roughseed  
 Carpetweed  
 Catchfly, Nightflowering  
 Chamomile, Corn  
 Chickweed, Common  
 Clovers (Annual)  
 Cockle, Corn  
 Cockle, Cow  
 Cocklebur, Common  
 Croton, Tropic  
 Croton, Woolly  
 Daisy, English  
 Evening Primrose, Cutleaf  
 Fleabane, Annual  
 Goosefoot, Nettleleaf  
 Henbit  
 Jimsonweed  
 Knotweed  
 Kochia  
 Ladysthumb  
 Lambsquarters, Common  
 Lambsquarters (triazine resistant)  
 Lettuce, Prickly  
 Mallow, Common  
 Mallow, Venice  
 Mare's Tail (Horseweed)  
 Mayweed  
 Moringglory, Ivyleaf  
 Moringglory, Tall  
 Mustard, Tansy  
 Mustard, Wild

Mustard (Yellowtops)  
 Nightshade, Black  
 Pennycress, Field (Fanweed, Frenchweed, Stinkweed)  
 Pepperweed, Virginia (Peppergrass)  
 Pigweed, Prostrate  
 Pigweed, Redroot (Carelessweed)  
 Pigweed, Rough  
 Pigweed, Smooth  
 Pigweed (triazine resistant)  
 Pigweed, Tumble  
 Poorjoe  
 Puncturevine  
 Purslane, Common  
 Pusley, Florida  
 Radish, Wild  
 Ragweed, Common  
 Ragweed, Giant (Buffaloweed)  
 Ragweed, Lance-Leaf  
 Rubberweed, Bitter (Bitterweed)  
 Sesbania, Hemp  
 Shepherdspurse  
 Sicklepod  
 Sida, Prickly (Teaweed)  
 Smartweed, Green  
 Smartweed, Pennsylvania  
 Sneezeweed, Bitter  
 Sowthistle, Annual  
 Sowthistle, Spiny  
 Spikeweed, Common  
 Spurge, Prostrate  
 Spurry, Corn  
 Starbur, Bristly  
 Sumpweed, Rough  
 Sunflower, Common (Wild)  
 Sunflower, Volunteer  
 Thistle, Russian  
 Velvetleaf  
 Waterhemp  
 Waterprimrose, Winged  
 Wormwood, Annual

**BIENNIALS**

Burdock, Common	Plantain, Bracted
Carrot, Wild (Queen Anne's Lace)	Ragwort, Tansy
Cockle, White	Starthistle, Yellow
Evening Primrose, Common	Sweetclover
Geranium, Carolina	Teasel
Gromwell	Thistle, Bull
Knapweed, Diffuse	Thistle, Milk
Knapweed, Spotted	Thistle, Musk
Mallow, Dwarf	Thistle, Plumeless

**PERENNIALS**

*Alfalfa	Milkweed, Climbing
Artichoke, Jerusalem	Milkweed, Common
Aster, Spiny	Milkweed, Honeyvine
Aster, Whiteheath	Milkweed, Western Whorled
Beadstraw, Smooth	Nettle, Stinging
Bindweed, Field	Nightshade, Silverleaf (White Horsenettle)
Bindweed, Hedge	Onion, Wild
Blueweed, Texas	*Plantain, Broadleaf
*Bursage (Bur Ragweed, Lakeweed, Povertyweed)	Plantain, Buckhorn
Bursage, Woollyleaf (Lakeweed)	Pokeweed
Buttercup, Tall	Ragweed, Western
Campion, Bladder	Redvine
Chickweed, Field	Sericia Lespedeza
Chickweed (Mouseear, Canada)	Smartweed, Swamp
Chicory	Snakeweed, Broom
*Clover, Hop	*Sorrel, Red (Sheep Sorrel)
*Dandelion, Common	Sowthistle
*Dock, Broadleaf (Bitterdock)	Sowthistle, Perennial
*Dock, Curly	Spurge, Leafy
Dogbane, Hemp	Sundrop, Halfshrub (Evening Primrose)
*Dogfennel (Cypressweed)	Thistle, Canada
Fern, Bracken	Toadflex, Dalmation
Garlic, Wild	Tropical Soda Apple
Goldenrod, Canada	Trumpet creeper (Buckvine)
Goldenrod, Missouri	Vetch
Goldenweed, Common	Waterhemlock
Hawkweed	Waterprimrose, Creeping
Henbane, Black	*Woodsorrel, Creeping Common Yellow
Horsenettle, Carolina	Wormwood, Common
Ironweed	Wormwood, Louisiana
Knapweed, Black	*Yankeeweed
Knapweed, Russian	Yarrow, Common

\*Noted perennials may be controlled using RIFLE at rates lower than those recommended for other listed perennial weeds. (See APPLICATION RATES AND TIMING section in this label.)

**WOODY**

- Alder
- Ash
- Aspen
- Basswood
- Beech
- Birch
- \*Blackberry
- \*Blackgum
- \*Cedar
- Cherry
- Chinquapin
- Cottonwood
- \*Creosotebush
- Cucumbertree
- \*Dewberry
- \*Dogwood
- Elm
- Grape
- \*Hawthorn, (Thornapple)
- Hemlock
- Hickory
- Honeylocust
- Honeysuckle
- Hornbeam
- Huckleberry
- Huisache
- Ivy, Poison
- Kudzu
- \*Growth Suppression
- Locust, Black
- Maple
- Mesquite
- Oak
- Oak, Poison
- Olive, Russian
- Persimmon, Eastern
- Pine
- \*Plum Sand (Wild Plum)
- Poplar
- Rabbitbrush
- \*Redcedar, Eastern
- \*Rose, McCartney
- \*Rose, Multiflora
- Sagebrush, Fringe
- Sassafras
- Serviceberry
- Spicebush
- Spruce
- Sumac
- \*Sweetgum
- Sycamore
- Tarbrush
- Willow
- Witchhazel
- \*Yaupon
- \*Yucca

**FIELD, SEED\*, POPCORN\* AND SILAGE CORN**

\*Do not apply RIFLE to seed corn or popcorn without first verifying with your local seed corn company (supplier) the RIFLE selectivity on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties.

Observe all precautions, mixing and application instructions.

RIFLE is not registered for use on sweet corn.

Direct contact of RIFLE with corn seed must be avoided. If corn seeds are less than 1 1/2 inches below the surface, delay application until corn has emerged.

Up to 2 applications of RIFLE may be made during a growing season. Do not exceed a total of 1 1/2 pints of RIFLE per treated acre per crop year. Allow two weeks or more between applications. See appropriate section for rate information. For combination options or sequential treatments, refer to appropriate section.

**RIFLE contains 0.5 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.**

Applications of RIFLE to corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3 to 7 days. Cultivation should be delayed until after corn is growing normally to avoid breakage.

Agriculturally approved surfactants or sprayable fertilizers (1/2 to 1 gallon per acre of 28%, 30% or 32% urea ammonium nitrate or 2.5 pounds per acre spray grade ammonium sulfate<sup>1</sup>) may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

<sup>1</sup>Not for use in California.

Do not use adjuvants containing penetrants such as petroleum-based oils after crop emergence or crop injury may result.

Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.

Several synthetic pyrethroid insecticides are labeled for tank mix applications of DICAMBA. Refer to their label for specific recommendations.

#### **WEEDS CONTROLLED**

RIFLE will control many annual broadleaf weeds or give growth suppression of many perennial broadleaf weeds commonly found in corn. (Refer to the GENERAL WEED LIST.)

For best performance, make application when weeds have emerged and are actively growing.

Preemergence control of cocklebur, velvetleaf, and jimsonweed may be reduced if conditions such as low temperature or lack of soil moisture cause delayed or deep germination of weeds.

#### **PREPLANT/PREEMERGENCE IN NO-TILLAGE CORN**

Applications of RIFLE may be made before, during, or after planting to emerged and actively growing broadleaf weeds. Apply RIFLE at 1 pint per treated acre on medium or fine textured soils containing 2% or greater organic matter. Use 1/2 pint per treated acre on coarse textured soils (sand, sandy loam, and loamy sand) or medium and fine textured soils with less than 2% organic matter.

When planting into a legume sod (e.g., alfalfa or clover), apply RIFLE after 4 to 6 inches of regrowth has occurred.

#### **PREEMERGENCE IN CONVENTIONAL OR REDUCED TILLAGE CORN**

RIFLE may be applied after planting and prior to corn emergence. Application at 1 pint per treated acre may be made to medium or fine textured soils which contain 2% or greater organic matter. DO NOT apply to coarse textured soils (sand, sandy loam, and loamy sand) until after crop emergence (see EARLY POSTEMERGENCE uses below).

Preemergence application of RIFLE does not require mechanical incorporation to become active. A shallow mechanical incorporation is recommended if application is not followed by adequate rainfall or sprinkler irrigation. Avoid tillage equipment (e.g., drags, harrows) which concentrate treated soil over seed furrow.

**EARLY POSTEMERGENCE (ALL TILLAGE SYSTEMS)  
(SPIKE THROUGH 8 INCH TALL CORN)**

RIFLE at 1 pint per treated acre may be applied during the period from corn emergence through the five-leaf stage or 8 inches tall, whichever comes first. Reduce the rate to ½ pint per treated acre if corn is growing on coarse textured soils (sand, sandy loam, and loamy sand). See LATE POSTEMERGENCE APPLICATIONS given below if the 6<sup>th</sup> true leaf is emerging from whorl or corn is greater than 8 inches tall.

**LATE POSTEMERGENCE (ALL TILLAGE SYSTEMS)  
(8 TO 36 INCH TALL CORN)**

Application of RIFLE at ½ pint per treated acre may be made from 8 to 36 inch tall corn or 15 days before tassel emergence, whichever comes first. For best performance, make applications when weeds are less than 3 inches tall.

Make directed spray application when (1) corn leaves prevent proper spray coverage; (2) sensitive crops are growing nearby; (3) tank mixing with 2,4-D.

DO NOT apply RIFLE when soybeans are growing nearby if any of these conditions exist:

- corn is more than 24 inches tall
- soybeans are more than 10 inches tall
- soybeans have begun to bloom

**OVERLAY (SEQUENTIAL) TREATMENTS**

RIFLE may be applied to ground previously treated with one or more of the following herbicides:

acetochlor (Surpass™, Harness® Plus)  
alachlor (Lasso®, Lasso® MT, Partner®)  
atrazine  
Broadstrike™ + Dual®  
Broadstrike™ Plus  
Bullet®  
butylate (Sutan® +)  
Clarity®  
dimethenamid (Frontier®)  
EPTC (Eradicane®)

glyphosate (Gly Star™ Original, Roundup®)  
Guardsman®  
halosulfuron (Battalion®, Permit®)  
Lariat®  
Dicamba plus Atrazine (Dicambazine™)  
metolachlor  
paraquat (Gramoxone®)  
pendimethalin (Prowl®)  
propachlor (Ramrod®)  
simazine (Princep®)

Apply RIFLE at ½ pint per treated acre to ground previously treated with full rates of Clarity® or Dicamba plus Atrazine herbicides. Allow at least 2 weeks between applications. Read and follow label directions for each of the above products.

**TANK MIX TREATMENTS FOR CORN**

RIFLE may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, and other restrictions.

**RATES AND TIMINGS**

RIFLE Plus	Preplant Preemergent (No Tillage Corn)	Preemergent (Conventional or Reduced Tillage Corn)	Early Postemergent (All Tillage Systems)	Late Postemergent (All Tillage Systems)	Additional Directions
ACCENT® (nicosulfuron)			½ - 1 oz. a.i./A	½-1 oz. a.i./A. To improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall.	Application may be made to emerged weeds before corn is greater than 24 inches tall. Use non-ionic surfactant at 0.25% (v/v) with this tank mixture.
Atrazine	1 ¼ -2 lbs. a.i./A	1 ¼-2 lbs. a.i./A	1 ¼-2 lbs. a.i./A. Crop oil concentrate may be used with this mixture if corn is 5 inches or less in height.	1 ¼-2 lbs. a.i./A. Do not apply if corn is greater than 12 inches tall.	Application may be made before grasses are 1 ½ inches tall. Follow all state and federal restrictions pertaining to atrazine applications.
BEACON® (primisulfuron)			0.31-0.62 oz. a.i./A	0.31-0.62 oz. a.i./A. To improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall.	Application may be made to emerged weeds when corn is 4 to 24 inches tall. Use non- ionic surfactant at 0.25% (v/v) with this tank mixture.
DUAL II S-metolachlor	1 to 2 lbs. a.i./A	1 to 2 lbs. a.i./A. Use only on fine or medium textured soils with 2.5% or greater organic matter.	1 to 2 lbs. a.i./A		Application may be made before grasses reach the 2-leaf stage and before corn is greater than 3 inches tall.
FRONTIER® (dimethenamid)	13-25 fl. oz. a.i./A	13-25 fl. oz./A. Use only on fine or medium textured	13-25 fl. oz. a.i./A		Application may be made up to 8-inch tall corn.



RIFLE Plus	Preplant Preemergent (No Tillage Corn)	Preemergent (Conventional or Reduced Tillage Corn)	Early Postemergent (All Tillage Systems)	Late Postemergent (All Tillage Systems)	Additional Directions
		soils with 2.5% or greater organic matter.			This treatment must be combined with a herbicide that provides postemergence control of grass weeds if they are greater than 1-inch tall at the time of application.
FRONTIER® 6.0 (dimethenamid)	16 to 32 fl. oz./A	16 to 32 fl. oz./A. Use only on fine or medium textured soils with 2.5% or greater organic matter.	16 to 32 fl. oz./A		Application may be made up to 8-inch tall corn. This treatment must be combined with a herbicide that provides postemergence control of grass weeds if they are greater than 1-inch tall at the time of application.
GRAMOXONE® (paraquat)	¼-1 lb. a.i./A	¼-1 lb. a.i./A			Application may be made to emerged weeds but prior to corn emergence.
HARNESS® PLUS or SURPASS™ (acetochlor)	1 ½-3 lbs. a.i./A	1 ½-3 lbs. a.i./A. Use only on fine or medium textured soils with 2.5% or greater organic matter.			Application should be made prior to corn emergence.
LASSO® (alachlor)	1 ½-4 lbs. a.i./A	1 ½-4 lbs. a.i./A. Use only on fine textured soils with greater than 2.5% organic matter.	1 ½-4 lbs. a.i./A		Application may be made before grasses reach the 2-leaf stage and before corn is greater than 3 inches tall. If microencapsula ted forms of

RIFLE Plus	Preplant Preemergent (No Tillage Corn)	Preemergent (Conventional or Reduced Tillage Corn)	Early Postemergent (All Tillage Systems)	Late Postemergent (All Tillage Systems)	Additional Directions
					alachlor are used (Lasso <sup>®</sup> MT, Partner <sup>®</sup> ), applications must be made prior to grass emergence.
PRINCEP <sup>®</sup> (simazine)	2-3 lbs. a.i./A	2-3 lbs. a.i./A			Application may be made prior to corn or weed emergence.
PROWL <sup>®</sup> (pendimethalin)		$\frac{3}{4}$ -1 $\frac{1}{2}$ lbs. a.i./A. Use only on fine or medium textured soils with 2.5% or greater organic matter.	$\frac{3}{4}$ -1 $\frac{1}{2}$ lbs. a.i./A		Application may be made immediately after planting but prior to weed emergence. Corn should not be beyond the 2-leaf stage of growth.
GLY STAR <sup>™</sup> ORIGINAL or ROUNDUP <sup>®</sup> (glyphosate)	1-3 lbs. a.i./A	1-3 lbs. a.i./A			Application may be made to emerged weeds but prior to corn emergence.
STINGER <sup>™</sup> (clopyralid)			0.035-0.07 lbs. a.i./A	0.035-0.07 lbs. a.i./A	Application may be made any time after corn emergence through 24-inch tall corn. Use drop nozzles to direct spray after corn exceeds the 8-inch stage. Apply when the majority of the thistle plants have emerged and are at least 4 inches in height, but before bud stage. Use

RIFLE Plus	Preplant Preemergent (No Tillage Corn)	Preemergent (Conventional or Reduced Tillage Corn)	Early Postemergent (All Tillage Systems)	Late Postemergent (All Tillage Systems)	Additional Directions
					higher rates listed for stand reduction of larger thistle plants or heavier infestations. Lower rates listed may provide seasonal thistle suppression only.
TOUGH® 3.75 EC (pyridate)			0.47 lb. a.i./A	0.47-0.94 lb. a.i./A	Applications may be made to emerged, actively growing weeds. Directed applications are recommended when corn is large enough to prevent proper spray coverage.
2,4-D	¼-1/2 lb. a.i./A	¼-1/2 lb. a.i./A	Not recommended	1/8 lb. a.i./A	Drop pipes are to be used when corn height is 8 inches or greater. Keeping the spray off the corn leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

### SORGHUM (MILO)

Observe all precautions, including the reference to crops growing under stress.

Read and follow MIXING AND APPLICATION instructions.

Applications of RIFLE to sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 to 14 days.

Do not graze or feed treated sorghum forage or silage prior to mature grain stage. If sorghum is grown for pasture or hay, refer to the PASTURE use section. Do not apply RIFLE to sorghum grown for seed production.

Make no more than one application per growing season.

**RIFLE contains 0.5 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.**

### WEEDS CONTROLLED

RIFLE, when applied at the recommended rate for sorghum, will control many actively growing annual broadleaf weeds and will reduce competition from established perennial broadleaf weeds as well as control their seedlings. (Refer to GENERAL WEED LIST.)

### RATES AND TIMINGS

RIFLE may be applied to emerged and actively growing weeds at least 15 days prior to planting. Postemergence application of RIFLE must be made after sorghum is in the spike stage (all sorghum emerged) but before sorghum is 15 inches tall. For best performance, make applications when sorghum is in the 3 to 5 leaf stage and weeds are small (less than 3 inches tall). Use drop pipes (drop nozzles) if sorghum is taller than 8 inches. Keeping the spray off the sorghum leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

**BROADCAST RATE PER TREATED ACRE: ½ pint (1/4 lb. a.i.)**

### TANK MIX TREATMENTS

#### RIFLE PLUS ATRAZINE

For improved control of emerged, actively growing broadleaf weeds including triazine-resistant species and added suppression of perennial broadleaf weeds, tank mix ½ pint RIFLE with 0.5 to 1.25 lbs. a.i. atrazine per treated acre. For control of grasses (less than 1.5 inches tall), tank mix ½ pint RIFLE with 2 lbs. a.i. atrazine per treated acre. For best performance and minimal crop injury, make application when sorghum is 3-8 inches tall and when broadleaf weeds are small (less than 6 inches tall). Application of atrazine must be made before sorghum is beyond 12 inches tall. The atrazine rate will depend upon soil texture and length of residual weed control desired. Follow all State and Federal restrictions pertaining to atrazine applications.

#### RIFLE PLUS BROX™ 2EC HERBICIDE or BUCTRIL®

For improved control of broadleaf weeds, tank mix ½ pint RIFLE with 1-1 ½ pints BROX™ 2EC Herbicide or Buctril® per treated acre. Make application at 4-leaf to 15-inch tall sorghum. Use drop nozzles to direct spray beneath sorghum leaves when sorghum is greater than 8 inches tall.

Read and follow the label of each tank mix product used for precautionary statements, directions for use, application rates and timings and other restrictions.

### OVERLAY (SEQUENTIAL) TREATMENTS

RIFLE may be applied to ground previously treated with one or more of the following herbicides registered for use in sorghum:

Herbicide	Maximum Rate Per Treated Acre (lbs. a.i.)
alachlor (Lasso <sup>®</sup> ) (Screen <sup>®</sup> -treated seed)	4
atrazine <sup>1</sup>	2.5
metolachlor ( ) (Concep <sup>®</sup> -treated seed)	1.67
propachlor (Ramrod <sup>®</sup> )	5

<sup>1</sup>Maximum use rate for atrazine is determined by soil type, tillage practices, surface residue, and state or local restrictions. Follow the more restrictive requirements when determining the maximum use rate for atrazine.

### PREHARVEST USES (FOR USE ONLY IN THE STATES OF TEXAS AND OKLAHOMA)

RIFLE may be applied for weed suppression any time after the sorghum has reached the soft dough stage. An agriculturally approved surfactant may be used to improve performance. For aerial application use at least 2 gallons of water-based carrier per treated acre. Delay harvest until 30 days after treatment.

BROADCAST RATE PER TREATED ACRE: ½ pint (1/4 lb a.i.)

### SMALL GRAINS (WHEAT, BARLEY AND OATS) NOT UNDERSEEDED TO LEGUMES

Observe all precautions. Read and follow cleaning, mixing and application instructions. If small grains are used for pasture or hay, the following restrictions apply:

- Animals cannot be removed from treated area for slaughter prior to 30 days after last application.
- There is no waiting period between treatment and grazing for non-lactating dairy animals.
- Treated areas may not be grazed by lactating dairy animals before 7 days after treatment.
- Do not harvest hay from treated areas before 37 days after treatment.

**RIFLE contains 0.5 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.**

Note: Observe all precautions and restriction on the labels of products used in tank mix treatments.

### WEEDS CONTROLLED

RIFLE or combinations with listed tank mix partners will provide control or suppression of the annual broadleaf weeds listed below. For improved control of listed weeds, it is recommended that RIFLE be applied in a tank mix with other herbicides. Refer to specific crop for tank mix options.

Alkanet<sup>1</sup>  
Bedstraw, Catchweed<sup>1</sup>

Mustard, Blue (Purple)<sup>1</sup>  
Mustard, Tansy

Bindweed, Field <sup>2</sup>	Mustard, Treacle <sup>1</sup>
Buckwheat, Tartary	Mustard, Tumble (Jim Hill) <sup>1</sup>
Buckwheat, Wild	Mustard, Wild <sup>1</sup>
Carpetweed <sup>1</sup>	Nightshade, Black
Chamomile, Corn	Nightshade, Cutleaf <sup>1</sup>
Chervil, Bur <sup>1</sup>	Nightshade, Silverleaf <sup>2</sup> (White Horsenettle)
Chickweed, Common <sup>1</sup>	Pennycress, Field (Fanweed, Frenchweed, Stinkweed)
Cockle, Corn	Pigweed, Redroot (Carelessweed)
Cockle, Cow	Pigweed, Rough
Cocklebur, Common	Pigweed, Tumble
Cornflower (Bachelorbutton) <sup>1</sup>	Pineappleweed <sup>1</sup>
Dandelion, Common <sup>2</sup>	Plantain, Broadleaf <sup>2</sup>
Dock, Curly <sup>2</sup>	Poppy, Red Horned <sup>1</sup>
Dragonhead, American <sup>1</sup>	Puncturevine <sup>1</sup>
Evening Primrose, Cutleaf <sup>1</sup>	Purslane, Common <sup>1</sup>
Falseflax, Smallseeded <sup>1</sup>	Radish, Wild <sup>1</sup>
Fiddleneck (Tarweed) <sup>1</sup>	Ragweed, Common <sup>1</sup>
Flixweed <sup>1</sup>	Ragweed, Giant (Buffaloweed) <sup>1</sup>
Fumitory <sup>1</sup>	Rocket, London <sup>1</sup>
Gromwell, Corn <sup>1</sup>	Rocket, Yellow <sup>1</sup>
Groundsel, Common <sup>1</sup>	Salsify (Goatsbeard) <sup>1</sup>
Hempnettle <sup>1</sup>	Shepherdspurse <sup>1</sup>
Henbit	Smartweed, Green
Jacobs Ladder <sup>1</sup>	Smartweed, Pennsylvania
Knawel (German Moss)	Sorrel, Red (Sheep Sorrel) <sup>1</sup>
Knotweed, Prostrate	Sowthistle, Annual
Kochia	Starthistle, Yellow <sup>1</sup>
Ladysthumb	Sunflower, Common (Wild)
Lambsquarters, Common	Thistle, Canada <sup>2</sup>
Lettuce, Miners <sup>1</sup>	Thistle, Russian
Lettuce, Prickly	Velvetleaf
Mallow, Common	Vetch <sup>1</sup>
Mayweed, Chamomile (Dogfennel) <sup>1</sup>	Yarrow, Common <sup>2</sup>
Pepperweed, Peppergrass <sup>1</sup>	

<sup>1</sup>These weeds will be controlled with RIFLE tank mixtures. Refer to tank mix label for specific weeds controlled.

<sup>2</sup>RIFLE tank mixes will provide suppression of established broadleaf weeds and control of their seedlings.

#### RATES AND TIMINGS

Application of RIFLE may be made before, during or after planting of small grains. For best performance, make applications when weeds are in the 2-3 leaf stage and rosettes are less than 2 inches across. Application of RIFLE to small grains during periods of rapid growth may result in crop leaning. This condition is temporary and will not reduce crop yields.

Use RIFLE at 2 to 4 fluid ounces per treated acre in wheat, fall seeded barley, and oats, and at 2 to 3 fluid ounces per treated acre in spring seeded barley. Use the higher level of listed rate ranges when treating difficult to control weeds such as kochia, Russian thistle and prickly lettuce or dense vegetative growth.

RIFLE used in a tank mix with other herbicides offers the best spectrum of weed control and herbicide tolerant or resistant weed management. Refer to specific crop for RIFLE rate and application timing.

For applications prior to the emergence of weeds or when sulfonylurea-resistant weeds are present or suspected, use a minimum of 3 fluid ounces per treated acre of RIFLE with a tank mix herbicide. Non-sulfonylurea herbicides, such as 2,4-D or MCPA tank mixed with RIFLE will offer more consistent control of sulfonylurea resistant weeds.

When tank mixing with sulfonylurea herbicides, such as Ally®, Amber®, Express®, Finesse®, Glean® and Harmony® Extra, use an agriculturally approved surfactant of at least 80% active ingredient at the rate of 1-4 pints/100 gallons of spray or not more than 0.25-0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature and difficult to control weeds or dense vegetative growth.

**FALL AND SPRING SEEDED WHEAT**

RIFLE must be applied to fall seeded wheat prior to the jointing stage. Applications to spring seeded wheat must be made before wheat reaches the 6-leaf stage.

**TANK MIX TREATMENTS**

RIFLE may be tank mixed with one or more of the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, and geographic and other restrictions.

**BROADCAST RATE PER TREATED ACRE<sup>1</sup>**

Apply 2-4 fluid ounces RIFLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D amine or ester	2,4-D	4 lb./gal.	8-12 fl. oz. (0.25-0.375 lb. a.i./A) <sup>2</sup>
MCPA amine or ester	MCPA	4 lb./gal.	8-12 fl. oz. (0.25-0.375 lb. a.i./A) <sup>2</sup>
Ally®	metsulfuron-methyl	60% DF	1/10 oz.
Amber®	Triasulfuron	75% DF	0.28 oz.
Express®	thifensulfuron + tribenuron-methyl	75% DF	1/6 oz.
Finesse®	chlorsulfuron + metsulfuron-methyl	75% DF	1/3 oz.
Glean®	Chlorsulfuron	75% DF	1/6 oz.
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	1/3 oz.
BROX™ 2EC Herbicide, Buctril®	bromoxynil <sup>3</sup>	2 lb./gal.	1-1 1/2 pts.
BROX™-M Herbicide, Bronate®	bromoxynil + MCPA	4 lb./gal.	1-2 pts.
Curtail™	clopyralid + 2,4-D	2.38 lb./gal.	2-2 2/3 pts.
Stinger™	Clopyralid	3 lb./gal.	1/4-1/3 pt.
Karmex® <sup>4</sup>	diuron <sup>3</sup>	80% DF	1/2 - 1 1/2 lbs.
Sencor® <sup>4</sup>	metribuzin <sup>3</sup>	75% DF	1-10 oz.
Dakota® <sup>5</sup>	fenoxaprop-ethyl + MCPA	3.1 lb./gal.	16 oz.
Tiller® <sup>5</sup>	fenoxaprop-ethyl +	2.7 lb./gal.	1-1.7 pts.

	MCPA + 2,4-D	
--	--------------	--

<sup>1</sup>Early developing wheat varieties such as TAM 107, MADISON, OR WAKEFIELD must receive application between early tillering and the jointing stage. Care should be taken in staging these varieties to be certain that the application occurs prior to the jointing stage.

<sup>2</sup>When using formulations other than 4 lb./gal., use pounds active/acre listed.

<sup>3</sup>Herbicides with the same active ingredient and/or different formulations may be used.

<sup>4</sup>Tank mixtures for fall seeded wheat only.

<sup>5</sup>Use 2 fluid ounces of RIFLE only. Do not use if wild oats is the target weed. Do not use RIFLE as a tank mix treatment with Dakota<sup>®</sup> or Tiller<sup>®</sup> on Durum wheat.

SPECIAL USE TANK MIXES FOR SPRING AND FALL SEEDED WHEAT  
(SEE FOOTNOTES FOR APPLICABLE USES)

Apply 3-4<sup>1</sup> fluid ounces RIFLE with:

PRODUCT <sup>2</sup>	ACTIVE INGREDIENT	FORMULATION	AMOUNT OF PRODUCT PER ACRE
2,4-D or MCPA amine	2,4-D or MCPA	4.0 lb./gal.	1-2 pts. <sup>3</sup> (0.5-1.0 lb. a.i./A) <sup>4</sup>
2,4-D or MCPA ester	2,4-D or MCPA	4.0 lb./gal.	1-1 1/2 pts. <sup>3</sup> (0.5-0.75 lb. a.i./A) <sup>4</sup>
Ally <sup>®</sup>	metsulfuron-methyl	60% DF	1/20-1/10 oz.
Amber <sup>®</sup>	triasulfuron	75% DF	0.14-0.28 oz.
Express <sup>®</sup>	thifensulfuron + tribenuron-methyl	75% DF	1/12-1/6 oz.
Finesse <sup>®</sup>	chlorsulfuron + metsulfuron-methyl	75% DF	1/6-1/3 oz.
Glean <sup>®</sup>	chlorsulfuron	75% DF	1/6 oz.
Harmony <sup>®</sup> Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6-1/3 oz.
Ally <sup>®</sup> + 2,4-D amine or ester <sup>5</sup>	metsulfuron-methyl + 2,4-D	60% DF + 4 lb./gal.	1/20-1/10 oz. + 8 fl. oz.
Amber <sup>®</sup> + 2,4-D amine or ester <sup>5</sup>	triasulfuron + 2,4-D	75% DF + 4 lb./gal.	0.14-0.28 oz. + 8 fl. oz.
Express <sup>®</sup> + 2,4-D amine or ester <sup>5</sup>	(thifensulfuron + tribenuron-methyl) + 2,4-D	75% DF + 4 lb./gal.	1/12-1/6 oz. + 8 fl. oz.
Finesse <sup>®</sup> + 2,4-D amine or ester <sup>5</sup>	(chlorsulfuron + metsulfuron-methyl) + 2,4-D	75% DF + 4 lb./gal.	1/6-1/3 oz. + 8 fl. oz.
Glean <sup>®</sup> + 2,4-D amine or ester <sup>5</sup>	chlorsulfuron + 2,4-D	75% DF + 4 lb./gal.	1/6 oz. + 8 fl. oz.
Harmony <sup>®</sup> Extra + 2,4-D amine or ester <sup>5</sup>	(thifensulfuron + tribenuron-methyl) + 2,4-D	75% DF + 4 lb./gal.	1/6-1/3 oz. + 8 fl. oz.
Roundup <sup>®</sup> RT <sup>6</sup>	glyphosate	3.0 lb./gal.	12-16 fl. oz.

<sup>1</sup>RIFLE may be used at 6 fluid ounces on fall seeded wheat in western Oregon as a spring application only. In CO, KS, NM, OK and TX up to 8 fluid ounces of RIFLE may be applied on fall seeded wheat after it exceeds the 3-leaf stage for suppression of perennial weeds, such as field bindweed. Applications may be made in fall following a frost but before a killing freeze. RIFLE may be tank mixed with 2,4-D amine at 8 fluid ounces after wheat begins to tiller. Periods of extended stress, such as cold and wet weather, may enhance the possibility of crop injury. For fall applications only, do not use if the potential for crop injury is not acceptable.



<sup>2</sup>Do not use low rates of sulfonylurea herbicides, such as Ally<sup>®</sup>, Amber<sup>®</sup>, Express<sup>®</sup>, Finesse<sup>®</sup>, Glean<sup>®</sup>, and Harmony<sup>®</sup> Extra, on more mature weeds and/or on dense vegetative growth.

<sup>3</sup>NOTE: For use on Fall Seeded Wheat Only. Do not use unless potential crop injury will be acceptable.

<sup>4</sup>When using formulations other than 4 lb./gal., use pounds active/acre listed.

<sup>5</sup>Use for improved control of Russian thistle, flixweed, gromwell, mayweed and fiddleneck.

<sup>6</sup>RIFLE may be applied at 2 fluid ounces with Roundup<sup>®</sup> RT or any glyphosate formulation labeled for use as a preplant application to small grains with no waiting period prior to planting. Read and follow label directions of the tank mix product for adjuvant use recommendations.

**FALL SEEDED BARLEY**

RIFLE must be applied to fall seeded barley prior to the jointing stage.

NOTE: For spring seeded barley varieties that are seeded during the winter months or later, follow the rates and timings given for spring seeded barley.

**TANK MIX TREATMENTS**

RIFLE may be tank mixed with one or more of the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions.

**BROADCAST RATE PER TREATED ACRE**

Apply 2-4 fluid ounces RIFLE with:

Product <sup>1</sup>	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D amine or ester	2,4-D	4 lb./gal.	8 fl. oz. (0.25 lb. a.i./A) <sup>2</sup>
MCPA amine or ester	MCPA	4 lb./gal.	8-12 fl. oz. (0.25-0.375 lb. a.i./A)
Ally <sup>®</sup>	metsulfuron-methyl	60% DF	1/20-1/10 oz.
Amber <sup>®</sup>	triasulfuron	75% DF	0.14-0.28 oz.
Express <sup>®</sup>	thifensulfuron + tribenuron-methyl	75% DF	1/12-1/6 oz.
Finesse <sup>®</sup>	chlorsulfuron + metsulfuron-methyl	75% DF	1/6-1/3 oz.
Glean <sup>®</sup>	chlorsulfuron	75% DF	1/6 oz.
Harmony <sup>®</sup> Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6-1/3 oz.
Sencor <sup>®</sup>	metribuzin <sup>3</sup>	75% DF	1-10 oz.
BROX <sup>™</sup> 2EC Herbicide, Buctrij <sup>®</sup>	bromoxynil	2 lb./gal.	1-1 1/2 pts.
BROX <sup>™</sup> -M Herbicide, Bronate <sup>®</sup>	bromoxynil + MCPA	4 lb./gal.	3/4-1 1/2 pts.

<sup>1</sup>Do not use low rates of sulfonylureas (Ally<sup>®</sup>, Amber<sup>®</sup>, Express<sup>®</sup>, Glean<sup>®</sup>, and Harmony<sup>®</sup> Extra) on more mature weeds and/or on dense vegetative growth.

<sup>2</sup>When using formulations other than 4 lb./gal., use pounds active/acre listed.

<sup>3</sup>Herbicides with the same active ingredient and/or different formulations may be used.

**SPRING SEEDED BARLEY**

RIFLE must be applied before spring seeded barley exceeds the 4-leaf stage.

**TANK MIX TREATMENTS**

RIFLE may be tank mixed with one or more of the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions.

**BROADCAST RATE PER TREATED ACRE**

Apply 2-3 fluid ounces of RIFLE with:

PRODUCT <sup>1</sup>	ACTIVE INGREDIENT	FORMULATION	AMOUNT OF PRODUCT PER ACRE
MCPA amine or ester	MCPA	4 lb./gal.	8-12 fl. oz. (0.25-0.375 lb. a.i./A) <sup>2</sup>
Ally <sup>®</sup>	metsulfuron-methyl	60% DF	1/20-1/10 oz.
Amber <sup>®</sup>	triasulfuron	75% DF	0.14-0.28 oz.
Express <sup>®</sup>	thifensulfuron + tribenuron-methyl	75% DF	1/12-1/6 oz.
Finesse <sup>®</sup>	chlorsulfuron + metsulfuron-methyl	75% DF	1/6-1/3 oz.
Glean <sup>®</sup>	chlorsulfuron	75% DF	1/6 oz.
Harmony <sup>®</sup> Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6-1/3 oz.
Sencor <sup>®</sup>	metribuzin <sup>3</sup>	75% DF	1-10 oz.
BROX <sup>™</sup> 2EC Herbicide, Buctril <sup>®</sup>	bromoxynil	2 lb./gal.	1-1 1/2 pts.
BROX <sup>™</sup> -M Herbicide, Bronate <sup>®</sup>	bromoxynil + MCPA	4 lb./gal.	3/4-1 1/2 pts.

<sup>1</sup>Do not use low rates of sulfonyleureas (Ally<sup>®</sup>, Amber<sup>®</sup>, Express<sup>®</sup>, Glean<sup>®</sup>, and Harmony<sup>®</sup> Extra) on more mature weeds and/or on dense vegetative growth.

<sup>2</sup>When using formulations other than 4 lb./gal., use pounds active/acre listed.

<sup>3</sup>Herbicides with the same active ingredient and/or different formulations may be used.

**FALL AND SPRING SEEDED OATS**

RIFLE must be applied before spring seeded oats exceed the 5-leaf stage. Applications to fall seeded oats must be made prior to the jointing stage.

**TANK MIX TREATMENTS**

RIFLE may be tank mixed with one or more of the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions.

**BROADCAST RATE PER TREATED ACRE**

Apply 2-4 fluid ounces of RIFLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
---------	-------------------	-------------	----------------------------

MCPA amine or ester	MCPA	4 lb./gal.	8-12 fl. oz. (0.25-0.375 lb. a.i./A) <sup>1</sup>
---------------------	------	------------	---

<sup>1</sup>When using formulations other than 4 lb./gal., use pounds active/acre listed.

**SUGARCANE**

Observe all precautions. Read and follow MIXING AND APPLICATION instructions.

Consult your local or state authorities for possible application restrictions, especially concerning aerial applications and advice concerning special local use situations.

**WEEDS CONTROLLED**

RIFLE, when applied at the recommended rates, will control many annual, biennial and perennial broadleaf weeds commonly found in sugarcane. (Refer to GENERAL WEED LIST.)

**RATES AND TIMINGS**

Application of RIFLE may be made any time after weeds have emerged and are actively growing but before the close-in stage of sugarcane. Application rates and timings of RIFLE are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage & Type	Product Amount	Broadcast Rate Per Treated Acre (lbs. a.i.)
Annual		
Small, actively growing	1/2-1 pt.	1/4-1/2
Established growth	1-1 1/2 pts.	1/2-3/4
Biennial	1-2 pts.	1/2-1
Perennial	2-4 pts.	1-2*

\*Application made over the top of actively growing sugarcane may result in crop injury. When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage.

Retreatments may be made as needed; however, do not exceed a total of 4 pints (2 lbs. a.i.) of RIFLE per treated acre during a growing season.

**TANK MIX TREATMENTS**

RIFLE may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled, geographic and other restrictions.

Herbicide	Rate Per Treated Acre (lbs. a.i.)
ametryn (Evik®)	2/5-8
asulam (Asulox®)	2-3 1/3
atrazine	2/5-4
2,4-D	1/2-3*

\*Application of RIFLE plus 2,4-D tank mix at the higher listed ranges may result in crop injury.

**PASTURE, HAY, RANGELAND AND GENERAL FARMSTEAD (NON-CROPLAND)**

RIFLE is recommended for use on pasture, hay, rangeland, general farmstead (non-cropland including fence rows and non-irrigation ditchbanks) for broadleaf weed and brush control. RIFLE may also be applied to non-cropland areas for the control of broadleaf weeds in noxious weed control programs.

Districts or areas including broadcast or spot treatment of roadsides and highways, utilities, railroad and pipeline rights-of-way. Noxious weeds must be recognized at the state level but programs may be administered at state, county or other level.

Observe all precautions. Read and follow mixing and application instructions.

RIFLE uses described in this section also pertain to small grains (such as barley, forage sorghum, oats, rye, sudangrass or wheat) grown for pasture use only.

Newly seeded areas, including small grains grown for pasture may be severely injured if rates of RIFLE greater than 1 pint/A are applied.

Established grass crops growing under stress can exhibit various injury symptoms that may be more pronounced if herbicides are applied. Furthermore, rates of RIFLE in excess of 2 quarts (2 lbs. a.i.) per treated acre may cause temporary injury to many grass species.

Bentgrass, carpetgrass, buffalograss and St. Augustine grass may be injured at rates exceeding 1 pint RIFLE (1/2 lb. a.i.) per treated acre. Usually colonial bentgrasses are more tolerant than creeping types. Velvetgrasses are most easily injured. Treatments will kill or injure alfalfa, clovers, lespedeza, wild winter peas, vetch and other legumes.

Animals cannot be removed from treated area for slaughter prior to 30 days after last application. There is no waiting period between treatment and grazing for non-lactating animals.

**TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT**

RIFLE Rate per Treated Acre	Days Before Grazing	Days Before Hay Harvest
Up to 1 pint (1/2 lb. a.i.)	7 days	37 days
Up to 1 quart (1 lb. a.i.)	21 days	51 days
Up to 2 quarts (2 lbs. a.i.)	40 days	70 days

NOTE: Observe all precautions and restrictions on labels of products used in tank mixtures.

**RIFLE contains 0.5 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.**

**MIXING AND APPLICATION**

RIFLE can be applied using water, oil in water emulsions including invert systems, or sprayable fluid fertilizer as a carrier. A compatibility test (see COMPATIBILITY TEST section) should be made prior to tank mixing.

To prepare oil in water emulsions, half-fill spray tank with water. Then add the appropriate amount of emulsifier with continuous agitation. Slowly add the herbicide and then the oil (such as diesel oil or fuel oil) or a premix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers. RIFLE may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply 3 to 600 gallons of diluted spray per treated acre. Volume of spray applied will depend on the height, density, and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment, apply 2 to 40 gallons of diluted spray per treated acre in a water-based carrier.

RIFLE may be applied to individual clumps or small areas (spot treatment) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to run-off) of foliage and stems.

Herbicide adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use recommendations and precautions on product label.

**WEEDS CONTROLLED**

RIFLE, when applied at recommended rates, will give control of many annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species commonly found in pasture, hay, rangeland and general farmstead (non-cropland) areas. (Refer to GENERAL WEED LIST.) Perennial weeds noted with an asterisk (\*) may be controlled with lower rates of either RIFLE D or RIFLE plus 2,4-D. See RATES AND TIMINGS below.

**RATES AND TIMINGS**

Application rates and timing of RIFLE are given below. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

WEED STAGE & TYPE	PRODUCT AMOUNT	BROADCAST RATE PER TREATED ACRE (LBS. A.I.)
Annual Small, actively growing Established weed growth	1/2-1 pt. 1-1 1/2 pts.	1/4-1/2 1/2-3/4
Biennial <sup>1</sup> Rosette diameter Less than 3 inches 3 inches or more Bolting	1/2-1 pt. 1-2 pts. 2-3 pts.	1/4-1/2 1/2-1 1-1 1/2
Perennial Suppression or top growth control Noted (*) Perennials	1/2-1 qt. 1-2 qts.	1/2-1 1-2*
Other Perennials	2 qts.	2*
Woody Brush & Vines Top growth suppression Top growth control <sup>2</sup> Stems and stem suppression	1/2-1 qt. 1-2 qts. 2 qts.	1/2-1 1-2* 2*

<sup>1</sup>For best performance, make application when biennial weeds are in the rosette stage.  
<sup>2</sup>Species noted in GENERAL WEED LIST section will require tank mixtures for adequate control.  
 \*Rates above 2.0 lb. a.i./A are spot treatments only. Do not broadcast apply more than 1 lb. a.i./A. Retreatments may be made as needed; however, do not exceed a total of 2 quarts (2 lbs. a.i.) of RIFLE per treated acre during a growing season.

**TANK MIX TREATMENTS**

Read and follow the label of each tank mix product used for precautionary statements, directions for use, application rates and other restrictions. RIFLE may be tank mixed with one or more of the following herbicides for control of grasses, additional broadleaf weeds, and woody brush and vines.

HERBICIDE	RATE PER TREATED ACRE (LBS. A.I.)
-----------	-----------------------------------

Pasture, hay, rangeland and general farmstead (non-cropland) use:	
glyphosate (Gly Star™ Original, Roundup®)	¾-3 ¾
metsulfuron methyl (Ally®)	0.0038-0.011
paraquat (Gramoxone®)	½-1
picloram (Tordon™)	1/8-3
tricolopyr (Garlon™)	¾-9
2,4-D	¼-6

Due to variations that may occur in formulated products and specific use ingredients (e.g. water supplies), a compatibility test (see COMPATIBILITY TEST section) is recommended prior to actual tank mixing.

**CUT SURFACE TREE TREATMENTS**

RIFLE may be applied as a cut surface treatment for control of unwanted trees and prevention of sprouts of cut trees. A mix of 1 part RIFLE with 1 to 3 parts water should be used in application. Use the lower dilution when treating difficult-to-control species.

FRILL OR GIRDLE TREATMENTS: Make a continuous cut or a series of overlapping cuts using an axe to girdle tree trunk. Spray or paint cut surface with the RIFLE /water mix.

STUMP TREATMENTS: Spray or paint freshly cut surface with the water mix. The area adjacent to the bark should be thoroughly wet.

NOTE: For more rapid foliar effects, 2,4-D may be added to the RIFLE /water mix.

**DORMANT APPLICATIONS FOR CONTROL OF MULTIFLORA ROSE**

RIFLE can be applied when plants are dormant as an undiluted SPOT-CONCENTRATE directly to the soil or as a LO-OIL BASAL BARK treatment using an oil-water emulsion solution.

SPOT-CONCENTRATE applications of RIFLE should be applied directly to the soil as close as possible to the root crown but within 6-8 inches of the crown. On sloping terrain, application should be made to the uphill side of the crown. Do not make application when snow or water prevents applying RIFLE directly to the soil. The use rate of RIFLE is dependent on the canopy diameter of the multiflora rose. Examples: Use RIFLE at ¼, 1 or 2 ¼ fluid ounces of product respectively, for 5, 10, or 15 feet canopy diameters. Do not exceed a total of 2 qts. RIFLE per acre per year.

LO-OIL BASAL BARK applications of RIFLE should be applied to the basal stem regions from the ground line up to a height of 12 to 18 inches. Spray until runoff, with special emphasis on covering the root crown. For best results, make application when plants are dormant. Do not make application after bud break or when plants are showing signs of active growth. Do not make application when snow or water prevents applying RIFLE to the ground line. Refer to MIXING AND APPLICATIONS above in this section for method of preparing oil-in-water emulsion. Example for making approximately 2 gallons of a LO-OIL spray mixture: combine 1 ½ gallons water plus 1 ounce emulsifier plus 1 pint RIFLE plus 2 ½ pints of No. 2 diesel fuel. Adjust amounts of materials used proportionately to the amount of final spray solution desired. Do not exceed 8 gallons of spray solution mix applied per acre per year.

**CONSERVATION RESERVE PROGRAM (CRP) ACRES**

RIFLE is recommended for use on both newly seeded and established grasses grown in Conservation Reserve or Federal Set-Aside Programs.

Observe all precautions, MIXING AND APPLICATION directions.

RIFLE treatment will injure or may kill alfalfa, clovers, lespedeza, wild winter peas, vetch, and other legumes.

Agriculturally approved surfactants may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

Do not use adjuvants containing penetrants such as petroleum based oils after grass emergence on newly seeded grasses.

**RIFLE contains 0.5 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.**

**NEWLY SEEDED AREAS**

RIFLE may be applied either preplant or postemergence to newly seeded grasses or small grains such as barley, oats, rye, sudangrass, wheat, or other grain species grown as a cover crop. Postemergence applications may be made after seedling grasses exceed the 3-leaf stage. Rates of RIFLE greater than 1 pint per treated acre may severely injure newly seeded grasses.

Preplant applications: Injury to new seedings may occur if intervals between application and grass planting are less than 45 days per pint of RIFLE per treated acre west of the Mississippi River or 20 days per pint east of the Mississippi River.

**ESTABLISHED GRASS STANDS**

Established grass stands are perennial grasses planted one or more seasons prior to treatment. Certain species, bentgrass, carpetgrass, smooth brome, buffalograss or St. Augustine grass, may be injured when treated with RIFLE at rates exceeding 1 pint per treated acre.

**WEEDS CONTROLLED**

RIFLE, when applied at recommended rates, will control many annual and biennial weeds and provide control or suppression of many perennial weeds. (Refer to GENERAL WEED LIST.)

**RATES AND TIMINGS**

Application rates and timings of RIFLE treatments are given below. Use the higher rate of the rate range when vegetation is either dense or tall, or when weeds are growing under stressed conditions such as drought or cool temperature.

Weed Type* & Stage	Broadcast Rate Per Treated Acre	
	Amount of formulated RIFLE (pts.)	Equivalent lbs. a.i.

Annuals		
Small, actively growing	1/4-1	1/8-1/2
Established weed growth	1	1/2
Biennials**		
Rosette diameter		
Less than 3 inches	1/2-1	1/4-1/2
3 inches or greater	1-2	1/2-1
Bolting biennial	2-3	1-1 1/2
Perennials**		
Suppression/Control	2-4	1-2

\*For best results, treat biennial weeds with RIFLE when they are in the rosette stage of growth. Retreatments may be made as needed; however, do not exceed a total of 2 quarts (2 lbs. a.i.) of RIFLE per treated acre during a growing season.

\*\*biennial and perennial weeds will require follow-up (sequential) treatments for seedling control and escapes.

**TANK MIX TREATMENTS**

To control grasses and additional broadleaf weeds, RIFLE may be tank mixed with other herbicides registered for use in Conservation Reserve Programs such as 2,4-D, glyphosate (Gly Star™ Original or Roundup®), paraquat (Gramoxone®), metsulfuron (Ally®) and others.

Read and follow the label of each tank mix product used for precautionary statements, directions for use, application rates, and other restrictions.

**ASPARAGUS**

FOR USE ONLY IN THE STATES OF CALIFORNIA, OREGON AND WASHINGTON

Observe all precautions. Read and follow mixing and application instructions.

If spray contacts emerged spears, crooking (twisting) of some spears may result. If such crooking occurs, discard affected spears.

Do not harvest prior to 24 hours after treatment.

Do not use in the Coachella Valley of California.

**Two applications** may be made per growing season. Do not exceed a total of 1 pint of RIFLE per treated acre per crop year.

**RIFLE contains 0.5 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.**

**RATES AND TIMINGS**

Apply RIFLE to emerged and actively growing weeds in 40 to 60 gallons of diluted spray per treated acre immediately after cutting the field, but at least 24 hours before the next cutting.

WEEDS	RATE PER TREATED ACRE
Mustard, Black	
Pigweed, Redroot (Carelessweed)	
Sowthistle, Annual	1/2-1 pt
*Thistle, Canada	(1/4-1/2 lb. a.i.)



330441

Thistle, Russian	
*Bindweed, Field	
Chickweed, Common	
Goosefoot, Nettleleaf	1 pt.
Radish, Wild	(1/2 lb. a.i.)
Thistle, Milk	

RIFLE may be applied in a tank mixture with either 2,4-D, Gly Star™ Original or Roundup® for improved control of noted (\*) weeds. Read and follow 2,4-D, Gly Star™ Original or Roundup® product labeling for precautionary statements, directions for use, application rates and timings, and other restrictions.

**TURF AND LAWNS**

**FOR USE IN GENERAL FARMSTEAD (NON-CROPLAND) AND SOD FARMS**

Observe all precautions. Read and follow mixing and application instructions.

To avoid injury to newly seeded grasses, application of RIFLE should be delayed until after the second mowing. Furthermore, application rates in excess of 1 pint (1/2 lb. a.i.) per treated acre may cause noticeable stunting or discoloration of sensitive grass species such as bentgrass, carpetgrass, buffalograss, and St. Augustine grass.

In areas where roots of sensitive plants extend, do not apply in excess of 1/4 pint (1/8 lb. a.i.) of RIFLE per treated acre on coarse textured (sandy-type) soils, or in excess of 1/2 pint (1/4 lb. a.i.) per treated acre on fine textured (clay-type) soils. Do not make repeat applications in these areas for 30 days and until previous applications of RIFLE have been activated in the soil by rain or irrigation.

**RIFLE contains 0.5 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.**

**WEEDS CONTROLLED**

RIFLE, when applied at recommended rates, will give control of many annual, biennial, and noted (\*) perennial broadleaf weeds commonly found in turf. RIFLE will also give growth suppression of many other listed perennial broadleaf weeds and woody brush and vine-species. (Refer to GENERAL WEED LIST.)

**MIXING AND APPLICATION**

Apply 30 to 200 gallons of diluted spray per treated acre (3 qts. to 4 1/4 gals. per 1,000 sq. ft.), depending on density or height of weeds treated and on the type of equipment used.

**RATES AND TIMINGS**

Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage & Type	RIFLE		
	Pints per treated acre	Pounds a.i. per treated acre	Teaspoons per 1000 sq. ft.

Annuals Small, actively growing Established weed growth	1/2-1 1-1 1/2	1/4-1/2 1/2-3/4	1-2 1/4 2 1/4-3 1/4
Biennials, Rosette diameter Less than 3 inches 3 inches or more	1/2-1 1-2	1/4-1/2 1/2-1	1-2 1/4 2 1/4-4 1/2
Perennials, Woody Brush and Vines	1-2	1/2-1	2 1/4-4 1/2

For best performance, apply when weeds are emerged and actively growing.

Retreatments may be made as needed; do not exceed a total of 2 pints (1 lb. a.i.) RIFLE per treated acre during a growing season.

**TANK MIX TREATMENTS**

Read and follow the label of each tank mix product used for precautionary statements, directions for use, application rates and timings and other restrictions.

Tank mix treatments of RIFLE may be made with 2,4-D, MCPA, MCPP, or bromoxynil for control of additional weeds listed on the tank mix product label.

Apply 1/5 to 1/2 pint (1/10 to 1/4 lb. a.i.) of RIFLE per treated acre with 1/2 to 1 1/2 lbs. acid equivalent of 2,4-D, MCPA, MCPP, or with 3/8 to 1/2 lb. a.i. of bromoxynil. Use the higher level of the listed rate ranges when treating established weeds. Repeat treatments may be made as needed; however, do not exceed 2 pints (1 lb. a.i.) of RIFLE per treated acre during the growing season.

**GRASS SEED CROPS**

**GRASSES GROWN FOR SEED SUCH AS BERMUDAGRASS, BLUEGRASS, FESCUE AND RYEGRASS**

Observe all precautions. Read and follow mixing and application instructions.

Refer to the PASTURE, HAY, RANGELAND, AND GENERAL FARMSTEAD (NON-CROPLAND) section for possible grazing and feeding restrictions.

Do not use on bentgrass unless possible crop injury can be tolerated.

**RIFLE contains 0.5 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.**

**WEEDS CONTROLLED**

RIFLE will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds plus additional weeds, it is recommended that RIFLE be applied in a tank mix with other herbicides.

Alfalfa<sup>1</sup>  
Bedstraw, Catchweed  
Bindweed, Field  
Buttercup, Corn

Hemlock, Poison  
Knapweed, Russian<sup>1</sup>  
Knawel  
Knotweed, Prostrate

Buttercup, Creeping  
 Buttercup, Western Field  
 Catchfly, Nightflowering  
 Chamomile, Corn  
 Chickweed, Common  
 Chickweed, Mouseear  
 Clover  
 Cockle, White  
 Dock, Broadleaf  
 Dock, Curly  
<sup>1</sup>Top growth only.

Kochia  
 Ladysthumb  
 Lambsquarters, Common  
 Lettuce, Prickly  
 Mayweed (Dogfennel)  
 Ragwort, Tansy  
 Sorrel, Red (Sheep Sorrel)  
 Sowthistle, Annual  
 Starwort, Little  
 Thistle, Canada<sup>1</sup>

### RATES AND TIMINGS

Apply ½ to 1 pint of RIFLE per treated acre on seedling grass after the crop reaches the 3 to 5 leaf stage. Apply up to 2 pints of RIFLE on well-established perennial grass. Do not apply after the grass seed crop begins to joint. For best performance, make applications when weeds are in the 2 to 4 leaf stage and rosettes are less than 2 inches across. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.

### TANK MIX TREATMENTS

For control of grasses or additional broadleaf weeds, RIFLE may be tank mixed with all broadleaf herbicides registered for use in Grass Seed Production. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

### BROADCAST RATE PER TREATED ACRE

Apply ½ to 2 pints RIFLE with:

PRODUCT	ACTIVE INGREDIENT	FORMULATION	AMOUNT OF PRODUCT PER ACRE
2,4-D amine or ester	2,4-D	4 lb./gal.	1-4 pts. (0.5-2.0 lbs. a.i./A) <sup>1</sup>
MCPA amine	MCPA	4 lb./gal.	1-2 pts. (0.5-1.0 lb. a.i./A) <sup>1</sup>
BROX™ 2EC Herbicide, Buctril®	bromoxynil <sup>2</sup>	2 lb./gal.	1-2 pts.
Curtail™	clopyralid + 2,4-D	2.38 lb./gal.	1 ¾-4 pts.
Karmex®	diuron <sup>2</sup>	80% DF	2-4 lbs.
Stinger™	clopyralid	3 lb./gal.	¼-1 pt.

<sup>1</sup>When using formulations other than 4 lb./gal., use pounds active/acre listed.

<sup>2</sup>Herbicides with the same common name and/or different formulations may be used.

### ANNUAL GRASS CONTROL

For suppression of annual grass weeds such as:

Brome, Downy (Cheatgrass)  
 Brome, Rippgut  
 Fescue, Rattail  
 Windgrass

Apply up to 4 pints of RIFLE per treated acre in the fall or late summer after harvest and burning of established grass seed crops. Applications should be made immediately following the first irrigation when the soil is moist and before weeds have more than 2 leaves.

**PREPLANT DIRECTIONS (POST HARVEST/FALLOW/CROP STUBBLE/SET-A-SIDE) FOR  
BROADLEAF WEED CONTROL BEFORE WHEAT, CORN, SORGHUM, SOYBEANS**

Observe all precautions. Read and follow mixing and application instructions.

**WEEDS CONTROLLED**

RIFLE may be applied alone or in tank mix combinations with other herbicides registered for this use.

RIFLE can be applied either post harvest in the fall, spring or summer, during the fallow period or to crop stubble/set-a-side acres. RIFLE, when applied at the recommended rates, will control many annual broadleaf weeds. See the WEEDS CONTROLLED section under small grains. In addition, RIFLE will control or suppress the following biennial and perennial broadleaf weeds:

Alfalfa <sup>1</sup>	Knapweed, Diffuse
Artichoke, Jerusalem	Nightshade, Silver
Bindweed, Field	Redvine
Bindweed, Hedge	Smartweed, Swamp
Blueweed, Texas	Sowthistle, Perennial <sup>1</sup>
Bursage (Bur Ragweed, Povertyweed, Lakeweed) <sup>1</sup>	Spurge, Leafy
Dandelion, Common <sup>1</sup>	Thistle, Bull
Dock, Curly <sup>1</sup>	Thistle, Canada <sup>2</sup>
Dogbane, Hemp	Thistle, Milk
Garlic, Wild <sup>2</sup>	Thistle, Musk
Horsenettle, Carolina	Thistle, Plumeless
Knapweed, Spotted	Thistle, Scotch
	Trumpetcreeper (Buckvine)

<sup>1</sup>Perennials may be controlled using RIFLE at rates lower than those recommended for other listed perennial weeds. (See RATES AND TIMINGS under this heading).

<sup>2</sup>See the SPECIAL TANK MIX TREATMENTS section under this heading for specific control programs for these weeds.

**RATES AND TIMINGS**

Apply RIFLE as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (post harvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer. Agriculturally approved spray additives, such as surfactants or oils, may be used to enhance spray coverage and the herbicide's penetration of weed foliage. See CROPPING RESTRICTIONS for recommended interval between application and planting to prevent crop injury.

For best performance, make application when annual weeds are less than 6 inches tall, when biennial weeds are in the rosette stage, and to perennial weed regrowth in late summer or fall following a mowing or tillage treatment. Most effective control of upright perennial broadleaf weeds, such as Canada thistle and Jerusalem artichoke, occurs if application is made when the majority of weeds, such as field bindweed and hedge bindweed, are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds which develop from seed or underground plant parts, such as rhizomes or bulblets, after the effective period for RIFLE. For seedling control, a follow-up program or other cultural practices could be instituted. For small grain in-crop uses of RIFLE, see the RATES AND TIMINGS section under the SMALL GRAINS heading for details.

**RIFLE RATES PER TREATED ACRE**

WEED TYPE	AMOUNT OF PRODUCT PER ACRE
-----------	----------------------------

WEED TYPE	AMOUNT OF PRODUCT PER ACRE
Annual	1/2-1 pt. (8 to 16 fl. oz.)
Biennial	1-2 pts. (16 to 32 fl. oz.)
Perennial	1-4 pts. (16 to 64 fl. oz.)
Perennial suppression	1-2 pts. (16 to 32 fl. oz.)
Noted <sup>1</sup> perennials	2-4 pts. (32 to 64 fl. oz.)
Other perennials	4 pts. (64 fl. oz.)

Retreatments may be made as needed; however, do not exceed a total of 4 pints of RIFLE per treated acre during any given period.

### TANK MIX TREATMENTS

RIFLE may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled, geographic and other restrictions.

### RIFLE BROADCAST RATE PER TREATED ACRE FOR ANNUAL WEED CONTROL

Apply 1/4 to 1 pint RIFLE with:

PRODUCT	ACTIVE INGREDIENT	FORMULATION	AMOUNT OF PRODUCT PER ACRE
Aatrex <sup>®</sup> 4L <sup>1</sup>	atrazine	4 lb./gal.	0.5-6 pts.
Aatrex <sup>®</sup> Nine-O <sup>®1</sup>	atrazine	90% DF	0.5-3 1/3 lb.
Ally <sup>®2</sup>	metsulfuron-methyl	75% DF	0.1 oz.
Amber <sup>®2</sup>	triasulfuron	75% DF	0.28-0.35 oz.
Cyclone <sup>®</sup>	paraquat	2 lb./gal.	1-2 pts.
Fallow Star <sup>™</sup> or Fallow Master <sup>™</sup>	glyphosate + dicamba	1.6 lb./gal.	22-44 fl. oz.
Finesse <sup>®2</sup>	chlorsulfuron + metsulfuron-methyl	75% DF	0.2 oz.
Gramoxone <sup>®</sup> Extra	paraquat	2.5 lb./gal.	1.5 pts.
Kerb <sup>®1</sup>	pronamide	50-W	0.5-1 lb.
Landmaster <sup>®</sup> BW	glyphosate + 2,4-D	2.4 lb./gal.	27-54 fl. oz.
Gly Star <sup>™</sup> Original, Roundup <sup>®</sup> or Roundup <sup>®</sup> RT	glyphosate	3 lb./gal.	8-48 fl. oz.
Sencor <sup>®</sup> DF <sup>1</sup>	metribuzin	75% DF	0.5-1 lb.
Sencor <sup>®</sup> 4 <sup>1</sup>	metribuzin	4 lb./gal.	0.75-1.5 pts.
2,4-D	2,4-D	4 lb./gal.	1-2 pts. (0.5-1 lb. a.i./A) <sup>3</sup>

<sup>1</sup>Tank mixes of RIFLE with these products may be subject to special restrictions. See the product label of the tank mix partner for intended use rates, restrictions and other precautions.

<sup>2</sup>When tank mixing with sulfonylurea herbicides, refer to the product label for rates and restrictions. Use a surfactant of at least 80% active ingredient at the rate of 1-2 quarts/100 gallons of spray or not more than 0.25-0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature weeds or dense vegetative growth. Sulfonylurea-resistant weeds may not be controlled by tank mixes of RIFLE and a sulfonylurea. Refer to the RIFLE tank mix section for alternative tank mixes.

<sup>3</sup>When using formulations other than 4 lb./gal., use pounds active/acre listed.

**RIFLE BROADCAST RATE PER TREATED ACRE FOR  
BIENNIAL AND PERENNIAL WEED CONTROL**

Apply 1 to 4 pints of RIFLE with:

PRODUCT	ACTIVE INGREDIENT	FORMULATION	AMOUNT OF PRODUCT PER ACRE
Curtail™	clopyralid + 2,4-D	2.38 lb./gal.	2-4 pts.
2,4-D	2,4-D	4 lb./gal.	2-6 pts. (1.0-3 lbs. a.i./A) <sup>1</sup>
Landmaster® BW	glyphosate + 2,4-D	2.4 lb./gal.	54 fl. oz.
Gly Star™ Original, Roundup®	glyphosate	3 lb./gal.	1-5 qts.
Roundup® RT	glyphosate	3 lb./gal.	1-5 qts.
Tordon™ 22K	picloram	2 lb./gal.	½-1 pt.

<sup>1</sup>When using formulations other than 4 lb./gal., use pounds active/acre listed.

**SPECIAL TANK MIX TREATMENTS**

For suppression of perennial weeds, apply ½-1 pint of RIFLE with 8-16 fluid ounces of Gly Star™ Original, Roundup® or Roundup® RT per treated acre.

For wild garlic control, apply 1 pint RIFLE with 3 pints of 2,4-D LV ester (4 lb./gal.) per treated acre. Apply when wild garlic is 4 to 8 inches tall.

For Canada thistle control, use RIFLE, RIFLE plus Curtail™, or RIFLE plus Gly Star™ Original, Roundup® or Roundup® RT tank mix treatments.

Application may be made during fallow periods for control of volunteer barley, bulbous bluegrass, downy brome, jointed goatgrass, common rye and volunteer wheat when they are actively growing. Use 1 pint RIFLE with ½-1 lb. Kerb® 50-W. Fall seeded wheat may be planted 9 months or more after application. For best performance, make application between mid-October and mid-December, prior to soil freeze up.

During fallow periods, apply RIFLE plus Landmaster® BW, Fallow Star™ or Fallow Master™ to give improved control of kochia, wild buckwheat, prickly lettuce, field bindweed and Canada thistle. Use 1/8-1/4 pint of RIFLE plus 22-54 fluid ounces of Landmaster® BW, Fallow Star™ or Fallow Master™ for annual weed control or ¼-1/2 pint RIFLE plus 22-54 fluid ounces of Landmaster® BW, Fallow Star™ or Fallow Master™ for perennial weed suppression.

**CROPPING RESTRICTIONS**

The following recommendations are based on RIFLE use rates up to 4 pints per treated acre.

Corn, sorghum, and soybeans may be planted in the spring following applications made during the previous year. If less than 1 inch of rainfall occurs between application and first killing frost, treated areas should be cultivated to allow herbicide to come in contact with moist soil. Cultivation may take place before or immediately after ground thaw.

Soybean injury may occur if the interval between application and planting is less than specified. In areas with greater than 30 inches of rainfall, delay planting for 30 days per pint of RIFLE per treated acre. In areas with less than 30 inches of rainfall, delay planting for 45 days per pint of RIFLE per treated acre. Exclude days when ground is frozen.

Wheat may be planted in the fall or spring following applications. Also, spot application may be made any time prior to crop emergence if crop injury can be tolerated in treated areas. Wheat injury may occur if the interval between application and planting is less than specified.

East of the Mississippi River, the interval is 20 days per pint of RIFLE per treated acre or 1.25 days per 1 ounce. Moisture is essential for RIFLE degradation. Exclude days when ground is frozen.

West of the Mississippi River, the interval is 45 days per pint of RIFLE per treated acre or 3 days per ounce. Moisture is essential for RIFLE degradation. Exclude days when ground is frozen.

Following a normal harvest of barley, oats, or wheat, any rotational crop may be planted. If the interval before harvest is shortened, such as when cover crops will be plowed under, do not follow up with the planting of a sensitive crop.

### **COTTON PREPLANT APPLICATION**

Observe all precautions. Read and follow mixing and application instructions. Refer to the GENERAL WEED LIST section of this label for a list of weeds controlled or suppressed.

**RIFLE contains 0.5 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.**

### **RATES AND TIMINGS**

Apply RIFLE as a broadcast or spot treatment to emerged and actively growing weeds at a rate of up to 8 fl. oz./acre prior to planting cotton. Most effective control of weeds occurs if application is made when weeds are in the 2-4 leaf stage and rosettes are less than 2" across.

### **CROPPING RESTRICTIONS**

Do not plant cotton for at least 21 days after application and after allowing for a minimum accumulation of 1" of rainfall or overhead irrigation. Do not apply west of the Rockies or to geographic areas with average annual rainfall less than 25".

### **TANK MIX TREATMENTS**

For control of grasses or additional broadleaf weeds, RIFLE may be tank mixed with Caparol<sup>®</sup>, Gramoxone<sup>®</sup> Extra, and Roundup Ultra<sup>®</sup> RT herbicides.

### **CONTROL OF PERENNIAL BROADLEAF WEEDS IN CROPLAND (SPOT APPLICATION ONLY)**

FOR USE ONLY IN THE STATES OF IDAHO, MONTANA, NEVADA, OREGON, UTAH, AND WASHINGTON

Observe all precautions. Read and follow mixing and application instructions.

Do not treat subirrigated cropland or areas where the soil remains saturated with water throughout the year.

Make only one application of RIFLE per year.

**RIFLE contains 0.5 pounds a.e. of dicamba per pint. When tank mixing with products that contain dicamba, do not exceed a combined total of 1.0 pound of a.e. per acre per application.**

**WEEDS CONTROLLED**

RIFLE, when applied at recommended rates, will control many broadleaf weeds including:

- |                              |                   |
|------------------------------|-------------------|
| Bindweed, Field              | Knapweed, Russian |
| Dock, Broadleaf (Bitterdock) | Ragwort, Tansy    |
| Dock, Curly                  | Spurge, Leafy     |
| Knapweed, Black              | Thistle, Canada   |

**RATES AND TIMINGS**

RIFLE may be applied at any time following a crop harvest to stubble, fallow or other cropland. Application should be made when weeds are actively growing and prior to a killing frost.

Apply 2 quarts (2 lbs. a.i.) of RIFLE per treated acre. Application may be made up to one month prior to the planting of wheat.

NOTE: Do not use unless injury to wheat or rotated barley will be acceptable.

Barley, oats, corn, sorghum (milo), annual or perennial grass crops may be planted into treated areas one year after application. Crops grown for seed (other than perennial grass seed) should not be planted into treated areas until three years after application. Do not plant broadleaf crops such as alfalfa, beans, peas, potatoes, or sugar beets into treated areas until two years after application.

In most cases, treatments will not kill perennial weed seedlings which germinate from seed one or two years after treatment. Once the effect of the chemical has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

**WIPER APPLICATION USES**

Important. Observe all precautions. RIFLE may be applied through wiper application equipment to control or suppress actively growing broadleaf weeds, brush and vines. Use a solution containing 1 part RIFLE to 1 part water. Do not contact desirable vegetation with herbicide solution. Wiper application should only be made to crops (including PASTURES) and NON-CROPLAND AREAS described in this label with the exception of GRAIN SORGHUM (MILO).

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**BEFORE BUYING OR USING THIS PRODUCT**, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO



OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

### REGISTERED TRADEMARKS

BROX™, Gly Star™ and Dicambazine™ are trademarks of Albaugh, Inc.

Clarity®, Frontier®, Guardsman® and Prowl® are registered trademarks of BASF Ag Products.

Asulox®, Bronate®, Buctril®, Dakota®, Sencor® and Tiller® are registered trademarks of Bayer CropScience.

Broadstrike™, Curtail™, Garlon™, Stinger™, Surpass™ and Tordon™ are trademarks of Dow AgroSciences LLC.

Accent®, Ally®, Express®, Finesse®, Glean®, Harmony® and Karmex® are registered trademarks of E.I. duPont de Nemours & Co., Inc.

Battalion®, Bullet®, Harness®, Landmaster®, Lariat®, Lasso®, Partner®, Permit®, Ramrod®, Roundup®, and Screen® are registered trademarks of the Monsanto Company.

Kerb® is a registered trademark of Rohm and Haas Company.

Aatrex®, Aatrex® Nine-O®, Amber®, Beacon®, Concep®, Cyclone®, Eradicane®, Evik®, Gramoxone®, Princep®, Sutan +® and Tough® are registered trademarks of Syngenta Crop Protection, Inc.

Rifle® is a registered Trademark of Loveland Products, Inc.