12/04/2004

s acket



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

NOTICE OF PESTICIDE:

Registration
X Reregistration
(under FIFRA, as amended)

EPA Reg. Number:	Date of Issuance:
34704-872	
,	DEC - 4 2009

Term of Issuance:

Name of Pesticide Product:

Ginmaster Cotton Defoliant

Name and Address of Registrant (include ZIP Code):

Loveland Products Inc.

P.O. Box 1286

Greeley, CO 80632-1286

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA provided that you:

1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.

Signature of Approving Official:	Date:
/s/	DEC - 4 2009
Joanne Miller	·
Senior Registration Specialist	
Herbicide Branch, Team 23	
Registration Division (7505P)	

Page 2 EPA Reg. 34704-872

- 2. On page 2, change the citrus spray drift text to "Thidiazuron products can not be applied by air within ½ mile of lettuce or within 5 miles upwind of citrus in flush (burst of new growth, as in springtime) in the Rio Grande valley of Texas. Ground applications are restricted to no less than 100 feet away from lettuce and no less than ½ mile upwind from citrus. (In general, a decrease in droplet size or increase in wind speed at the time of application will result in risk to non-target organisms. Alternatively, if droplet size is coarser or wind speeds are lower, exposures due to drift would be reduced)."
- 3. Delete the text "Do not allow persons or pets in treated areas until dusts have settled or sprays have dried" from the Hazards to Humans and Domestic Animals section of the label.
- 4. Per the acute toxicity review, the third paragraph of the handler PPE section must be revised to read:

"Mixers, loaders, and other applicators and other handlers must wear:

Coveralls over short-sleeved shirt and short pants,

Chemical-resistant shoes plus socks,

Goggles or faceshield,

Chemical-resistant gloves,

Chemical-resistant headgear for overhead exposure,

Chemical-resistant apron when mixing, loading, or cleaning equipment or spills,

A NIOSH approved particulate filtering respirator equipped with N, R, or P class filter media. The respirator should have a NIOSH approval number prefix TC-21-C. It is recommended that you require that the respirator wearer be fit tested, and trained in the use, maintenance, and limitations of the respirator.

See engineering controls for additional requirements."

- 5. The text "or rinsate" must be added to the end of the statement "Do not contaminate water by cleaning of equipment or disposal of equipment washwaters..." in the Environmental Hazard section of the label.
- 6. Per the Diuron RED, add a 21-day retreatment interval.
- 7. Delete "General" from "General Information".
- 8. On page 2, change "the spray boom should be mounted" to "the spray boom must be mounted".
- 9. On page 2, delete" recommended" from "timing of application is recommended". Change "recommendations" to "directions".

Page 3 EPA Reg. 34704-872

- 10. On page 3, delete "general" from "general use rate".
- 11. Clarify the heading "to achieve an application rate of" and the amounts as it looks as the numbers apply to diuron.

A stamped copy of the label is enclosed for your records. You must submit one copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this letter or the next printing of the label whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of the product constitutes acceptance of these conditions. This label supersedes all other previously accepted labels. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.





GINMASTER COTTON DEFOLIANT

ACCEPTED with COMMENTS In EPA Letter Dated:

DEC - 4 2009 Under the Federal Insecticide. Fungicide, and Rodenticide Acı as amended, for the pesticide registered under EPA Reg. No.

34704-872

For Agricultural Use Only

•	TOTAL	100%
INERT INGREDIENTS:		. 82%
Diuron: 3-(3,4-dichlorophenyl)-1,1-dimethylurea*		
Thidiazuron: N-phenyl-N'-1,2,3-thidiazol-5-ylurea*		
ACTIVE INGREDIENTS:		

Contains 1 lb. Thidiazuron per gallon and 0.5 lb. Diuron per gallon. *Protected by U.S. Patent Nos. 4,294,605 and 4,613,354.

KEEP OUT OF REACH OF CHILDREN DANGER — PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, have someone explain it to you in detail.)

EPA REG. NO. 34704-872

EPA EST. NO. 34704-MS-1

NET CONTENTS 21/2 GALS. (9.46 L)

EXP 01/09

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Causes irreversible eye damage. Causes skin irritation. Harmful if inhaled or absorbed through skin. Avoid contact with eyes, skin, clothing, and avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Do not allow persons or pets in treated areas until dusts have settled or sprays have dried.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are made of barrier laminate or butyl rubber. If you want more options, follow the instructions for category B on an EPA chemical-resistance selection chart.

All pilots and flaggers must wear: • Long-sleeved shirt and long pants,

- · Shoes and socks.

All mixers, loaders, other applicators, and other handlers must wear:

- · Long-sleeved shirt and long pants,
- · Shoes plus socks, and
- · Chemical-resistant gloves
- · A NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21-C or any N2, R, P, or HE filter.
- Goggles, face shield or glasses.
- Chemical-resistant apron when mixing, loading or cleaning equipment or spills, and chemical-resistant headgear for overhead exposure.

In addition to the PPE above, all handlers except for ground boom applicators, pilots, and flaggers must also wear

· A NIOSH approved particulate filtering respirator equipped with N, R, or P class filter media. The respirator should have a NIOSH approval number prefix TC-21-C. It is recommended that you require that the respirator wearer be fit tested, and trained in the use, maintenance, and limitations of the respirator.

See Engineering Controls for additional requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

Flaggers supporting aerial applications must use an enclosed cab that meets the definition in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(5)] for dermal protection. In addition, flaggers must wear:

- · Long-sleeved shirt, long pants,
- Shoes and socks.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling before eating,
- drinking, chewing gum, using tobacco, or using the toilet.

 Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

FIRST AID

If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contact lenses, if present, after the first five minutes,
	then continue rinsing.
	Call a poison control center or doctor for treatment advice.
If on skin	Take off contaminated clothing.
or clothing:	 Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
If inhaled:	Move person to fresh air.
	If person is not breathing, call 911 or an ambulance, then give
	artificial respiration, preferably mouth-to-mouth if possible.
	Call a poison control center or doctor for further treatment
L	advice.
If swallowed:	Immediately call a poison control center or doctor for treatment advice.
	Do not induce vomiting unless told to do so by a poison control center or doctor.
	Have person sip a glass of water if able to swallow.
	Do not give anything by mouth to an unconscious person.
	,

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:

1-866-944-8565.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

Note To Physician: Probable mucosal damage may contraindicate the use of gastric

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Do not apply when weather conditions favor drift from the target area.

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to product runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

This chemical has properties and characteristics associated with chemicals detected in ground water. Use of this chemical in areas where soils are permedicile, particularly where the water table is shallow, may result in ground grater contamination.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. O O In case of spillage, cover with an absorbent such as soda ash? I.me, clay, or sawdust. Sweep up and bury. Wash area thoroughly with dutorgent and Water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manne? Inconsistent with its labeling.

GINMASTER COTTON DEFOLIANT EPA REG. NO. 34704-872

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protections of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralis,
- · Shoes plus socks,
- · Chemical-resistant gloves made of any waterproof material,
- Protective evewear.

GENERAL INFORMATION

This product is a defoliant to be used as a pre-harvest aid for cotton. This product has performed well under both cool and warm weather conditions.

Use Precautions

Do not apply this product through any type of irrigation system.

Rainfall within 12 hours after application can reduce the effectiveness of this product.

Do not feed foliage from treated cotton plants or gin trash to livestock.

Do not plant the following crops earlier than the specified periods after application of this product:

(except carrots, onions) two (2) months
Legumes (including alfalfa) or leafy vegetables
(except lettuce) two (2) months
Cole crops, garlic, safflower, tomatoes, and
watermelon two (2) months
Carrots three (3) months
Onions four (4) months
Cantaloupe, honeydew melon/casaba melon,
muskmelon, or peppers five (5) months
Lettuce two (2) months
with deep-plowing of soil (12-15 inches);
or nine (9) months when soil is only disced (4-6 inches)

Cover Crops: Small grains and/or legumes intercropped with the cotton crop to which this product will be applied may only be used as cover crops and may not be harvested for food or feed. Small grains and/or legumes planted earlier than two (2) months following application of this product may only be used as cover crops and may not be harvested for food or feed.

Do not plant any other rotational crops (except those specified above) within one year of application of this product.

Do not use immature crops for food or feed.

Do not allow spray drift to contact trees or crops other than the target crop of mature cotton, or cotton you desire to defoliate, as this product may injure or defoliate other crops.

Particular care should be taken when applying this product adjacent to lettuce, citrus, or cantaloupe.

PRACTICES TO LOWER THE POTENTIAL FOR DRIFT ONTO NON-TARGET CROPS

- During applications, particularly under windy conditions, this product may drift to nontarget crops. To help reduce the drift potential, use the following practices:
- Do not apply this product by ground or air when wind speeds exceed ten (10) miles per hour at the time of application. Follow local recommendations if wind speeds of less than ten (10) miles per hour are specified in those recommendations.
- Use of low nozzle pressure (20-30 psi) is recommended.
- Do not apply this product when a temperature inversion is present or when conditions favor an inversion prior to completing applications(s).

PRACTICES TO LOWER DRIFT POTENTIAL ONTO NON-TARGET CROPS IN THE RIO GRANDE VALLEY OF TEXAS

- To help reduce the drift potential, use the following practices:
- Do not apply this product by ground or air when wind speeds exceed ten (10) miles per hour at the time of application. Follow local recommendations if wind speeds of less than ten (10) miles per hour are specified in those recommendations.

- Use of low nozzle pressure (20-30 psi) is recommended.
- Do not apply this product when a temperature inversion is present or when conditions favor an inversion prior to completing applications(s).
- Use the largest nozzle orifice possible, which permits proper deposition and coverage of product.
- Thidiazuron products can not be applied by air within ½ mile of lettuce or 5 miles downwind of the point of application when citrus is in flush (burst of new growth, as in springtime) in the Rio Grande valley of Texas. Ground applications are restricted to 100 feet away from lettuce and ½ mile downwind from the point of application for citrus. (In general, a decrease in droplet size or increase in wind speed at the time of application will result in risk to non-target organisms. Alternatively, if droplet size is coarser or wind speeds are lower, exposures due to drift would be reduced).

Requirements for reducing spray drift for ground and aerial applications:

Use best practices to avoid drift to all other crops and not-target areas. Do not apply when conditions favor drift from the target areas. The interaction of many equipment-and weather-related factors determine the potential for spray drift. Avoiding spray drift at the application site is the responsibility of the applicator. The applicator must follow the most restrictive precautions to avoid drift, including those found in this labeling as well as applicable state and local regulations and ordinances. A drift control agent may reduce drift, however, it may also decrease weed

Make aerial or ground applications only when the wind speed is less than or equal to 10 miles per hour.

Do not make aerial or ground applications into temperature inversions.

Apply with medium or coarser spray (according to ASAE standard 572) for standard nozzles.

Additional requirements for ground applications:

When applying to crops, apply with nozzle height no more than 2 feet above the ground or crop canopy. When applying to non-crop areas, use lowest nozzle height consistent with safety and efficacy. Direct spray into target vegetation.

Additional requirements for aerial applications:

The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter

Use upwind swath displacement.

When applying to crops, do not release spray at a height greater than 6 to 10 feet above the ground or crop canopy. When applying to non-crop areas, apply at a minimum safe altitude above the area being treated.

Do not apply by air if sensitive non-target crops are within 100 feet of the application site.

Time Of Application

Apply this product only to mature cotton plants when the last boll you expect to harvest is mature. A boll can be described as "mature" when it is to hard to be dented when squeezed between thumb and fingers, is difficult to slice with a sharp knife, and/or when seeds cut in cross sections have fully developed cotyledons, as evidenced by an absence of jelly within the seed.

Apply this product at least 5 days prior to anticipated harvest.

Precaution: The addition of adjuvants can cause desiccation and/or leaf freezing during periods of high temperature. The use of compounds that desiccate leaf tissue is not recommended.

Use of this product under extremely cool or adverse conditions can result in less than desirable defoliation and/or growth inhibition.

Application

This product may be applied by air or ground equipment. Apply specified dosages in 10-25 gallons of spray per acre with ground equipment or 2-10 gallons per acre by aircraft.

Dosage

Apply this product at a rate of 0.4 to 1.0 pint of formulated product per acre prior to harvest (see Application Rate Table). At some locations, following the initial application of product, it may be necessary to make a second application of this product (Do Not exceed 1.0 pint/acre per year), or an application of a standard defoliant. Apply a maximum of two applications per year. Minimum retreatment interval is 21 days.

Mixing Instructions

Fill the spray tank with one-half of the required amount of water. Add the proper amount of this product and start the spray tank agitator. Finish filling the tank with the balance of water needed. Maintain sufficient agitation during toth mixing and apparation to ensure uniformity of the spray mixture.

PRECONDITIONING FOR USE IN CALIFORNIA

This product may be used as a pre-conditioner of enhance the activity of a defoliant application. Apply this product at 4-6 oz. formulative product per acre by ground or 2-10 gallons of water per acre by aerial application. Timing of application is recommended 7-10-developing of value per acre by aerial application. Timing of application is recommended 7-10-developing to a defoliation application of this product (see Application Rate Table) or the use of another defoliant, Refer to the second product label prior to use for complete recommendations.

ر ر د د د د د

953

GINMASTER COTT DEFOLIANT EPA REG. NO. 34704-872

GINMASTER COTTON DEFOLIANT

To Achieve an Application Rate of:	Use This Amount of Ginmaster Cotton Defoliant	At the Indicated Rate, One Gallon of Ginmaster Cotton Defoliant Will Treat:
0.05 lbs. ai/Acre	0.4 pts./A (6.4 oz./A)	20 Acres
0.07 lbs. ai/Acre	0.55 pts./A (8.8 oz./A)	15 Acres
0.1 lbs. ai/Acre	0.8 pts./A (12.8 oz./A)	10 Acres
0.125 lbs. ai/Acre	1.0 pts./A (16.0 oz./A)	8 Acres

General Use Rate Information (under normal conditions) for Cotton:

Do not exceed the single use rate of 1.0 pts product

(0.125 lbs Thidiazuron ai)/A.

Do not exceed the maximum application rate of 2.4 pts product

(0.3 lbs Thidiazuron ai)/A/year

Do not exceed a maximum of 2 applications and do not exceed 2.4 pts product (0.3 lbs Thidiazuron ai)/A/vear.

General Use Rate Information (under variable conditions) for Cotton:

Do not exceed the maximum single application rate of 1.6 pts product (0.2 lbs Thidiazuron ai)/A.

Do not exceed the maximum total application rate of 2.4 pts product (0.3 lbs Thidiazuron ai)/A/year.

Do not exceed the maximum of 2 applications and do not exceed 2.4 pts product (0.3 lbs Thidiazuron ai)/A/year.

Cleanout Instructions

Do not allow the spray solution to dry in the application equipment. Immediately following application, clean all equipment (mix tanks, pumps, transfer lines, application tanks, sumps, booms, nozzles, and all related equipment) thoroughly with cleaner and water.

Should small quantities of this product remain in inadequately cleaned equipment, they may be released during subsequent applications and may cause damage to crops. Loveland Products, Inc. accepts no liability for damage to crops due to inadequately cleaned equipment.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

PESTICIDE STORAGE: Store in original container and keep closed. Store in a cool, dry place. Do not use or store near heat or open flame. Do not contaminate food or

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org. If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. (For packages up to 5 gallons:) Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(For packages greater than 5 gallons or 50 lbs:) Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/2 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(For square bottom caged totes greater than 55 gals.): Triple rinse or pressure rinse as follows: Empty the remaining contents into application equipment or a mix Storage & Disposal cor

tank and continue to drain for 10 seconds after the flow begins to drip. Fill the container about 1/4 full with water, rinsing down all sides inside the container thoroughby Recirculate water with the pump for 2 minutes. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

(For refillable containers:) Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.

Italicized type is for container size identification.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVE-LAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS PRODUCT IS SOLD AS IS TO THE EXTENT ALLOWED BY APPLICABLE LAW. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PUR-POSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND, TO THE EXTENT REQUIRED BY APPLICABLE LAW, WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, 7251 WEST 4TH STREET, GREELEY, CO 80634.

TO THE EXTENT ALLOWED BY APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT, TO THE EXTENT ALLOWED BY APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAM-AGES IN THE NATURE OF A PENALTY

