

34704-875

6/27/2012

1/10



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Premjit Halarnkar PhD MPA
Loveland Products Inc
P O Box 1286
Greeley CO 80632 1286

JUN 27 2012

Dear Dr Halarnkar

Subject Amendment Correct application rates for cotton and pecans per RED
Holster Insecticide
EPA Reg No 34704 875
Your submission dated May 7 2012

The amendment referred to above submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act (FIFRA) section 3(c)(7)(a) is acceptable and a stamped copy of the label is enclosed for your records

Please be reminded that 40 CFR Part 156 140(a)(4) requires that a batch code lot number or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers The code may appear either on the label (and can be added by non notification/PR Notice 98 10) or durably marked on the container itself

If you have any questions regarding this letter please contact Dana Pilitt PhD of my staff at (703) 305 7071 or via e mail at pilitt.dana@epa.gov

Sincerely

A handwritten signature in black ink, appearing to read "Richard Gebken".

Richard Gebken
Product Manager 10
Insecticide Branch
Registration Division (7505P)

RESTRICTED USE PESTICIDE

Due to Toxicity to fish or aquatic invertebrates

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification



HOLSTER™ INSECTICIDE

ACCEPTED

JUN 27 2012

Under the Federal Insecticide
Fungicide and Rodenticide Act
as amended for the pesticide

EPA Reg. No. 34704-875

ACTIVE INGREDIENT

Cypermethrin *

30.6%

OTHER INGREDIENTS *

69.4%

100.0%

(±) a cyano (3 phenoxyphenyl)methyl (±) *cis/trans* 3 (2,2-dichloroethenyl) 2,2-dimethylcyclopropane carboxylate

Cis/trans ratio Max 55% (±) *cis* and min 45% (±) *trans*

Contains xylene range aromatic solvents

Contains 2.5 pounds cypermethrin per gallon

Contains petroleum distillate

KEEP OUT OF REACH OF CHILDREN WARNING—AVISO

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle (If you not understand this label find someone to explain it to you in detail)

FIRST AID

If inhaled	Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible Call a poison control center or doctor for further treatment advice Note to Physician Contains petroleum distillate Vomiting may cause aspiration pneumonia
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes Call a poison control center or doctor for treatment advice
If in eyes	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
If swallowed	Immediately call a poison control center or doctor Do not give any liquid to the person Do not induce vomiting unless told to do so by the poison control center or doctor Do not give anything to an unconscious person

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL 1 866 944 8565

Note to Physician Like the natural pyrethrins the synthetic derivative is expected to have relatively minor toxicity in humans in fact any significant acute toxic effects are more likely from a carrier hydrocarbon solvent. Consequently induction of vomiting may increase the likelihood of the most important toxic potential chemical pneumonia and so should either be avoided or done only under medical supervision. Ingestion of a large amount calls for gastric lavage with care (Trendelenburg position suction available cuffed endotracheal tube if patient is unconscious) to avoid intrapulmonary aspiration. A saline cathartic (sodium or magnesium sulfate) 15 to 30 gm dissolved in water should be given as should 15 to 20 gm activated charcoal as a slurry in water. Digestible fats oils or alcohol may increase absorption and so should be avoided. Skin contact (vapor or powder) may be followed by transient tingling or numbness usually of the face but this subsides without treatment.

See below for additional precautionary information

EPA REG NO 34704 875
EPA EST NO 279 FL 001
NET CONTENTS 1 GAL (3.78 L)

EXP 05B12

3/10

**HOLSTER™ INSECTICIDE
EPA REG NO 34704 875**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING Causes moderate eye irritation Do not get on skin or clothing Harmful if swallowed Harmful if absorbed through skin Avoid contact with eyes Harmful if inhaled Avoid breathing vapor or spray mist Causes moderate eye irritation User should wash hands before eating drinking chewing gum using tobacco or using the toilet Remove contaminated clothing and wash clothing before reuse

Product should be opened and poured in a well ventilated area

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are barrier laminate or viton (14 mils or greater) If you want more options follow the instructions for category G or H on an EPA chemical resistance category selection chart

Pilots must wear

Long sleeved shirt and long pants and
Shoes and socks

See engineering controls for additional requirements

Mixers loaders applicators and other handlers must wear the following

Coveralls worn over short sleeved shirt and short pants
Chemical resistant footwear plus socks and
Chemical resistant gloves

See engineering controls for additional requirements

User Safety Requirements

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product s concentrate Do not reuse them

Engineering controls

Pilots must use an enclosed cockpit that meet the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170 240(d)(6)]

Human flagging is prohibited Flagging to support aerial application is limited to use of the Global Positioning System (GPS) or mechanical flaggers

USER SAFETY RECOMMENDATIONS

Users should

Wash hands with plenty of soap and water before eating drinking chewing gum using tobacco or using the toilet
Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish aquatic invertebrates oysters and shrimp Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean water mark Do not apply when weather conditions favor drift from treated areas Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas Do not contaminate water when disposing of equipment wash waters

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area

PHYSICAL/CHEMICAL HAZARDS

Do not use or store near heat or open flame

**DIRECTIONS FOR USE
RESTRICTED USE PESTICIDE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

4/10

Resistance Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with any thing that has been treated, such as plants, soil, or water is:

- Coveralls
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material

BUFFER ZONES

Vegetative Buffer Strip

Construct and maintain a minimum 10 foot wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds, estuaries, and commercial fish farm ponds).

Only apply products containing cypermethrin (Holster™ Insecticide) onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers:

Conservation Buffers to Reduce Pesticide Losses: Natural Resources Conservation Services, USDA NRCS, 2000, Fort Worth, Texas, 21 pp. <http://www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf>

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast) Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for ULV Aerial Application Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for Non ULV Aerial Application Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Spray Drift Requirements

Wind Direction and Speed

Only apply this product if the wind direction favors on target deposition.

Do not apply when the wind velocity exceeds 15 mph.

Temperature Inversion

Do not make aerial or ground applications into temperature inversions.

Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only medium or coarser spray nozzles (for ground and non ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications

Wind speed must be measured adjacent to the application site on the upwind side immediately prior to application
For ground boom applications apply using a nozzle height of no more than 4 feet above the ground or crop canopy
For airblast applications turn off outward pointing nozzles at row ends and when spraying the outer two rows To minimize spray loss over the top in orchard applications spray must be directed into the canopy

Additional Requirements for Aerial Applications

The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter
Flight speed and nozzle orientation must be considered in determining droplet size
Spray must be released at the lowest height consistent with pest control and flight safety Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety
When applications are made with a cross wind the swath will be displaced downwind The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind

CHEMIGATION USE DIRECTIONS

Apply this product only through sprinkler including center pivot lateral move end tow side (wheel) roll traveler big gun solid set or hand move irrigation systems Do not apply this product through any other type of irrigation system Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system

Crop injury lack of effectiveness or illegal residues in the crop can result from nonuniform distribution of treated water If you have questions about calibration you should contact State Extension Service specialists equipment manufacturers or other experts A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise

The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow

The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump

The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops

The irrigation line or water pump motor must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected

Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock

Holster Insecticide should be applied continuously for the duration of the water application Holster Insecticide should be diluted in sufficient volume to insure accurate application over the area to be treated When using chemigation a minimum of 0.5 inch per acre of irrigation water is recommended Agitation is not required when a suitable diluent is used

APPLICATION INSTRUCTIONS

Rotational Crops

Rotational crops may not be planted within 30 days after last application
Use low rate under light to moderate infestation Higher rates should be used under heavy insect pressure The rate of application is variable according to insect pressure timing of spray and field scouting

Tank Mixture

Holster Insecticide may be applied in tank mixtures with other products approved for use on cotton brassicas bulb vegetables head lettuce and pecans Observe all restrictions and precautions which appear on the labels of these products

For improved control of aphids and mites on cotton Holster Insecticide may be tank mixed with Curacron® 8E The tank mixture is most effective when pest populations are first observed Subsequent applications may be needed to maintain control

6/10

HOLSTER™ INSECTICIDE
EPA REG NO 34704 875

Maximum Usage When Applying Both Zeta Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season

Do not apply more than the maximum seasonal total for either product when used alone and do not apply more than the combined maximum seasonal total for both products as outlined in the table below

Crop	Maximum Seasonal Total for Either Product Used Alone (lbs AI/acre)			Maximum Seasonal Total When Applying Both Products to the Same Crop (lbs AI/acre)
	Zeta cypermethrin*		Cypermethrin *	Zeta cypermethrin* plus Cypermethrin**
	Fury® or Mustang®	Mustang Max™**		
Cotton	0.3	0.15	0.4	0.4
Head and Stem brassica	0.3	0.15	0.6	0.6
Leafy brassica greens	0.3	0.15	0.4	0.4
Head lettuce	0.3	0.15	0.6	0.6
Bulb vegetables	0.25	0.125	0.5	0.5
Pecans	0.3	0.15	0.5	0.5

Fury or Mustang (1.5 EC or 1.5 EW) Mustang Max (0.8 EC)
 Any cypermethrin product approved for crop use

Cotton
(14 day phi)
Specific Crop Restrictions

Maximum single application rate of 0.1 pound active ingredient per acre Minimum re treatment interval of 5 days
 Maximum annual application rate of 0.4 pound active ingredient per acre per year

Do not make more than 10 synthetic pyrethroid applications (of one product or combinations of products) to cotton in 1 growing season

PEST	DOSAGE		REMARKS
	LB AI/A	FL OZS/A	
Preemergent Use Cutworms	0.025 to 0.10	1.3 to 5.0	Use Holster Insecticide in the time period from 14 days prior to planting up to emergence of the crop. Apply as a broadcast spray by ground or air banded (including T band) or in furrow spray using sufficient spray volume to achieve adequate coverage. Reduced volumes of water may be used with specialized equipment. Use a minimum of 1.0 gal of water/A by air. Use the higher rates of Holster insecticide when incorporating into the soil.
Foliar Use Cutworms Soybean (banded) thrips Tobacco thrips	0.025 to 0.10	1.3 to 5.0	
Foliar Use Boll weevil Cabbage looper Cotton bollworm Cotton fleahopper Cotton leaf perforator European corn borer Fall armyworm Lygus bugs Other plant bugs Pink bollworm Saltmarsh caterpillar Tarnished plant bug Tobacco budworm White flies Yellow striped armyworm	0.04 to 0.10 0.10	2.0 to 5.0	Holster Insecticide may be applied in water or refined vegetable oil. When water is used, apply a minimum of 1.0 gal of finished spray/A by air or 5.0 gal of finished spray with ground equipment. When applying in water by air, 1.0 qt of emulsified oil may be substituted for 1.0 qt of water in the finished spray. When using oil, use a minimum of 1.0 qt/A in the finished spray. Holster Insecticide may be injected into overhead sprinkler irrigation water provided 1) an anti backflow check valve is present between the injection port and the water source 2) a check valve is present in the line to prevent irrigation water from entering the chemical supply tank and 3) the injection system has interlocking on off irrigation switches. For Boll weevil control, apply Holster Insecticide at a 5 day interval until pest numbers are reduced to acceptable levels. A maximum of 0.4 lb AI may be applied/A/season. Do not graze or feed cotton for forage.
Foliar use Beet armyworm	0.06 to 0.10	3.0 to 5.0	

Aids in control

For control of Beet armyworms only in the high plains of Texas, Arizona, and California

7/10

HOLSTER™ INSECTICIDE
EPA REG NO 34704 875

Head and Stem brassica vegetables including Broccoli Chinese broccoli (Gai Ion White flowering broccoli) Brussels sprouts Cauliflower Cavalo broccolo Kohlrabi Cabbage Chinese cabbage (Napa) (tight heading varieties only) (1 day phi)

Specific Crop Restrictions

Maximum application rate of 0.1 pound active ingredient per acre
Minimum re treatment interval of 7 days
Maximum seasonal application rate of 0.6 pound active ingredient per acre

PEST	DOSAGE		REMARKS
	LB AI/A	FL OZS/A	
Aphids Corn earworm Cucumber beetles Cutworms Flea beetles Imported cabbageworm Leafhoppers Lygus bugs Saltmarsh caterpillar Tobacco budworm Whiteflies	0.05 to 0.10	2.5 to 5.0	Apply in water as necessary for insect control using a minimum of 15.0 gals of finished spray/A with ground equipment and 5.0 gals/A by air. Lower rates of Holster Insecticide should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations but never to exceed the maximum allowable rate. Do not make applications less than 7 days apart. A maximum of 0.6 lb AI may be applied/A/season. In areas where arid climatic conditions persist such as California and Arizona higher than minimum recommended rates may be required.
Armyworms Crickets Ground beetles Loopers Onion thrips Stinkbugs Wireworm (adults) Aids in control	0.075 to 0.10	3.75 to 5.0	

Leafy brassica greens including Broccoli raab (rapini) Chinese cabbage (Bok choy) Chinese mustard cabbage (Gai choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens (1 day phi)

Specific Crop Restrictions

Maximum application rate of 0.1 pound active ingredient per acre
Minimum re treatment interval of 7 days
Maximum seasonal application rate of 0.4 pound active ingredient per acre

PEST	DOSAGE		REMARKS
	LB AI/A	FL OZS/A	
Aphids Corn earworm Cucumber beetles Cutworms Flea beetles Imported cabbageworm Leafhoppers Lygus bugs Saltmarsh caterpillar Tobacco budworm Whiteflies	0.05 to 0.10	2.5 to 5.0	Apply in water as necessary for insect control using a minimum of 15.0 gals of finished spray/A with ground equipment and 5.0 gals/A by air. Lower rates of Holster Insecticide should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations but never to exceed the maximum allowable rate. Do not make applications less than 7 days apart. A maximum of 0.4 lb AI/A/season. In areas where arid climatic conditions persist such as California and Arizona higher than minimum recommended rates may be required.
Armyworms Crickets Ground beetles Loopers Onion thrips Stinkbugs Wireworm (adults) Aids in control	0.075 to 0.10	3.75 to 5.0	

8/10

HOLSTER™ INSECTICIDE
EPA REG NO 34704 875

Lettuce head

(5 day phi)

Specific Crop Restrictions

Maximum application rate of 0.1 pound active ingredient per acre

Minimum re treatment interval of 7 days

Maximum seasonal application rate of 0.6 pound active ingredient per acre

PEST	DOSAGE		REMARKS
	LB AI/A	FL OZS/A	
Corn earworm Cucumber beetles Cutworms Diamondback moth Flea beetles Imported cabbageworm Leafhoppers Lygus bugs Saltmarsh caterpillar Tobacco budworm	0.05 to 0.10	2.5 to 5.0	Apply in water as necessary for insect control using a minimum of 15.0 gals of finished spray/A with ground equipment and 5.0 gals/A by air. Lower rates of Holster Insecticide should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations but never to exceed the maximum allowable rate. A maximum of 0.6 lb AI may be applied/A/season. In areas where arid climatic conditions persist such as California and Arizona higher than minimum recommended rates may be required.
Armyworms Crickets Loopers Onion thrips Stinkbug	0.075 to 0.10	3.75 to 5.0	

Bulb vegetables (*Allium* spp) including Garlic, Garlic Great headed (elephant), Green eschalots, Japanese bunching onions, Leeks, Onion, Dry bulb and green, Onion, Welch, Shallots, Dry bulb and green, Spring onion or Scallions

(5 day phi)

Specific Crop Restrictions

Maximum application rate of 0.1 pound active ingredient per acre

Minimum re treatment interval of 7 days

Maximum seasonal application rate of 0.5 pound active ingredient per acre

PEST	DOSAGE		REMARKS
	LB AI/A	FL OZS/A	
Onion thrips	0.08 to 0.10	4.0 to 5.0	Apply Holster Insecticide in a minimum of 20.0 gals/A with ground equipment or in a minimum of 3.0 gals/A by aircraft. Begin applications when pests appear. Do not make applications less than 7 days apart.
Aphids Armyworms Cutworms Leafminers Onion maggot adults Stink bugs	0.04 to 0.10	2.0 to 5.0	

To control Onion thrips
Use higher rates as population increases but do not exceed the maximum allowable rate and avoid rescue situations.
Use of a crop oil concentrate at 1.0 fl oz/A is recommended.
A maximum of 0.5 lb AI may be applied/A/season.
Do not graze livestock in treated areas or cut treated crops for feed.

9/10

HOLSTER™ INSECTICIDE
EPA REG NO 34704 875

Pecans
(21 day phi)

Specific Crop Restrictions

Maximum single application rate of 0.1 pound active ingredient per acre
 Minimum re treatment interval of 7 days
 Maximum annual application rate of 0.5 pound active ingredient per acre per year

PEST	DOSAGE		REMARKS
	LB AI/A	FL OZS/A	
Black pecan aphid Hickory shuckworm Pecan nut casebearer Pecan weevil Yellow pecan aphid	0.06 to 0.10	3.0 to 5.0	Applications at the lower rate should be made when pest populations are low. Rates should be increased as the pest pressure increases. Apply by ground equipment to the point of drip. Use 100 gals of dilute spray/A for smaller trees. For larger trees which require higher gallonage to achieve adequate coverage, apply in 200 to 300 gals of water. In order to calculate the correct number of gals of water needed to spray 1 acre of your trees to the point of drip, you may need to conduct a test. If you do not know how to conduct such a test with your equipment, you should request assistance from your equipment dealer. Up to 0.5 lb AI/A/season may be applied prior to shuck split. Do not graze livestock in treated orchards or cut treated cover crops for feed.

DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY

STORAGE AND DISPOSAL

PESTICIDE STORAGE Do not store below 10 F (12 C)

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL Nonrefillable container Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

In case of spill: avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1 800 424 9300

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

HOLSTER™ INSECTICIDE
EPA REG NO 34704 875

10/10

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD AS IS, AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC. ATTENTION LAW DEPARTMENT, P O BOX 1286, GREELEY, CO 80632 1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING: AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER, DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Fury and Mustang are registered trademarks and Mustang MAX™ is a trademark of FMC Corporation. Curacron is a registered trademark of a Syngenta Group Company.

Curacron, Fury, Mustang, and Mustang MAX are restricted use pesticides.

FORMULATED FOR
LOVELAND PRODUCTS, INC.
P O BOX 1286 GREELEY, COLORADO 80632 1286