
 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <u> X </u> Registration <u> </u> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number:	Date of Issuance:
	34704-913	JAN 10 2006
	Term of Issuance: Conditional	
Name of Pesticide Product: SMZ 4L		
Name and Address of Registrant (include ZIP Code): Loveland Products, Inc. 7251 W 4 th Street, PO Box 1286 Greeley, CO 80632-1286		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided you agree in writing to:</p> <ol style="list-style-type: none"> 1. Change EPA Reg. No. on label from 34704- to 34704-913. 2. Change the PRECAUTIONARY STATEMENTS to the following: "Harmful if absorbed through skin. Causes moderate eye irritation. Harmful if swallowed or inhaled. Avoid breathing vapors or spray mist. Avoid contact with skin, clothing, or eyes." 3. On page 4, in the statement "Apply the mixture in 10-40 gals of water per acre as a postemergence spray to the weeds at the appropriate weed growth stage given on the Roundup label." add "or Touchdown" after "Roundup". 4. On page 5, in the second column, remove the section CHRISTMAS TREE PLANTINGS AND SHELTERBELTS up until the section NURSERIES, CHRISTMAS TREE PLANTINGS AND SHELTERBELTS, as this information is already in the section NURSERIES, CHRISTMAS TREE PLANTINGS AND SHELTERBELTS. 		
Signature of Approving Official:  James Tompkins, Product Manager (25) Herbicide Branch, Registration Division (7505C)	Date: 1-10-06	

5. On page 6, in the section Warranty Disclaimer And Notice, add "To the extent allowed by law" before the statement "Buyers or seller's total liability, shall be limited to damages not exceeding the cost of the product." In the statement "Except as expressly stated herein, Loveland Product Inc, the Manufacturer or Seller makes no warranty of results to be obtained by the use of this product" add "other" before the word "warranty."

You will submit one (1) copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). A stamped copy of labeling is enclosed for your records.

If you have any questions, please contact Hope Johnson at 703-305-5410.



James Tompkins
Product Manager (25)
Herbicide Branch
Registration Division (7505C)

378

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 10 2006

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

34704-913



SMZ 4L

Herbicide

FOR WEED CONTROL IN CERTAIN CROPS

ACTIVE INGREDIENT:	
Simazine: 2-chloro-4,6-bis(ethylamino)-s-triazine	41.9%
OTHER INGREDIENTS:	58.1%
TOTAL	100.0%

SMZ 4L contains 4 lbs. active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:
1-800-301-7976.

EPA REG NO. 34704-

EPA EST. NO. 34704-MS-1

NET CONTENTS 2½ GALS. (9.46 L)

HT

01/06

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes eye irritation. Harmful if inhaled or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with eyes, skin, or clothing.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton, and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separate from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets

the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Simazine is a chemical which can travel (seep or leach) through soil and enter ground water which may be used as drinking water. Simazine has been found in ground water as a result of its use as a herbicide. Users of this product are advised not to apply Simazine where the water table (ground water) is close to the surface and where the soils are very permeable, i.e., well-drained soils, such as loamy sands. Users are advised to consult with their local agricultural agencies to obtain information on the location of ground water and the type of soil in their area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Coveralls, chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton, and shoes plus socks.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

PESTICIDE STORAGE: Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

PESTICIDE DISPOSAL: Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent), puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.

4 8 8

SMZ 4L
EPA REG. NO. 34704-

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR WEED CONTROL, AND/OR ILLEGAL RESIDUES.

Do not apply this product through any type of irrigation system.

This product is not for use in formulating end-use products for aquatic use.

GENERAL INFORMATION

Apply SMZ 4L herbicide before weeds emerge or after removal of weed growth. SMZ 4L controls a wide variety of annual broadleaf and grass weeds when used at selective rates in agricultural crops.

Where a range of application rates is given, use the low rate on coarse-textured soil and soil lower in organic matter; use the high rate on fine-textured soil and soil higher in organic matter.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

Since SMZ 4L enters weeds mainly through their roots, moisture is needed to move it into the root zone. Very dry soil conditions following application may necessitate shallow cultivation or rotary hoeing.

SMZ 4L is nonflammable.

Annual Weeds Controlled

alyssum	fivehook bassia	rattail fescue
annual bluegrass	Flora's paintbrush	redmaids
annual morning glory	Florida pusley	Russian thistle
annual ryegrass	foxtails	shepherdspurse
barnyardgrass	goosegrass	signalgrass
(watergrass)	groundsel	(Brachiaria spp.)
burclover	henbit	silver hairgrass
carelessweed	juglerice	smartweed
carpetweed	knawel (German moss)	spanishneedles
common chickweed	common lambsquarters	speedwell
crabgrass (Digitaria spp.)	nightsshade	tansymustard
downy brome	pepperweed	wild mustard
(cheatgrass)	pigweed	wild oats
fall panicum	pineappleweed	witchgrass
fiddleneck	prickly lettuce	yellow flower
filaree	common purslane	pepperweed
fireweed	ragweed	

Resistance Management

SMZ 4L is a Group 5 Herbicide (contains the active ingredient simazine).

Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected, and weeds controlled by this product are expected to be present along with resistant biotypes, we recommend the use of this product in registered combinations or in sequence with other registered herbicides which are not triazines, i.e. which are not solely a Group 5 Herbicide. If only resistant biotypes are expected to be present, use a registered herbicide which is not solely a Group 5 Herbicide. Consult with your State Agricultural Extension Service for specific recommendations.

Perennial Crop Replanting

If replanting perennial crops, do not apply SMZ 4L in the year preceding planting these crops, unless otherwise state on this label, or crop injury may occur.

APPLICATION PROCEDURES

Ground application: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 20 gals. of spray mixture per acre.

Use a pump with capacity to: (1) maintain 35-40 psi at nozzles, (2) provide sufficient agitation in tank to keep mixture in suspension, and (3) to provide a minimum of 20% bypass at all times. Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gals./minute/100 gal. tank size circulated through a correctly positioned sparger tube or jets.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom.

and where required, at the nozzles. Check nozzle manufacturer's recommendations. For band applications, calculate amount to be applied per acre as follows:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \frac{\text{broadcast rate}}{\text{per acre}} = \frac{\text{amount needed}}{\text{per acre of field}}$$

Aerial application: Use aerial application only where specified in the use directions. Apply in a minimum of 1 gal. of water for each quart of SMZ 4L applied per acre. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the [Aerial Drift Reduction Advisory Information](#).

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

Controlling Droplet Size

- **Volume** – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** – Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** – Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** – Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger than other nozzle types.
- **Boom Length** – For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.
- **Application** – Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a

SMZ 4L EPA REG. NO. 34704-

ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

MIXING PROCEDURES

All Uses: (1) Be sure sprayer is clean and not contaminated with any other materials, or crop injury or sprayer clogging may result. (2) Fill tank ¼ full with clean water, nitrogen solution, or complete fluid fertilizer. (3) Start agitation. (4) Be certain that the agitation system is working properly and creates a rippling or rolling action on the liquid surface. (5) If any dry herbicides are being tank mixed, add them to the tank and allow them to become thoroughly suspended. (6) Then pour SMZ 4L directly from jug into tank. (7) Continue filling tank until 90% full. Increase agitation if necessary to maintain surface action. (8) Add other liquid tank mix herbicide(s) after this product is thoroughly suspended. (9) Finish filling tank. Maintain agitation to avoid separation of materials. (10) Empty tank as completely as possible before refilling to prevent buildup of emulsifiable concentrate residue from possible tank mix herbicides. (11) If an emulsifiable concentrate film starts to build up in tank, drain it and clean with strong detergent solution or solvent. (12) Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent.

Compatibility test: To determine the tank mix compatibility of SMZ 4L with liquid fertilizer, crop oil, spreaders, or other recommended pesticides, use this test method. Nitrogen solutions or complete fluid fertilizers may replace all or part of the water in the spray in corn. Since liquid fertilizers can vary, even within the same analysis, check compatibility each time before use. Be especially careful when using complete suspension or fluid fertilizers, as serious compatibility problems are more apt to occur. Commercial application equipment may improve compatibility in some instances. Check compatibility using this procedure:

1. Add 1 pt. of fertilizer to each of two 1-qt. glass jars with tight lids.
2. To one of the jars, add ¼ tsp. of compatibility agent approved for this use (¼ tsp. is equivalent to 2 pts./100 gals. spray). Cap and shake until mixed. Examples of compatibility agents include Compex® and Unite®.
3. To both jars, add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. After each addition, cap and shake until thoroughly mixed. The appropriate amount of herbicides for this test follows (assuming a spray volume of 25 gals./A):
Dry Herbicides: For each pound per acre, add 1.5 teaspoons to each jar. Fluff up wettable powder products before measuring.

Liquid herbicides: For each pint per acre, add 0.5 teaspoon to each jar.

For a spray volume other than 25 gals./A, change the teaspoons added to each jar as follows:

$$\frac{25 \text{ gals.} \times \text{no. teaspoons given above}}{\text{desired gals. spray volume per acre}} = \text{teaspoons to add to each jar}$$

4. After adding all ingredients, put lids on and tighten. Shake jars vigorously one minute. Let the mixtures stand 15 minutes and then look for separation, large flakes, precipitates, grease, gels, medium to heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the 2 jars. If either mixture separates, but can be remixed readily, the mixture can probably be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) slurry the dry herbicide(s) in water before addition, or (B) add ½ of the compatibility agent to the fertilizer and the other ½ to the emulsifiable concentrate or flowable herbicide before addition to the mixture. If still incompatible, do not use the materials mixed in the same spray tank.

CROP USE DIRECTIONS

FRUIT AND NUT CROPS

Apply the spray to the orchard or vineyard floor, avoiding contact with fruit, foliage, or stems. Recommended rates are based on broadcast treatment. For band applications or spot applications around trees in fruit or nut plantings, reduce the broadcast rate of SMZ 4L and water per acre in proportion to the area actually sprayed.

Precautions (all fruit and nut crops): To avoid crop injury, (1) Apply only to orchards or groves where trees have been established 12 months or more, unless specified differently. (2) Make only one application per year, except as noted otherwise. (3) Do not use on gravelly, sand, or loamy sand soil. (4) Immediately following application, limit overhead sprinkler irrigation to ½ inch.

Almonds, Peaches and Nectarines (CA Only)

Apply 1-2 qts. of SMZ 4L per treated acre in a 2-4 ft. band on each side of the tree row. Apply before weeds emerge in late fall or early winter. Weeds controlled by 1 qt. of SMZ 4L include burdock, common chickweed, wild mustard, and shepherdspurse. Apply only once per year.

Precautions: To avoid crop injury, (1) Do not treat trees established in the grove less than 3 years. (2) Do not treat the Mission (TX) variety of almonds. (3) Do not apply to almond trees propagated on plum rootstocks. (4) Do not replant almonds, peaches, or nectarines in treated soil for 12 months after treatment. (5) Do not apply on soil with less than 1% organic matter. (6) Do not treat areas where water will accumulate.

Apples, Pears, and Sour Cherries

Apply 2-4 qts./A.

Avocados CA and FL Only:

Apply 2-4 qts./A after final preparation of grove.

Blueberries and Caneberries (Blackberries, Boysenberries, Loganberries, and Raspberries)

Apply 2-4 qts./A in the spring or apply a split application of 2 qts./A in the spring plus 2 qts./A in the fall. Apply in a minimum of 40 gals. of water per acre.

On plantings less than 6 months old, use ½ the above rate.

To control quackgrass, apply 4 qts./A in the fall or split the application, applying 2 qts./A in the fall plus 2 qts./A in the spring, when quackgrass is actively growing.

Note: Do not apply when fruit is present, or illegal residues may result.

Cranberries

MA: Apply up to 4 qts./A, either before spring growth begins or in the fall after harvest.

Other areas: Apply 2 qts./A before spring growth begins.

Filberts

Apply 2-4 qts./A in the fall or apply a split application of 2 qts./A in the fall plus 2 qts./A in the spring.

Precaution: If trees are planted on a hillside, excessive soil erosion may result from the elimination of weeds.

Note: Do not apply when nuts are on the ground during the harvest period, or illegal residues may result.

Grapes

Apply 2-4.8 qts./A any time between harvest and early spring.

Precaution: Do not use in vineyards established less than 3 years, or crop injury may occur.

Grapefruit, Lemons, and Oranges

AZ (Lemons and Oranges Only): Apply a split application of 1.6 qts./A in the spring plus 1.6 qts./A in the fall.

CA: Apply 2-4 qts./A in a single application, or apply 2 qts./A in the fall and 2 qts./A in the spring.

Precautions: Do not use in the Imperial, Coachella, or Palo Verde Valleys, or crop injury may occur.

FL (Grapefruit and Oranges Only): Apply 4 qts. of SMZ 4L to weed-free soil during the spring and/or fall to control weeds expected to emerge during these periods. Apply prior to emergence of weeds or if weeds have emerged, apply in tank mixture with a contact herbicide. Use caution to keep the treatment off the foliage, fruit, or trunk of citrus trees. For control of difficult species, such as balsam poplar vine and spanishneedles, and partial control of honeyvine milkweed, apply 8 qts. of SMZ 4L as a single application in the spring as a 50% band application to the grove acre. Apply in the spring growing season between January and April. Do not make a fall SMZ 4L application if this treatment was used in the spring. When emerged weeds are present, apply SMZ 4L in tank mixture with a recommended contact herbicide. Follow all directions, precautions, limitations, etc. on the tank mix products.

TX (Grapefruit and Oranges Only): Apply 4-4.8 qts./A.

Precautions (all areas): To avoid crop injury, (1) Do not use in nurseries. (2) Do not apply to bedded grapefruit, lemons, or oranges (except for FL grapefruit and oranges). (3) Do not apply to trees under stress from freeze damage for one year after the freeze. (4) In FL, do not exceed 8 qts. of SMZ 4L during any one growing season. (5) Do not exceed 4 qts./acre/year of SMZ 4L on grove planted trees 1 year old or less, or crop injury may occur. **Note:** Dark-red grapefruit hybrids, e.g. "Star Ruby", have a higher risk of crop injury than non-dark-red types.

Macadamia Nuts

Apply 2-4 qts. in 50 gals. of water per acre before harvest and just prior to weed emergence. Repeat application as necessary.

Note: Do not apply when nuts are on the ground during the harvest period, as illegal residues may result.

Oilives

Apply 2-4 qts./A following grove preparation in the fall. Repeat annually in midwinter.

Peaches, Plums, and Sweet Cherries

Apply 1.6-4 qts./A. Apply in late fall to early spring prior to weed emergence.

Precautions: To avoid crop injury, (1) Peaches: use only in AR, LA, MO, OK, TX and states east of the Mississippi River. For CA, see specific directions in the Almonds, Peaches and Nectarines (CA Only) section. (2) Plums and sweet cherries: use only in MO and states east of the Mississippi River, except TN.

Pecans

Apply 2-4 qts./A before weeds emerge in the spring.

Precautions: To avoid crop injury, (1) Do not use west of the Pecos River in TX or in AZ, CA, or NM. (2) Do not make applications to transplanted trees that have been established less than 2 years in the grove.

Note: Do not apply when nuts are on the ground, or illegal residues may result. Do not allow animals to graze treated areas.

Strawberries

OR and WA: For control of chickweed, groundsel, mustard and shepherdspurse, apply broadcast 1 qt./A. in fields where overhead irrigation is used to activate this product; apply

6 8 8

SMZ 4L EPA REG. NO. 34704-

after harvest at time of bed renovation. In fields where overhead irrigation is not available, apply during early October through November.
Precautions: To avoid crop injury, (1) Make only one application per growing season. (2) Do not apply within 4 months after transplanting.

Walnuts

Apply 2-4 qts/A. Leveling and trowing operations after application will lessen effectiveness of weed control.
Note: Do not apply when nuts are on the ground, or illegal residues may result.

SMZ 4L Plus Bromacil 80W for Grapefruit and Oranges (FL Only)

Use in grapefruit and oranges in FL for control of balsamapple, black nightshade, carpetweed, crabgrass, cudweed, dayweed, Florida pusley, horseweed, pepperweed, pigweed, poorjoe, ragweed, rattlebox, spanishneedles, and sandbur, and for partial control of bermudagrass, bahiagrass, pangolagrass, paragrass, and torpedograss. Apply 4 qts. of SMZ 4L plus 3-4 lbs. of bromacil 80W per acre beneath trees in a minimum of 40 gals. of water per acre before or soon after weed growth begins. Use caution to keep the spray off the foliage, fruit or trunk of citrus trees. When mixing, add bromacil 80W slurry to water in spray tank, agitate thoroughly, then add SMZ 4L and agitate thoroughly again. Use the lower rates for light weed infestations or all applications in bedded citrus areas. Use the higher rates for heavy weed infestations only in ridge grown citrus areas. Temporary yellowing of citrus leaves may occur following treatment.

Precautions: To avoid crop injury, (1) Do not use in nurseries, or where trees are under stress from freeze damage for one year after the freeze. (2) Do not use on soil with less than 1% organic matter or on poorly drained soil. (3) Do not treat trees planted in irrigation furrows. (4) Do not treat diseased trees such as those with foot rot. (5) Do not use in groves interplanted with other trees or desirable plants, nor in home grapefruit or orange plantings, or in areas where roots of other valuable plants or trees may extend. (6) Treated areas may be planted to citrus trees one year after application. (7) Do not rotate to other crops within 2 years after application.

Note: Apply only once per year and avoid contact with foliage and fruit with spray or mist, or illegal residues may result.

SMZ 4L Plus Gramoxone® Max

This tank mix is effective in the following fruit and nut crops for kill of existing vegetation and for residual control of the annual broadleaf and grass weeds claimed for SMZ 4L applied alone. This combination is also effective for top kill and suppression of perennial weeds. In FL, this mixture may be applied in spring or fall to emerged weeds.

Use this tank mix on these crops:

almonds (CA only)	macadamia nuts
apples	olives
avocados (CA and FL ¹ only)	oranges ²
cherries (sour and sweet ⁴)	peaches ³
filberts	pears
grapefruit (CA and TX only)	pecans
grapes	plums ⁴
lemons (AZ and CA only)	walnuts

Apply the SMZ 4L rate given under the appropriate crop on this label plus the appropriate labeled rate of Gramoxone Max in 50-200 gals. (30-50 gals. for pecans) of water per acre to the orchard floor, avoiding contact with fruit, foliage, or stems. Add a nonionic surfactant, such as X-77®, at 0.5 pt./100 gals. of spray. Apply when weeds are succulent and new growth is 1-6 inches tall. For mature woody weeds or difficult to control perennial weeds, re-treat or spot treat with Gramoxone Max if regrowth occurs.

Add SMZ 4L to the spray tank first (refer to Mixing Procedures section of this label), then add Gramoxone Max, and add the surfactant last. Provide constant agitation during mixing and application to keep the mixture in suspension.

Precautions: To avoid crop injury, (1) Apply the tank mix only once per year. (2) Use a shield for young trees or vines.

Refer to the SMZ 4L and Gramoxone Max labels for further directions, specific weeds controlled, and precautions and limitation on each crop.

- ¹ In avocados in FL, this tank mix also controls balsamapple, rattail amaranth, and at the higher rate of each herbicide, it suppresses coral vine.
- ² In oranges in FL, apply 4 qts. of SMZ 4L per acre per application. Do not exceed 8 qts. of SMZ 4L during any one growing season.
- ³ Limited to AR, CA, LA, MO, OK, TX, and states east of the Mississippi River. As appropriate, refer to SMZ 4L sections Almonds, Peaches and Nectarines (CA only) or Peaches, Plums, and Sweet Cherries for rate of SMZ 4L and other information.
- ⁴ Limited to MO and states east of the Mississippi River, except TN.

Tank Mixture with Roundup® or Touchdown Brand Herbicides

This tank mixture is effective in grape vineyards and in the following bearing or non-bearing tree crops for control of existing vegetation and for residual control of the annual broadleaf and grass weeds claimed for this product applied alone. This combination is also effective for partial control of perennial weeds contracted by the spray mixture during application.

almonds** (CA only)
apples*
avocados* (CA and FL only)
cherries (sour and sweet****2)
filberts**
grapefruit*** (CA, FL, and TX)
grapes*
lemons* (AZ and CA)

macadamia nuts**
oranges*** (AZ, CA, FL, and TX)
peaches****1
pears*
pecans**
plums****2
walnuts**

Use the appropriate rate given elsewhere on this label for SMZ 4L applied alone to the crop being treated. Add to the spray tank the appropriate labeled rate of a Roundup or Touchdown brand product per acre, depending on weeds present and their growth stage. Also add an agriculturally approved nonionic surfactant at 0.5% by volume of spray solution. Apply the mixture in 10-40 gals. of water per acre as a postemergence spray to the weeds at the appropriate weed growth stage given on the Roundup label.

Add SMZ 4L to the spray tank first, then add the Roundup or Touchdown brand product. Provide constant agitation during mixing and application to keep the mixture in suspension. Refer to Application Procedures section of this label for further directions.

Precaution: To avoid crop injury, take extreme care to avoid contact of herbicide solution, spray, drift, or mist with foliage or green bark of trunk, branches, suckers, fruit, or other parts of trees or vines. Observe precautions on both the SMZ 4L and Roundup or Touchdown labels for each crop involved.

Always refer to the labels of both herbicides for further directions, specific weeds controlled, and precautions and limitation on each crop.

- *Allow a minimum of 14 days between last application and harvest.
- **Allow a minimum of 21 days between last application and harvest of these crops.
- ***Allow a minimum of 1 day between last application and harvest of these crops
- ****Allow a minimum of 17 days between last application and harvest of these crops.

- ¹ Limited to AR, CA, LA, MO, OK, TX, and states east of the Mississippi River. For CA, see specific directions in Almonds, Peaches and Nectarines (CA only) section.
- ² Plums and sweet cherries: limited to MO and states east of the Mississippi River, except TN.

Tank Mixture and Solicam®

For improved control of such weeds as clover, cutleaf evening primrose, dandelion, henbit, horseweed or marehail, lambsquarters, and puncturevine, apply SMZ 4L in tank mixture with Solicam DF on these crops:
oranges grapefruit lemons

Apply 4 qts./A of SMZ 4L plus 4-5 lbs. of Solicam DF Herbicide (78.6% active ingredient) in 20-100 gals. of water per acre. SMZ 4L + Solicam may be applied in tank mixture with Gramoxone Extra or Roundup.

Follow the labeled directions and restrictions of use on labels of the herbicides used in mixtures with SMZ 4L.

Precaution: Keep SMZ 4L + Solicam mixtures from contacting foliage, fruits, and stems of citrus trees during spraying.

SMZ 4L Plus Surflan®

Use this tank mix in the following crops for preemergence control of all weeds claimed on both labels:

almonds	filberts	peaches
apples	grapes	pears
avocados	grapefruit	pecans
canberries	lemons	plums
cherries	oranges	English walnuts

Apply the SMZ 4L rate given under the appropriate crop on this label plus 2.67-5.33 lbs. of Surflan 75W or 2-4 qts. of Surflan A.S. in 20-40 gals. of water per acre. Refer to the Surflan 75W or Surflan A.S. label for complete tank mix directions, and use 0.8 qt. of SMZ 4L for each pound of SMZ 4L for each pound of Princep 80W. Observe all precautions and limitations on the SMZ 4L and Surflan labels.

FIELD CROPS

Corn

Nitrogen solution or complete liquid fertilizer may replace all or part of the water in the spray. Determine the physical compatibility of SMZ 4L with fertilizer before use. Refer to the General Information section of this label for a compatibility test procedure. Do not apply after corn has emerged, as there is danger of liquid fertilizer causing crop injury. Use a minimum of 10 gals. of spray mixture per acre.

Preemergence: Apply before weeds and corn emerge. Use the appropriate rate in the following table.

Preplant: Broadcast in the spring after plowing at the appropriate rate in the following table. Apply before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation of SMZ 4L. Best results will be obtained when SMZ 4L is applied within 2 weeks before planting. Under dry weather conditions, preplant applications may give better weed control than preemergence. If weeds develop, particularly under relatively dry conditions, a shallow cultivation will generally result in better weed control.

7 8 8

SMZ 4L
EPA REG. NO. 34704-

Soil Texture	Broadcast Rate Per Acre*
Sand, silt, and loam that is low in organic matter	4 pts.
Soil containing moderate amounts of clay and organic matter	4.8 pts.
Loam that is high in organic matter and clay (including dark prairie soils of Corn Belt)	6 pts.
Peat, muck, and high-organic clay	8 pts.

* For calculation of band treatment rate, see the General Information section.

Quackgrass Control: Broadcast 6-8 pts./A in the fall. Plow 2-3 weeks later, or, if erosion is a problem, delay plowing until spring.

Winter Annual Broadleaf Control - Preemergence Fall Application

For preemergence control of winter annual weeds, such as common chickweed, henbit, shepherds-purse, tansymustard, wild mustard, annual bluegrass, downy brome, and others, broadcast 1 qt./A of SMZ 4L after harvest of the preceding crop and prior to weed emergence on land to be planted to corn the following year. A tillage operation may precede the application. Do not apply to frozen ground. If SMZ 4L is used in the fall corn weed control program, do not exceed 2 qts. of SMZ 4L preemergence in the spring.

Precautions: (1) Do not apply more than 8 pts. of SMZ 4L per acre of corn per year. (2) Do not plant any crop except corn until the following year, or injury may occur. (3) Following harvest, plow and thoroughly till the soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of the rate used. (4) If SMZ 4L is used at a rate higher than 6 pts./A or equivalent band rate, a crop of untreated corn should precede the next rotational crop. (5) Do not apply SMZ 4L preplant incorporated for weed control in corn in the High Plains and Intermountain areas of the West (including central and western KS, western NE, western OK, and the Panhandle of TX) where rainfall is sparse and erratic or where irrigation is required. (6) In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use SMZ 4L only when corn is to follow corn or when a crop of untreated corn is to precede other rotational crops. (7) In western MN and eastern parts of the Dakotas, KS, and NE, do not rotate to soybeans if the rate applied was more than 4 pts./A or equivalent band rate, or injury may occur. (8) Injury may occur to soybeans planted in north-central and northwest IA, south-central and southwest MN, northeast NE, southeast SD, and other areas the year following application on soils having a calcareous surface layer. (9) Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes and grasses the year following SMZ 4L application, or injury may occur.

Note: Do not graze treated areas, or illegal residues may result.

Tank Mixtures on Corn

AAtrex®: Use SMZ 4L in a preplant or preemergence tank mixture with AAtrex Nine-O®, or AAtrex 4L for control of many annual weeds, including carpetweed, crabgrass, fall panicum, foxtail, lambsquarters, morningglory, pigweed, ragweed, and velvetleaf.

Apply at rates given below. Use the 1:1 ratio for most weeds and the 2:1 ratio for expected heavy infestations of crabgrass and fall panicum.

Soil Texture	Broadcast Rate Per Acre			
	1:1 Ratio		2:1 Ratio	
	SMZ 4L	AAtrex 4L*	SMZ 4L	AAtrex 4L*
Sand, loamy sand, sandy loam	1 qt.	1 qt.	1.33 qts.	0.67 qt.
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with low organic matter	1.2 qts.	1.2 qts.	1.6 qts.	0.8 qt.
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with medium to high organic matter, and clay (including dark prairie soils of the Corn Belt)	1.44 qts.	1.44 qts.	1.92 qts.	0.96 qt.

*When using AAtrex 4L or AAtrex Nine-O, use equivalent atrazine rates. One qt. of AAtrex 4L equals 1.1 lb. of AAtrex Nine-O.

Refer to the AAtrex label for complete directions and use 1 qt. of this product for each 1.1 lb of Princep Caliber 90. Also refer to the AAtrex label and the Corn section of this label for precautions and limitations.

Gramoxone Max: Use in a tank mixture where corn will be planted directly in a cover crop, established sod, or previous crop residues. This combination controls existing vegetation and provides residual control of the annual broadleaf and grass weeds listed under the General Information section.

Add SMZ 4L to the spray tank, mix thoroughly with water, and then add Gramoxone Max and a nonionic surfactant. Provide constant agitation during mixing and application to keep the mixture in suspension.

Apply 4-6 pts. of SMZ 4L plus the appropriate labeled rate of Gramoxone Max in 20-60 gals. of water per acre as a broadcast spray, either before or after planting, but before corn emerges. Add a nonionic surfactant at the rate of 0.5 pt./100 gals. of spray volume.

For further information, see general information, caution and warning statements, precautions, and notes on the SMZ 4L and the Gramoxone Max labels.

CHRISTMAS TREE PLANTINGS AND SHELTERBELTS

Remove weed growth before application. Apply 2-4 qts. in at least 25 gals. of water per acre after transplanting. Use the same rate for annual maintenance applications.

For quackgrass control, apply 4 qts./A in the fall or apply a split application of 2 qts./A in the fall plus 2 qts./A in early spring, after quackgrass begins growth.

Precautions: To avoid tree injury, (1) Do not use on seedbeds or cutting beds. (2) In CA, OR, and WA, do not apply to Christmas trees or shelterbelts sooner than one year after transplanting. In other areas, do not apply to Christmas trees or shelterbelt transplants less than 2 years of age. (3) Do not use until soil is firmly settled around roots. (4) Do not apply more than once a year, except as directed for quackgrass control.

Apply SMZ 4L to these species of trees and shrubs, as recommended above:

Conifers

arborvitae	Scotch pine	white fir
Austrian pine	white pine	hemlock
Bishop pine*	blue spruce	juniper
knobcone pine	Norway spruce	red cedar
lodgepole pine (shore pine)	red spruce	white cedar
Monterey pine*	white spruce	yew (Taxus spp.)
Mugho pine	balsam fir	
Red pine (Norway pine)	Douglas fir	
	Fraser fir	

* For CA only

Surflan Tank Mix: On Christmas tree plantings, use this tank mix for preemergence control of weeds listed on the Surflan 75W (or Surflan A.S.) and SMZ 4L label. Use on field grown conifer species listed on the labels for each herbicide plus grand fir, alpine fir, Englemann spruce, black spruce, Colorado blue spruce, Coulter pine, giant redwood, and Veitch fir. Broadcast the mixture as a directed spray to the soil surface or as an over-top spray, using 2-4 qts. of SMZ 4L and 2.67- 5.33 lbs. of Surflan 75W (or 2-4 qts. of Surflan A.S.). Apply in sufficient water per acre to uniformly treat the area. Follow over-top sprays with sprinkler irrigation to move the herbicide from leaf surfaces to the soil. Remove weed growth before application. Mix weed residues, prunings, or trash into the soil, or remove them before treatment. Soil should be in good till and free of clods at time of application. Shallow cultivation (1-2 inches) after treatment will not reduce weed control. Observe all precautions and limitations on the SMZ 4L and Surflan labels.

Note: Length of weed control may be reduced when continuous wet soil conditions follow herbicide application.

Precautions: To avoid plant injury, (1) Do not use on seedbeds or on unrooted cuttings. (2) Do not use in greenhouses or other enclosed areas.

NURSERIES, CHRISTMAS TREE PLANTINGS, SHELTERBELTS

Nurseries (see list below)

Apply 2-3 qts. in at least 25 gals. of water per acre in fall or spring.

Precautions: To avoid injury, do not apply for at least one year after transplanting.

Christmas Tree Plantings and Shelterbelts (see list below)

Remove weed growth before application. Apply 2-4 qts. in at least 25 gals. of water per acre after transplanting. Use at the same rate for annual maintenance applications. For quackgrass control apply 4 qts./A in the fall or apply a split application of 2 qts./A in the fall plus 2 qts./A in early spring, after quackgrass begins growth.

Precautions: to avoid tree injury, (1) Do not use SMZ 4L on seedbeds or cuttingbeds. (2) In CA, OR, and WA, do not apply to Christmas trees or shelterbelts sooner than one year after transplanting. In other areas, do not apply to Christmas trees or shelterbelts less than 2 years of age. (3) Do not use until soil is firmly settled around roots. (4) Do not apply more than once a year, except as directed for quackgrass control.

Apply SMZ 4L to these species of trees and shrubs, as recommended above:

Conifers

arborvitae	Red pine (Norway pine)	Douglas fir
Austrian pine	Scotch pine	Fraser fir
Bishop pine*	white pine	white fir
knobcone pine*	blue spruce	hemlock
lodgepole pine (shore pine)	Norway spruce	juniper
Monterey pine*	red spruce	red cedar
Mugho pine	white spruce	white cedar
	balsam fir	yew (Taxus spp.)

*For CA only

Deciduous Trees and Woody Ornamentals

American elm	eucalyptus	red oak
Siberian elm	holly (Ilex spp.) (max. 3.2 qts./A)	Russian olive
barberry		oleander*
boxelder	honey locust	palm (2 qts./A)*
bush honeysuckle	Oregon grape (Mahonia spp.)	bottle brush*
caragana		carob*
cottonbaster	Pieris spp. (max. 3.2 qts./A)	
dogwood		

*For CA only

8 7 8

SMZ 4L EPA REG. NO. 34704-

Pennant® Tank Mix: SMZ 4L may be tank mixed with Pennant Liquid Herbicide to control weeds in containerized, field and liner* grown ornamentals and commercial landscapes. See the Pennant label for ornamental species, weeds controlled with this tank mixture, and for other information.

*Plants transplanted normally in rows in a nursery or similar area for further growth prior to transplanting to final growing place.

Surflan Tank Mix: On Christmas tree plantings, use this tankmix for preemergence control of weeds listed on the Surflan 75W (or Surflan A.S.) and SMZ 4L labels. Use on field grown conifer species listed on the labels for each herbicide plus grand fir, alpine fir, Englemann spruce, black spruce, Colorado blue spruce, Coulter pine, giant redwood and Veitchi fir.

Broadcast the mixture as a directed spray to the soil surface or as an overtop spray, using 2-4 qts. of SMZ 4L and 2.67 - 5.33 lbs. of Surflan 75W (or 2-4 qts. of Surflan A.S.). Apply in sufficient water per acre to uniformly treat the area. Follow overtop sprays with sprinkler irrigation to move the herbicide from leaf surfaces to the soil. Remove weed growth before application. Mix weed residues, prunings, or trash into the soil, or remove them before treatment. Soils should be in good tilth and free of clods at time of application. Shallow cultivation (1-2 inches) after treatment will not reduce weed control. Observe all precautions and limitations on the SMZ 4L and Surflan labels.

Note: Length of weed control may be reduced when continuous wet soil conditions follow herbicide application.

Precautions: To avoid plant injury, (1) Do not use on seedbeds or on unrooted cuttings. (2) Do not use in greenhouses or other enclosed areas.

TURFGRASSES FOR SOD (FL ONLY)

St. Augustinegrass, Centipedegrass, and Zoysiagrass

Apply 2-4 qts./A, according to soil texture as indicated below.

Muck or peat	4 qts.	Old beds	within 2 days after lifting of sod
		New beds	3-4 days after sprigging or plugging
Sandy soil	2 qts.	Old beds	within 2 days after lifting of sod
		New beds	7-10 days after sprigging or plugging

If weeds regrow, apply an additional 2 qts. on muck or peat, or 1 qt. on sandy soil.

Precautions: To avoid crop injury, (1) Do not apply within 30 days prior to cutting or lifting. (2) Do not apply in combination with surfactants or other spray additives. (3) Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases. (4) On newly sprigged turfgrass, temporary slowing of growth may follow application.

TURFGRASS FOR FAIRWAYS, LAWNS, SOD PRODUCTION*, AND SIMILAR AREAS

*In states other than FL. For use on turfgrass for sod in FL, see Turfgrasses for Sod (FL only) section.

Bermudagrass, Centipedegrass, St. Augustinegrass, and Zoysiagrass

Apply SMZ 4L after September 1 (after October 1 for annual bluegrass) before emergence of winter annual weeds. Fall applications of SMZ 4L will control annual bluegrass, burclover, lawn burweed, common and mouseear chickweed, corn speedwell, henbit, hop clover, spurweed, and parsley-piert. SMZ 4L will also control these weeds soon after emergence. For control of summer annual weeds listed in the General Information section of this label, apply SMZ 4L in late winter before the weeds emerge. Apply in a minimum of 15 gals. of water per acre.

Irrigate with ½ inch of water if rainfall does not occur within 10 days after preemergence treatment.

Where annual bluegrass is the major weed, use 1 qt. of SMZ 4L per acre (22 ml. or 0.75 fl. oz. per 1,000 sq. ft.). Use 1-2 qts./A (22-44 ml. or 0.75-1.5 fl. oz. per 1,000 sq. ft.) for control of other weeds. However, do not exceed 1 qt./A per treatment on newly sprigged turfgrass or on hybrid bermudagrass.

For control of summer annuals which emerge after the initial application, apply an additional 1 qt./A at least 30 days after the initial application, but not after June 1. However, do not make more than 2 applications per year.

Precautions: On newly sprigged turfgrass, hybrid bermudagrass, nondormant bermudagrass, or nondormant zoysiagrass, temporary slowing of growth and yellowing may occur following application. To avoid turf injury, (1) Use only on turfgrass reasonably free of infestations of insects, nematodes, and diseases. (2) Do not use on golf greens. (3) Do not use north of NC (except in the VA Coastal Plains) or on soils with a pH above 7.8. (4) Do not use on muck or alkaline soils. (5) Do not apply over the rooting area of trees or ornamentals not listed on this label. (6) Do not seed or overseed with desirable turfgrass within 4 months before or 6 months after treatment. (7) Do not apply this product to newly seeded grasses until they have overwintered and have a well-developed rhizome system. (8) Do not exceed 2 qts./A within 12 months of seeding grasses.

Note: Do not graze or feed turf clippings to animals, or illegal residues may result.

WARRANTY DISCLAIMER AND NOTICE

THE DIRECTIONS FOR USE OF THIS PRODUCT ARE BELIEVED TO BE ADEQUATE AND SHOULD BE FOLLOWED CAREFULLY. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT DUE TO SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OR ABSENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER.

THE PRODUCTS SOLD TO YOU ARE FURNISHED "AS IS" BY LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER, AND ARE SUBJECT ONLY TO THE MANUFACTURER'S WARRANTIES, IF ANY, WHICH APPEAR ON THE LABELS TO THE PRODUCTS SOLD TO YOU. EXCEPT AS EXPRESSLY PROVIDED HEREIN, LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD OR USE OF THE PRODUCT, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. EXCEPT AS EXPRESSLY STATED HEREIN, LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER MAKES NO WARRANTY OF RESULTS TO BE OBTAINED BY USE OF THE PRODUCT. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND LOVELAND PRODUCTS, INC.'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE LIMITED TO DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT. NO AGENT OR EMPLOYEE OF LOVELAND PRODUCTS, INC. OR SELLER IS AUTHORIZED TO AMEND THE TERMS OF THIS WARRANTY DISCLAIMER OR THE PRODUCT'S LABEL OR TO MAKE A REPRESENTATION OR RECOMMENDATION DIFFERENT FROM OR INCONSISTENT WITH THE LABEL OF THIS PRODUCT.

IN NO EVENT SHALL LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES AND THE BUYER AND USER WAIVE ANY RIGHT THEY MAY HAVE TO SUCH DAMAGES.

AAtrex®, AAtrex® Nine-C, Gramoxone®, Solicam® and Touchdown® are trademarks of Syngenta Group Company
Complex®, trademark of KALO Agricultural Chemicals, Inc.
Roundup® trademark of Monsanto Company
Surflan® trademark of Dow AgroSciences
Unite® trademark of HACO, Inc.
X-77® trademark of Loveland Industries, Inc.

FORMULATED FOR



P.O. BOX 1286, GREELEY, COLORADO 80634-1286