

34704-920

03/03/2006

1/2



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:
34704-920

Date of Issuance:
MAR -3 2006

NOTICE OF PESTICIDE:
 Registration
 Reregistration

Term of Issuance:
Conditional

Name of Pesticide Product:
Quinclorac 75DF

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

**Loveland Products Inc.
PO Box 1286
Greeley, CO 80632**

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided you agree in writing to:

1. Change the EPA Registration Number on the label from 34704- to 34704-920.
2. Add an appropriate EPA Establishment Number to the label.
3. Add "Do not induce vomiting unless told to do so by a poison control center or doctor." to the IF SWALLOWED subsection of the FIRST AID statements.
4. Change the PRECAUTIONARY STATEMENTS to the following: "Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, or clothing. Causes moderate eye injury. Avoid breathing dust or spray mist. May cause allergic skin response."
5. After the heading "Personal Protective Equipment (PPE)" and before "Applicators and other handlers must wear:" add the following statement: "Some materials that are chemical resistant to this product are issued below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart."

Signature of Approving Official

Date:

**James Tompkins, Product Manager (25)
Herbicide Branch, Registration Division (7505C)**

3-3-06

2/12

6. In the PPE section, change “chemical resistant gloves made of any waterproof material, i.e. barrier laminate, butyl rubber or nitrile rubber” to “chemical resistant gloves Category A, such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber \geq 14 mils.”
7. In the Agricultural Use Requirements box, change “waterproof gloves” to “chemical resistant gloves Category A, such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber \geq 14 mils.”
8. On page 3, in the section STORAGE AND DISPOSAL, change “Container Storage” to “Pesticide Storage.” Change the order of the subsections in the STORAGE AND DISPOSAL section to 1) Pesticide Storage 2) Pesticide Disposal 3) Container Disposal.
9. On page 8, under STATE SPECIFIC RESTRICTIONS, add “in” before “Arkansas” but after “Quinclorac 75DF” in the first sentence. Change “on” to “one” in the second sentence. **BOLD** “no aerial application” in the third sentence.
10. In the Warranty Disclaimer and Notice section, add the statement “To the extent allowed by law” after the statements “Except as expressly provided herein,”. Place the statement “To the extent allowed by law” before “Buyer’s or user’s exclusive remedy.....” Also place the statement “To the extent allowed by law” before “In no event shall Loveland Products Inc, the manufacturer.....” in the last paragraph of the Warranty Disclaimer.

The Basic Formulation CSF [dated 1/27/2006] of the product referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act is acceptable. The Basic CSF will be added to your file.

You will submit one (1) copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). A stamped copy of labeling is enclosed for your records.

James Tompkins
James Tompkins
Product Manager (25)
Herbicide Branch
Registration Division (7505C)

3/12

Quinclorac 75DF Herbicide

Active Ingredient

3,7-dichloro-8-quinolinecarboxylic acid..... 75.0%

Inert Ingredients 25.0%

Total 100.0%

EPA Reg. No 34704-

EPA Est. No.

Net Contents: Five 1-pound water-soluble bags

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR -3 2006

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

34704-920



Loveland
PRODUCTS INC.

P.O. Box 1286

Greeley, CO 80632

4/12

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for complete Precautionary Statements, First Aid, Directions For Use, and Conditions of Sale and Warranty.

FIRST AID	
If swallowed	Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-10 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-10 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control or doctor or going for treatment. FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-301-7976.	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin, eyes, or clothing. Avoid breathing dust or spray mist. Causes moderate eye injury. May cause allergic skin response.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, i.e. barrier laminate, butyl rubber or nitrile rubber and shoes plus socks.

Wash thoroughly with soap and water after handling. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water.

Keep and wash PPE separately from other laundry.

Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical where soils are permeable, particularly where water table is shallow, may result in groundwater contamination.

Keep out of lakes, ponds and streams. Do not apply directly to water, areas where surface water is present, or to intertidal areas below the mean high water mark, except as specified on this label for use in rice. Do not contaminate water by cleaning of equipment or disposal of rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** are to be followed. This labeling must be in the user's possession during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Container Storage: Do not store box under wet conditions. Handle box carefully when stored at temperatures less than 50°F to avoid breaking the water-soluble bags.

Container Disposal: Completely empty box into application equipment. Then dispose of packaging in a sanitary landfill or by incineration, or, if allowed by state and local

6/12

authorities, by burning. If burning, stay out of smoke.

Pesticide Disposal: Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.

For help with any spill, leak, fire or exposure involving this material, call day or night
CHEMTREC – 1-800-424-9300.

I. General Information

Quinclorac 75DF can be used for weed control in dry-seeded and water-seeded rice planting and production cultures. Quinclorac 75DF is formulated as dry flowable designed for dilution with water and spraying with common agricultural spray equipment. It is packaged in pre-measured, one pound, water-soluble bags that are added to the mixing vat or directly into the spray tank.

Crop Tolerance

Rice is tolerant to Quinclorac 75DF when used according to label use directions and under typical growing conditions. Adverse weather conditions or high use rate from spray overlap or other sources may contribute to leaf twisting, buggy whipping, or other abnormal growth characteristics. In broadcast or water-seeded rice, seed on the soil surface in direct contact with Quinclorac 75DF is the most sensitive. These symptoms are typically short-lived and rice usually recovers without a significant stand loss or other injury.

Water Management

(Irrigation and Flood Water)

Optimum weed control with Quinclorac 75DF is highly dependent on proper use of irrigation, including effective flush irrigation to maintain moist soil conditions and timely establishment of permanent flood water. Soil applications and residual activity from foliar applications require moist soil conditions for weeds to uptake the herbicide and be controlled. Therefore, keep the soil moist to maintain weed control. If the soil is permitted to dry and weeds emerge, flush irrigate the field to reactivate the residual activity of the herbicide while weeds are small (1" or less). If required, make additional Quinclorac 75DF applications as needed, but limit total usage to 0.67 pound per acre per season. In water-seeded rice plantings and in pin-point flood culture, drain all water from the rice field and ensure seedling rice has at least two leaves before applying Quinclorac 75DF. Rice seedlings without 2 leaves may be injured. Flood water levees should be formed prior to applying Quinclorac 75DF for more consistent weed control. Residual weed control on the levee is dependent on moist soil conditions on the levee. If soil on the levee dries, erratic weed control may result.

If a heavy rain occurs after applying Quinclorac 75DF, drain the excess water from the rice field to avoid possible rice injury.

II. Application Instructions

Quinclorac 75DF may be applied to rice fields to control barnyardgrass, propanil-resistant barnyardgrass, other annual grasses, and certain broadleaf weeds.

Application Equipment:

Whenever possible, spray mixtures should be applied using ground spray equipment. Ensure ground and aircraft spray equipment is properly calibrated and spray coverage is uniform. Always use spray nozzles and other equipment designed to reduce accidental spray drift. Always use drift control products and limit spray applications to periods

when wind and other weather conditions do not favor spray drift beyond the border of the rice field.

Soil Applications: Quinclorac 75DF can be applied to the soil surface before, during, or after planting of dry-seeded rice. When applied to the soil surface and activated by rainfall or irrigation, roots of susceptible grasses and broadleaf weeds uptake the herbicide resulting in commercially acceptable control before weed competition reduces rice productivity. Soil texture and clay content determines the proper use rate for optimum weed control, with heavier soil textures and higher clay content requiring higher use rates as recommended in Table 1.

Foliar Application: Quinclorac 75DF can be applied to the foliage of susceptible grasses and broadleaf weeds in dry-seeded and water-seeded rice. When applied to weed foliage, leaves and stems partially uptake the herbicide. It is essential that rice be flushed after a foliar application to maximize root absorption resulting in commercially acceptable weed control. Additionally, the herbicide reaching the soil surface moves into the soil with rainfall or irrigation providing residual weed control. In general, smaller weeds are more effectively controlled with lower use rates, with larger weeds requiring higher use rates for more complete control. The use rates in Table 1 are recommended for foliar applications to provide commercially acceptable control of susceptible weeds based on weed size or growth stage.

Ground Application

Whenever possible, spray mixtures containing Quinclorac 75DF should be applied using ground spray equipment. Do not make spray applications when wind speed is greater than 10 mph, when air temperatures exceed 90°F, or when environmental conditions exist for temperature inversions.

Application Information:

Preplant/Pre-emergence and Delayed Pre-emergence:

Water Volume: Apply 10-40 gallons of water per broadcast acre.

Spray Pressure: Use 25-40 psi

Postemergence:

Water Volume: Apply 10-20 gallons of water per broadcast acre

Spray Pressure: Use 25-40 psi.

Table 1. Timing and Application Rate Table

Weed Species	Soil Applications (Rate per Acre)			Foliar Applications	
	Light-textured sandy loams	Medium-textured silts, loams, silt loams, sandy clay loams	Heavy-textured such as silty clays, silty clay loams, clay loams, clays, gumbo and buckshot	Small weeds controlled and short-term soil residual	Larger weeds controlled and long-term soil residual
Annual Grasses Barnyardgrass, junglerice, broadleaf signalgrass, large crabgrass	0.33-0.44 pounds	0.50 pounds	0.67 pounds	0.33-0.50 pounds up to 2 inches	0.33 – 0.67 pounds up to 3 inches
Broadleaf Weeds Hemp sebania.	0.33- 0.44 pounds	0.50 pounds	0.67 pounds	0.33 – 0.50 pounds up to	0.50 – 0.67 pounds up to

8/12

jointvetches, morningglories (cypressvine, entireleaf, ivyleaf, palmleaf, purple moonflower, pitted, tall), eclipta				2 leaves	3 leaves
Alligatorweed (partial control)*	n/a	n/a	n/a	0.67	n/a
*Rice must be in at least the 2 leaf stage. For best control, establish permanent flood within 2 days after Quinclorac 75DF application.					

Air Application

If application with ground spray equipment is not possible, application by aircraft is acceptable, provided aerial applicator understands the risks and assumes the liability associated with accidental spray drift from aerial application. Do not make spray applications when wind speed is greater than 8 mph, when air temperatures exceed 90°F, or when environmental conditions exist for temperature inversions.

Application Information:

Water Volume: Apply a minimum of 5 gallons of water per acre

Spray Pressure: Use a maximum of 40 psi.

Cleaning Spray Equipment

All mixing equipment and all spray equipment should be thoroughly cleaned before and after mixing and applying Quinclorac 75DF.

III. Additives

For postemergence applications only, adding 2 pints of crop oil concentrate per acre will improve leaf and stem uptake of the herbicide and enhance weed control.

Drift Control products

Drift control products should always be added to the spray solution to affect spray droplet size and other characteristics, reducing the potential of off-target accidental spray drift.

IV. Mixing Order

- 1) Water: Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
 - 2) Agitation: Maintain constant agitation throughout mixing and application.
 - 3) Products in PVA bags: (such as Quinclorac 75DF) Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
 - 4) Water-dispersible products (such as wettable powders, suspension concentrates, or suspo-emulsions)
 - 5) Water-soluble products
 - 6) Emulsifiable concentrates: If an inductor is used, rinse it thoroughly after the component has been added.
 - 7) Water-soluble additives: If an inductor is used, rinse it thoroughly after the component has been added.
 - 8) Remaining quantity water
- Maintain constant agitation during application. For more information, refer to section V. General Tank Mixing Information.

V. General Tank Mixing Information

While Quinclorac 75DF is effective in controlling a broad spectrum of annual grasses and broadleaf weeds, more effective weed control may be obtained or additional weeds may be controlled by tank mixing Quinclorac 75DF with other herbicides labeled for weed control in rice. The table below describes some weed situations where tank mixing is appropriate. Read and follow all use directions, precautions, and restrictions for each herbicide in the spray mixture. The most restrictive labeling applies to tank mixes.

Table 2. Tank Mixtures

Weed	Tank Mix Information
Cocklebur	Quinclorac 75DF: 0.33 – 0.67 pound Basagran® herbicide: 1.5 – 2.0 pints
Dayflower	Quinclorac 75DF: 0.33-0.67 pound Basagran herbicide: 1.5-2.0 pints
Hemp Sesbania	a) Quinclorac 75DF: 0.33-0.67 pound Blazer® herbicide: 0.5-1.0 pint ¹ b) Quinclorac 75DF: 0.33-0.67 pound Command®3ME: 0.8-1.6 pints
Sprangletop	a) Quinclorac 75DF: 0.33-0.67 pound Bolero®8 EC herbicide: 0.5-1.0 pint ² b) Quinclorac 75DF: 0.33-0.67 pound Prowl® 3.3 EC herbicide: 2.4 pints ³ c) Quinclorac 75DF: 0.33-0.67 pound Command®3ME: 0.8-1.6 pints
Yellow Nutsedge	Quinclorac 75DF: 0.33-0.67 pound Basagran herbicide: 1.5-2.0 pints
Morningglory	Quinclorac 75DF: 0.33 – 0.67 pound Command®3ME: 0.8 – 1.6 pints
Heavy infestations of broadleaf weeds	Quinclorac 75DF: 0.33 – 0.67 pound Storm® herbicide: 1.5 pints
For weeds and grasses not controlled by Quinclorac 75DF	Quinclorac 75DF: 0.33 – 0.67 pound Propanil: 2-4 pounds a.i.
¹ Apply tank mix after rice has reached the 3 leaf stage ² Apply tank mix to the soil surface 1-5 days before rice emergence. ³ Apply this tank mix to soil surface after planting, before rice emergence, and before sprangletop emergence.	

VI. General Restrictions and Limitations – Rice

Maximum seasonal use rate: Do not apply more than 0.67 pounds of Quinclorac 75DF herbicide per acre, per season.

Preharvest Interval (PHI): Do not apply Quinclorac 75DF within 40 days of harvest. Do not apply Quinclorac 75DF to rice that is heading.

10/12

State Specific Restrictions: Because there are additional state restrictions in Arkansas, contact the Arkansas Plant Board or a representative for specific instructions about applying Quinclorac 75DF Arkansas.

In Arkansas Quinclorac 75DF (Quinaclorac) must not be applied in an area from one mile west of Highway #1 to one mile east of Highway #163 from the Craighead – Poinsett County line to the Cross-Poinsett County line. Furthermore, no aerial application is allowed in the area of Poinsett County one mile west of Highway #1 to two miles west of Highway #1 and one mile east of Highway #163 to Ditch #10, from the Craighead – Poinsett County line to the Cross – Poinsett county line.

Crop Rotation Restrictions: Do not plant any crop other than rice for a period of 309 days following application.

- Eggplants and tobacco should not be planted within 12 months on the fields treated with Quinclorac 75DF.
- Tomatoes and carrots should not be planted within 24 months on fields treated with Quinclorac 75DF.

In case of crop failure, only rice may be immediately replanted.

Soil Restrictions:

- Do not use Quinclorac 75DF on precision-cut fields until the second rice crop as injury can occur.
- Do not use Quinclorac 75DF on sand and loamy sand soils.
- Do not apply to rice fields with a history or poor water-holding capacity (porous subsoil), as erratic weed control may result.
- Do not apply Quinclorac 75DF on any rice soil that does not have an impermeable hard pan to provide good water holding capacity.

Drift Concerns: Do not allow Quinclorac 75DF to drift outside of the intended target areas.

- Ground application: Do not apply when wind speed is greater than 10 mph.
- Aerial application: Do not apply when wind speed is greater than 8 mph.

Temperature Inversions: Do not apply Quinclorac 75DF when air temperatures exceed 90°F.

Do not use rice straw or processing byproducts (such as chaff, hulls, etc.) as soil amendments or mulch for high-value crops such as bedding stock, vegetable transplants, or ornamental and fruit trees.

Do not use treated rice fields for the aquaculture of edible fish and crustacea (crayfish).

Do not use water from rice cultivation after a Quinclorac 75DF application to irrigate any crop other than rice.

Do not use in California or Florida.

Quinclorac 75DF cannot be used to formulate or reformulate any other pesticide product.

Do not apply this product through any type of irrigation system.

11/12

Table 3. Restrictions and Limitations

Crop	Minimum Time from Application to Harvest (PHI)	Maximum Rate Per Acre per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application
Rice	40 days	0.67 pound	0.67 pound	n/a	Yes

Crops
This product can be used on the following crops
Rice
Look inside for complete Restrictions and Limitations and Application Instructions

Weeds listed in this label:	
Common Name	Scientific Name
Alligatorweed	<i>Alternanthera philoxeroides</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Cocklebur	<i>Xanthium strumarium</i>
Crabgrass, Large	<i>Digitaria sanguinalis</i>
Dayflower Species	<i>Commelina</i> spp.
Eclipta	<i>Eclipta alba</i>
Jointvetch Species	
Northern	<i>Aeschynomene virginica</i>
Indian	<i>Aeschynomene indica</i>
Junglerice	<i>Echinochloa colonum</i>
Morningglory species	
Tall (common)	<i>Ipomoea purpurea</i>
Cypressvine	<i>Ipomoea quamoclit</i>
Entireleaf	<i>Ipomoea hederacea integruscula</i>
Ivyleaf	<i>Ipomoea hederacea</i>
Palmleaf	<i>Ipomoea wrightii</i>
Pitted	<i>Ipomoea lacunosa</i>
Purple Moonflower	<i>Ipomoea muricata</i>
Sesbania, Hemp	<i>Sesbania exaltata</i>
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>
Sprangletop	<i>Leptochloa</i> spp.
Yellow Nutsedge	<i>Cyperus esculentus</i>

WARRANTY DISCLAIMER AND NOTICE

THE DIRECTIONS FOR USE OF THIS PRODUCT ARE BELIEVED TO BE ADEQUATE AND SHOULD BE FOLLOWED CAREFULLY. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT DUE TO SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OR ABSENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH

12/12

ARE BEYOND THE CONTROL OF LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER.

THE PRODUCTS SOLD TO YOU ARE FURNISHED "AS IS" BY LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER, AND ARE SUBJECT ONLY TO THE MANUFACTURER'S WARRANTIES, IF ANY, WHICH APPEAR ON THE LABELS TO THE PRODUCTS SOLD TO YOU. EXCEPT AS EXPRESSLY PROVIDED HEREIN, LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD OR USE OF THE PRODUCT, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. EXCEPT AS EXPRESSLY STATED HEREIN, LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER MAKES NO WARRANTY OF RESULTS TO BE OBTAINED BY USE OF THE PRODUCT. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND LOVELAND PRODUCTS, INC.'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE LIMITED TO DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT. NO AGENT OR EMPLOYEE OF LOVELAND PRODUCTS, INC. OR SELLER IS AUTHORIZED TO AMEND THE TERMS OF THIS WARRANTY DISCLAIMER OR THE PRODUCT'S LABEL OR TO MAKE A REPRESENTATION OR RECOMMENDATION DIFFERENT FROM OR INCONSISTENT WITH THE LABEL OF THIS PRODUCT.

IN NO EVENT SHALL LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES AND THE BUYER AND USER WAIVE ANY RIGHT THEY MAY HAVE TO SUCH DAMAGES TO THE EXTENT ALLOWED BY LAW.