



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Washington, D.C. 20460

September 10, 2008

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

Kristie Raymond  
Registration Specialist  
Loveland Products, Inc.  
7251 W 4<sup>th</sup> Street  
P.O. Box 1286  
Greeley, CO 80632-1286

Subject: Product Name: DM Fosphite Fungicide  
EPA Reg. No: 34704-924  
Application for label Notification Dated June 17, 2008 to Revise the Container  
Disposal Language per PR Notice 2007-4

Dear Ms. Raymond:

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under 98-10 dated above. A preliminary screen of this request has been conducted for its applicability under PR Notice 98-10 and it has been determined that the action(s) requested falls within the scope of PR Notice 98-10. Our records have been duly noted, and the label submitted with this application has been stamped "Notification, received and reviewed" and will be placed accordingly in our records.

Questions concerning this action should be directed to Ms. Diana Hudson at (703) 308-8713 or email at [hudson.diana@epa.gov](mailto:hudson.diana@epa.gov).

Sincerely,

*Linda Hollis*

Linda Hollis, Chief  
Biochemical Pesticides Branch  
Biopesticides and Pollution Prevention  
Division (7511P)



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number 34704-924	2. EPA Product Manager Linda Roberts	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) DM Fosphite Fungicide	PM# 91	
5. Name and Address of Applicant (Include ZIP Code) Loveland Products, Inc. P.O. Box 1286 Greeley, Colorado 80632-1286 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	<b>NOTIFICATION</b> Date Reviewed: 9-10-08 Reviewed By: D. Hudson
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 15.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

### Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted				<input checked="" type="checkbox"/> Glass	<input type="checkbox"/> Paper
	If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container 1 gal	5. Location of Label Directions <input type="checkbox"/> On label booklet			
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input checked="" type="checkbox"/> Other self adhesive, pressure sensitive booklet		

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Kristie Raymond kristie.raymond@uap.com	Title Registration Specialist	Telephone No. (Include Area Code) 970-347-1347
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Kristie Raymond</i>	3. Title Registration Specialist	
4. Typed Name Kristie Raymond	5. Date June 17, 2008	




---

Performance	Quality	Value
-------------	---------	-------

---

June 17, 2008

Document Processing Desk (NOTIF)  
 Office of Pesticide Programs (7504P)  
 Linda Roberts  
 U.S. Environmental Protection Agency  
 One Potomac Yard, 4<sup>th</sup> Floor, Room S-4900  
 2777 South Crystal Drive  
 Arlington, VA 22202

Subject: DM Fosphite Fungicide (EPA Reg. No. 34704-924); Notification

Enclosed, is the Loveland Products, Inc. (LPI) submission for a Notification of DM Fosphite Fungicide (EPA Reg. No. 34704-924) per PR Notice 2007-4.

Please find the following enclosed:


1. Application for Registration (EPA Form 8570-1);
2. 1 copy of the DM Fosphite Fungicide label


If you have any questions, please feel free to contact me at 970-347-1347 or [kristie.raymond@uap.com](mailto:kristie.raymond@uap.com).

Sincerely,

Kristie Raymond  
 Registration Specialist

Enclosures





# DM FOSPHITE FUNGICIDE

## NOTIFICATION

Date Reviewed: 9-10-08

Reviewed By: D. Hudson

**A FUNGICIDE FOR THE CONTROL OF DOWNY MILDEW, PHYTOPHTHORA, PHYTHIUM AND VARIOUS OTHER DISEASES ON AGRICULTURAL AND GREENHOUSE CROPS, INDOOR AND OUTDOOR ORNAMENTALS, BEDDING PLANTS AND FORESTRY APPLICATIONS AND DOMESTIC TREES INCLUDING CONIFERS, PINES, OAKS AND TURF**

<b>ACTIVE INGREDIENTS:</b>	
Mono- and dipotassium salts of Phosphorous Acid*	53%
<b>OTHER INGREDIENTS:</b>	47%
<b>TOTAL</b>	<b>100%</b>

\*Contains 6.22 lbs./gallon of active ingredients, mono- and dipotassium salts of Phosphorous Acid. Equivalent to 3.90 lbs. Phosphorous Acid/gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION

See inside label booklet for First Aid and additional precautionary statements.

### FIRST AID STATEMENTS

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
--------------------	---

Have the container or label with you when calling a poison control center or doctor, or going for treatment. **FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1 800 301-7976.**

**EPA REG. NO. 34704-924  
EPA EST. NO. 34704-MS-001  
NET CONTENTS 1 GAL. (3.78 L)**

IHT 022608 V2D 06/08

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water

when disposing of equipment washwaters or rinsate.

### PHYSICAL AND CHEMICAL HAZARDS

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other person, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Workers Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants,
- Socks, shoes,
- Chemical resistant gloves
- Protective eyewear.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

### GENERAL APPLICATION INSTRUCTIONS

Apply DM Fosphite by various application methods, including foliar sprays, trunk injection, soil drench, soil incorporation, and bare root dip. For foliar sprays, apply DM Fosphite with sufficient water for adequate coverage of foliage, according to crop and growth stage. To insure good coverage, spray to wet.

**GENERAL APPLICATION INSTRUCTIONS (TRUNK INJECTION ONLY):** DM Fosphite is a systemic fungicide for the prevention and treatment of downy mildew, Sudden Oak Death (*Phytophthora ramorum*), root rots (*Phytophthora* spp.), pythium and various other disease of agricultural trees, indoor and outdoor ornamentals, forestry applications including domestic trees, conifers, pines and oaks. This product may be applied up to the day of harvest.

### MIXING INSTRUCTIONS

1. Fill the spray tank with 1/2 of the required volume of water.
2. Add DM Fosphite slowly to the tank and agitate by hydraulic or mechanical means.
3. Continue to fill the tank with water to the desired volume while agitating.
4. Continue agitation when applying.

For 1 gallon (3.785 liters) container sizes: this package can treat up to 34,848 sq. ft. at the maximum treatment rate.

For 2 1/2 gallon (9.463 liters) container sizes: this package can treat up to 2 acres at the maximum treatment rate.

For 5 gallon (18.925 liters) container sizes: this package can treat up to 4 acres at the maximum treatment rate.

### LOW VOLUME AND ULTRA-LOW VOLUME APPLICATIONS

Label rates are written for conventional application equipment. For LV/ULV equipment, use per acre rate of DM Fosphite and adjust amount of water to equipment needs. Always conduct a phytotoxicity test on a small area before applying to a large area in order to assess any potential risk to plants.

# DM FOSPHITE EPA REG. NO. 34704-924

## COMPATIBILITY

Mixing DM Fosphite with certain surfactants, foliar fertilizers or other pesticides may cause crop injury. DM Fosphite is a slightly acidic buffer solution. Avoid mixing DM Fosphite with strongly acidic or alkaline materials. Do not tank mix without first testing the mixture's compatibility nor apply it without assessing its safety to the crop (phytotoxicity). It is not possible to test all material combinations and environmental tank mix combinations. Always conduct a test with the intended tank mix on a small area prior to any large scale applications. To determine if a combination is phytotoxic to a specific crop, spray a small area of foliage and fruit, and then evaluate 3-7 days later for visual symptoms.

Test the compatibility of spray adjuvants (i.e. stickers, spreaders, wetting agents) with DM Fosphite before use. Do not use acidifying type compatibility agents.

To determine the compatibility of DM Fosphite with other products, use a jar compatibility test. Add the correct proportions of each product and the appropriate quantity of water to clean container, thoroughly mix, then let stand for 3-5 minutes. If the mixture remains in solution or can be remixed readily, the products are considered compatible.

**COMPATIBILITY (TRUNK INJECTION ONLY):** Use only DM Fosphite alone when used as a trunk injection. To avoid undesirable phytotoxicity, do not apply DM Fosphite to trees treated with other pesticides or copper based compounds at intervals less than 20 days, unless instructed to do so by your consultant.

## CHEMIGATION

### APPLICATIONS THROUGH SPRINKLER AND DRIP IRRIGATION SYSTEMS

Apply this product through microjet, drip, solid set and center pivot sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

**PREPARATION OF INJECTION EQUIPMENT:** Remove scale, pesticide residues and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

**APPLICATION INSTRUCTIONS:** Fill tank with 1/2 to 3/4 of desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of DM Fosphite, then the remaining volume of water. Set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the solution of DM Fosphite into the irrigation water line so as to deliver the desired rate per acre. Inject the DM Fosphite solution with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. For other questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

**NOTE:** For Microjet and Drip Irrigation Systems: Avoid further irrigation of the treated area for 24 to 48 hours after treatment with DM Fosphite has been completed. For Solid Set and Center Pivot Irrigation Systems: Avoid further irrigation of the DM Fosphite treated area until after the foliage has completely dried. This will help to prevent washing the chemical off the crop.

### GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER AND DRIP IRRIGATION SYSTEMS

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally dosed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

**Maintain continuous agitation in mix tank during mixing and application to assure uniformity solution.** Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution, or when lines containing the product must be dismantled or drained. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of chemigation water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut down and adjust the system as needed.

Do not connect an irrigation system (including greenhouse system) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

## GENERAL INSTRUCTIONS FOR APPLICATIONS FOOD CROPS

DM Fosphite fungicide is used for control of Downy Mildew, Powdery Mildew, Erwinia amylovora, Pythium, Fusarium, Rhizoctonia and Phytophthora disease (such as purple blotch, late blight, blue mold, soft rot, root rot, brown rot, foot rot, crown/collar rot, canker, fruit/leaf rot, fire blight, heart rot, pink rot, pythium leak, red stele, leather rot, etc.) of

agronomic crops. (See specific groupings for additional diseases.) This product may be applied up to the day of harvest.

DM Fosphite is applied to plants including:

**Berry Crop Grouping:** Blackberry (including bingberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Grapes (all types), Himalayaberry, hullberry, Lavacaberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these), Blueberry, Cane Berries, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Raspberry (black and red) and Strawberry.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Xanthomonas campestris Anthracnose	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in a minimum of 100 liters of water/hectare)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	
	Root Dip	2 quarts/100 gallons of water, 2/3 fluid ounce/gallon of water; (5 liters/1000 liters of water)	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
Downy Mildew Powdery Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in a minimum of 100 liters of water/hectare)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	

\* Check with your local Extension Agent if you are unsure these conditions exist

**Brassica Crop Grouping:** Broccoli (all types), Brussels sprouts, Cabbage (all types), Cauliflower, Cavalo broccoli, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach and Rape greens.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

**DM FOSPHITE**  
EPA REG. NO. 34704-924

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Xanthomonas campestris Anthracnose	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in minimum of 100 liters of water/hectare)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	or		
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	
	Root Dip	1 quart/100 gallons of water, 1/3 fluid ounce/ gallon of water; (2.5 liters/ 1000 liters of water)	
	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days
Downy Mildew Powdery Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in minimum of 100 liters of water/hectare)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	or		
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Cereal Grains, Forage and Fiber Crop Grouping:** Alfalfa, Barley, Buckwheat, Canola, Clover (all types), Corn (all types), Cotton, Hay, Kudzu, Lespedeza, Lupin, Millet, Oats, Popcorn, Rice, Rye, Safflower, Saintoin, Sorghum, Sunflower, Teosinte, Trefoil, Triticale, Vetch, Wheat and Wild rice.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Xanthomonas campestris Anthracnose	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in minimum of 100 liters of water/hectare)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	or		
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	

6/13

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in minimum of 100 liters of water/hectare)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	or		
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Citrus Crop Grouping (Bearing and Non-Bearing) including Avocado:** Avocado, Calamondin, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin, Orange (all types), Pummelo and Satsuma madarin.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Xanthomonas campestris	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre; (5 to 7 liters in a minimum of 1000 liters of water/hectare)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	or		
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	
	Root Dip	2 quarts/100 gallons of water, 2/3 fluid ounces/gallon of water; (5 liters/1000 liters of water)	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	Trunk Injection	3 teaspoons per 100 square yard of canopy width or 2 inch of trunk diameter at breast height (DBH) (15 milliliters/linear meter of canopy width or 5 centimeters DBH.) Inject directly in to tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.

**DM FOSPHITE**  
**EPA REG. NO. 34704-924**

Disease	Application Method	Rate	Application Program
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre; (5 to 7 liters in a minimum of 1000 liters of water/hectare)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	or  Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters/hectare)	

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Cucurbit Crop Grouping:** Chayote, Chinese waxgourd, Citron melon, Cucumber (all types), Gherkin, Gourd, Muskmelon, Pumpkin, Squash (summer and winter) and Watermelon.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Anthracnose	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in a minimum of 100 liters of water/hectare)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	or  Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	
	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	

Disease	Application Method	Rate	Application Program
Downy Mildew Powdery Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in a minimum of 100 liters of water/hectare)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	or  Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare)	

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Herbs and Spices Crop Grouping:** Allspice, Angelica, Anise, Annatto, Balm, Basil, Borage, Bumel, Camomile, Caper buds, Caraway, Cardamom, Cassia bark and buds, Catnip, Celery seed, Chervil, Chive, Cinnamon, Clary, Clove, Cocoa, Coriander, Costmary, Cilantro, Cumin, Curry, Dill, Fennel, Fenugreek, Grains of paradise, Hops, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage, Mace, Marigold, Marjoram, Mint, Mustard, Nasturtium, Nutmeg, Parsley, Pennyroyal, Pepper, Poppy seed, Rosemary, Rue, Saffron, Sage, Savory, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff and Wormwood.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in a minimum of 100 liters of water/hectare)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	or  Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	
	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	
Downy Mildew Powdery Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in a minimum of 100 liters of water/hectare)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	or  Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare)	

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Legume Vegetables Crop Grouping:** Bean (all types), Broad bean (fava bean), Chick pea (garbanzo bean), Guar, Jackbean, Lablab bean, Lentil, Pea (all types), Peanut, Pigeon pea, Soybean and Sword bean.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

**DM FOSPHITE  
EPA REG. NO. 34704-924**

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Early and Late Leaf Spot on Peanut ( <i>Cercospora arachidicola</i> and <i>Cercosporidium personatum</i> ) Xanthomonas campestris Anthracnose	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in a minimum of 100 liters of water/hectare)	<b>Disease Prevention Program:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	
	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	
Downy Mildew Powdery Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in a minimum of 100 liters of water/hectare)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program for Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Pome Fruits Crop Grouping (Bearing and Non-Bearing) including Papaya:** Apple, Crabapple, Loquat, Mayhaw, Papaya, Pear, and Quince.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Fire Blight ( <i>Erwinia amylovora</i> ) Xanthomonas campestris	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre; (5 to 7 liters in a minimum of 1000 liters of water/hectare)	<b>Disease Prevention Program:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	or	1 to 3 quarts in a minimum of 100 gallons of water/acre; (5 to 7 liters in a minimum of 1000 liters of water/hectare)	
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters/hectare.)	
	Root Dip	2 quarts/100 gallon of water, 2/3 fluid ounces/gallon of water; (5 liters/1000 liters of water)	

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Fire Blight ( <i>Erwinia amylovora</i> )	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	Trunk Injection	3 teaspoons/linear yard of canopy width or 2 inch of trunk diameter at breast height (DBH) (15 milliliter/linear meter of canopy width or 5 centimeters DBH.) Inject directly in to tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
Downy Mildew Powdery Mildew	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre; (5 to 7 liters in a minimum of 1000 liters of water/ hectare)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters/hectare)	

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Stone Fruits Crop Grouping including Olives:** Apricot, Cherry, Nectarine, Olive, Peach, Plum, Plumcot and Prune.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Xanthomonas campestris	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/acre; (5 to 7 liters in a minimum of 1000 liters of water/hectare)	<b>Disease Prevention Program:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	or	1 to 3 quarts in a minimum of 100 gallons of water/acre; (5 to 7 liters in a minimum of 1000 liters of water/hectare)	
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre; (5 to 7 liters in a minimum of 75 liters/hectare.)	
	Root Dip	2 quarts/100 gallon of water, 2/3 fluid ounces/gallon of water; (5 liters/1000 liters of water)	
	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.



**DM FOSPHITE  
EPA REG. NO. 34704-924**

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Trunk Injection	3 teaspoons/linear yard of canopy width or 2 inch of trunk diameter at breast height (DBH) (15 milliliter/linear meter of canopy width or 5 centimeters DBH.) Inject directly in to tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
Downy Mildew Powdery Mildew	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/ acre; (5 to 7 liters in a minimum of 1000 liters of water/hectare)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	or	1 to 3 quarts in a minimum of 10 gallons of water/ acre; (5 to 7 liters in a minimum of 75 liters/hectare)	
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/ acre; (5 to 7 liters in a minimum of 75 liters/hectare)	

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Tree Nuts Crop Grouping including Vines:** Almond, Beech nut, Betelnut Palm, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Grapes (all types), Hickory nut, Kiwi, Macadamia nut, Pecan, Pistachio and Walnut.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Xanthomonas campestris	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/ acre; (5 to 7 liters in a minimum of 1000 liters of water/hectare)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	or	1 to 3 quarts in a minimum of 10 gallons of water/ acre; (5 to 7 liters in a minimum of 75 liters/hectare)	
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/ acre; (5 to 7 liters in a minimum of 75 liters/hectare)	
	Root Dip	2 quarts/100 gallon of water, 2/3 fluid ounces/gallon of water; (5 liters/1000 liters of water)	Apply as pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping.
	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
Trunk Injection	3 teaspoons/ linear yard of canopy width or 2 inch of trunk diameter at breast height (DBH) (15 milliliter/ linear meter of canopy width or 5 centimeters DBH.) Inject directly into tree. <b>Betelnut Palm:</b> ¼ to 1 fluid ounces per tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.	

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Xanthomonas campestris <i>cont'd.</i>	Soil Drench	1- 2 gallons/100 gallons of water or 1.255 to 2.5 fluid ounces per gallon of water (10 - 20 liters/1000 liters of water)	Apply 5 - 10 gallons of diluted solution per tree. Adjust volume as needed to thoroughly drench area around trunk. Apply at 3 - 4 week intervals and repeat as needed. Do not apply DM Fosphite at intervals less than 3 days.
Downy Mildew Powdery Mildew	Foliar	1 to 3 quarts in a minimum of 100 gallons of water/ acre; (5 to 7 liters in a minimum of 1000 liters of water/ hectare)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	or	1 to 3 quarts in a minimum of 10 gallons of water/ acre; (5 to 7 liters in a minimum of 75 liters/hectare.)	
	Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/ acre; (5 to 7 liters in a minimum of 75 liters/hectare.)	

\* Check with your local Extension Agent if you are unsure these conditions exist.

**Vegetables Crop Grouping Including Miscellaneous Food Crops:** (Leafy, Root and Tuber) Amaranth (Chinese spinach, tampala, etc.) Arracacha, Arugula, Arrowroot, Artichoke (all types), Asparagus, Beet (all types), Burdock, Canna, Cardoon, Carrot, Cassava, Celeric, Celery, Chayote, Chenil (turnip-rooted), Chicory, Chilies, Chrysanthemum (edible), Chufa, Cinchoca, Coconut, Coffee, Corn (all types), Cress (all types), Dandelion, Dasheen, Dock, Eggplant, Endive, Fennel, Garlic, Ginger, Ginseng, Horseradish, Leren, Lettuce (all types), Okra, Onion, Orach, Parsley, Parsnip, Peppers (all types), Pineapple, Purslane, Potato (all types), Radicchio, Radish (all types), Rhubarb, Rutabaga, Salisify (all types), Skiret, Spinach (all types), Sugarbeet, Sugarcane, Sweet Potato, Swiss chard, Tanier, Tobacco, Tomato, Turmeric, Turnip, Yam (all types).

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Xanthomonas campestris Anthracnose	Foliar	1 to 4 quarts in a minimum of 20 gallons of water/ acre; (5 to 7 liters in minimum of 100 liters of water/hectare)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days.
	or	1 to 4 quarts in a minimum of 10 gallons of water/ acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	
	Aerial Application	1 to 4 quarts in a minimum of 10 gallons of water/ acre; (5 to 7 liters in a minimum of 75 liters of water/hectare.)	
	Root Dip	2 quarts/100 gallon of water, 2/3 fluid ounces/gallon of water; (5 liters/1000 liters of water)	Apply as pre-plant dip immediately prior to planting. (100 gallons is intended to treat the number of slips required to plant one acre).
	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water; (7-9 liters in at least 1000 liters of water) Drip: 2-3 quarts in at least 100 gallons of water; (5-7 liters in at least 1000 liters of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.

**DM FOSPHITE**  
EPA REG. NO. 34704-924

Disease	Application Method	Rate	Application Program
Downy Mildew Powdery Mildew	Foliar  or  Aerial Application	1 to 4 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in minimum of 100 liters of water/hectare)  1 to 4 quarts in a minimum of 10 gallons of water/acre (5 to 7 liters in a minimum of 75 liters of water/hectare.)	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
Pink Rot ( <i>Phytophthora erythroseptica</i> ) Late Blight ( <i>Phytophthora infestans</i> )	Foliar or Aerial Application	3 to 5 quarts in a minimum of 20 gallons of water/acre; (5 to 7 liters in minimum of 100 liters of water/hectare) *Note: 1 gallon container will not be sufficient for treating pink rot and late blight at the highest given application rate. An additional 1 gallon container or a larger container size will be necessary to achieve the maximum treatment rate.	Apply at 2-3 week intervals. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
Late Blight ( <i>Phytophthora infestans</i> ) Pink Rot ( <i>Phytophthora erythroseptica</i> )	Post-harvest treatment of potatoes: Spray, rinse or inject into humidification water prior to or during storage.	Dilute DM Fosphite fungicide at 1:20 ratio with application water. 6.4-12.8 fluid ounces/0.5 gallon of water/ton of tubers (95-190 milliliter/liter of water)	Inject concentrate into water used in humidification or process water used for post-harvest storage, rinses and associated tanks, flumes and lines.

\* Check with your local Extension Agent if you are unsure these conditions exist.

**ORNAMENTALS**

DM Fosphite fungicide is used for control of Pythium, Downy Mildew, Powdery Mildew, Phytophthora and various other diseases of ornamentals grown in commercial nurseries and greenhouses, golf courses, parks and commercial landscapes. DM Fosphite may be applied to all ornamental and landscape plants including (but not limited to): Aglaonema, Aphelandra, Arborvitae, Ash, Azaleas, Bougainvilleas, Boxwood, Bushes, Cattleya skinneri, Cedars, Ceanothus, Cotoneaster, Cissus, Diffenbachia, Dogwood, Ferns, Ficus, Flowering Trees, Hibiscus, Hard Wood Trees, Holly, Home Lawns, Juniper, Magnolia Trees, Maple, Pines, Oaks, Palms, Peperomia, Photinia, Pittosporum, Philodendron, Pieris, Polthos, Red woods, Rhododendron, Roses, Schefflera, Sedum, Sempervivum, Shrubs, Spatiphyllum, Sycamore, Taxus media, Zygocactus.

Apply DM Fosphite prior to disease development in conjunction with good cultural practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or under heat or moisture stress. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Fusarium Pythium Rhizoctonia Fire Blight ( <i>Erwinia amylovora</i> ) Bacterial Blight ( <i>Xanthomonas campestris</i> ) Geranium Will ( <i>Ralstonia</i> spp.) Anthracnose	Foliar Spray	1 to 2 quarts/100 gallons of water or 1/3 fluid ounce to 2/3 fluid ounce per gallon of water; (2.5 to 5 liters/1000 liters of water)	<b>Disease Prevention Program*</b> : Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program</b> : Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply DM Fosphite at intervals less than 3 days. Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply DM Fosphite at intervals less than 3 days.
	Soil Drench	12-24 fluid ounce/100 gallon of water or 1/8 to 1/4 fluid ounce per gallon of water; (1 to 2 liters/1000 liters of water)	Apply 1-2 pts. of diluted solution per sq. foot. Adjust volume as needed to thoroughly wet media. Apply at 2-3 week intervals and repeat as needed. Do not apply DM Fosphite at intervals less than 3 days.
Phytophthora Fusarium Pythium Rhizoctonia Fire Blight ( <i>Erwinia amylovora</i> ) Bacterial Blight ( <i>Xanthomonas campestris</i> ) Geranium Will ( <i>Ralstonia</i> spp.)	Soil Incorporation	1 to 2 pints/cubic yard of soil mix (300-600 milliliters/cubic meter)	Mix 1 to 2 pints of DM Fosphite with enough water to wet one cubic yard of soil; mix immediately before potting. If conditions are favorable for disease development, it is recommended to make additional applications as a foliar spray or soil drench.
	Irrigation	Overhead: 1-2 quarts/100 gallons of water; (2.5 - 5 liters/1000 liters of water) Drip: 1-2 quarts/100 gallons of water; (2.5 - 5 liters/1000 liters of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	Trunk Injection	3 teaspoons/linear yard of canopy width or 2 inch of trunk diameter at breast height (DBH) (15 milliliters/linear meter of canopy width or 5 centimeters DBH.) Inject directly into tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications. Repeat 2-4 times a year until control is reached.
	Root Dip	2 quarts/100 gallons of water or 2/3 fluid ounce/gallon of water; (5 liters/1000 liters of water)	Dip transplants into diluted solution of DM Fosphite. Thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.
Downy Mildew Powdery Mildew	Foliar Spray	1 to 2 quarts/100 gallons of water or 1/3 fluid ounce to 2/3 fluid ounce/gallon of water; (2.5 to 5 liters/1000 liters of water)	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Follow <b>Disease Prevention Program</b> or <b>Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.

\* Check with your local Extension Agent if you are unsure these conditions exist.

**BEDDING PLANTS**

DM Fosphite fungicide is used for control of Downy Mildew, Powdery Mildew, Pythium, Phytophthora and various other diseases of bedding plants grown in commercial nurseries and greenhouses, golf courses, parks and commercial landscapes. DM Fosphite may be applied to all bedding plants including: Ageratum, Ageria, Ivy, Alyssum, Anthurium, Artemisia, Aster, Baby's Breath, Begonia, Caladium, Carnation, Chrysanthemum, Coleus, Columbine, Daisy, Delphinium, Easter Lily, English Ivy, Foxglove, Gaillardia, Geranium, Gloxinia, Impatiens, Marigold, Pansy, Petunia, Phlox, Pinks, Poinsettia, Primrose, Prostrate Rosemary, Salvia, Snapdragon, Verbena, Vinca, and Zinnia.

**DM FOSPHITE**  
EPA REG. NO. 34704-924

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia Fire Blight ( <i>Erwinia amylovora</i> ) Anthracnose	Foliar Spray	1 to 2 quarts/100 gallons water or 1/3 fluid ounce to 2/3 fluid ounce/gallon of water; (2.5 to 5 liters/1000 liters water)	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply DM Fosphite at intervals less than 3 days.
	Root Dip	2 quarts/100 gallons of water or 2/3 fluid ounce/gallon of water; (5 liters/1000 liters of water)	Dip transplants into diluted solution of DM Fosphite. Thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.
	Soil Drench	12-24 fluid ounce/100 gallon of water or 1/8 to 1/4 fluid ounce per gallon of water; (1 to 2 liters/1000 liters of water)	Apply 1-2 pts. of diluted solution per sq. foot. Adjust volume as needed to thoroughly wet media. Apply at 2-3 week intervals and repeat as needed. Do not apply DM Fosphite at intervals less than 3 days
	Irrigation	Overhead: 1-2 quarts./100 gallons of water; (2.5 - 5 liters/1000 liters of water) Drip: 1-2 quarts/100 gallons of water; (2.5 - 5 liters/1000 liters of water)	Apply with normal irrigation schedule. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.
	Foliar Spray	1 to 2 quarts/100 gallons water; (1/3 fluid ounce to 2/3 fluid ounce/gallon of water) (2.5 to 5 liters/1000 liters of water)	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply more than 6 times per crop cycle or year. Follow <b>Disease Prevention Program or Disease Control Program</b> as stated above. Do not apply DM Fosphite at intervals less than 3 days.

\*Check with your local Extension Agent if you are unsure these conditions exist.

**CONIFERS AND TREES IN COMMERCIAL NURSERIES, PLANTATIONS AND FORESTS, LANDSCAPES AND PARKS (INCLUDING CHRISTMAS TREES, OAKS and PINES)**

DM Fosphite applied in conjunction with good cultural management practices will control Phytophthora root rot of conifers including (but not limited to): Douglas Fir, spruce and pines also other Phytophthora disease of Oaks (Sudden Oak Death) and forest trees. Use the higher rates when disease pressure is severe. Do not apply to trees that are under heat or moisture stress.

Disease	Application Method	Rate	Application Program
Phytophthora	Foliar Spray	1 to 2 quarts/100 gallon of water/acre; (2.5 to 5 liters/1000 liters water/hectare)  or  1/3 fluid ounce to 2/3 fluid ounce/gallon of water. Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed.	<b>Disease Prevention Program*:</b> Apply lower rate at 2-4 week intervals after plants become established. Do not apply DM Fosphite at intervals less than 3 days. <b>Disease Control Program:</b> Apply higher rate at 2-3 week intervals until control is reached. Do not apply DM Fosphite at intervals less than 3 days.
	Soil Drench	12-24 fluid ounce/100 gallons of water or 1/8 to 1/4 fluid ounce/gallon of water; (1 to 2 liters/1000 liters of water)	Apply 1-2 pts. of diluted solution per sq. foot. Adjust volume as needed to thoroughly wet media. Apply at 2-3 week intervals and repeat as needed. Do not apply DM Fosphite at intervals less than 3 days.
Fusarium Pythium Rhizoctonia Fireblight ( <i>Erwinia amylovora</i> ) Bacterial Blight ( <i>Xanthomonas campestris</i> ) Anthracnose ( <i>Phytophthora</i> )	Trunk Injection	3 teaspoons/linear yard of canopy width or 2 inch of trunk diameter at breast height (DBH) (15 milliliters/linear meter of canopy width or 5 centimeters DBH.) Inject directly into tree.	Apply according to injection equipment instructions. Adjust rate based on injection equipment suggestions for specific applications.
	Root Dip	2 quarts/100 gallons of water; (2/3 fluid ounce/gallon of water) 5 liters/1000 liters water	Apply as a pre-plant dip to the roots for 15-30 minutes. Plant within 24 hours after dipping

Do not graze livestock in treated conifer nurseries and/or plantations. Do not feed forage to livestock from treated nurseries and/or plantations.

**TURF (INCLUDES RESIDENTIAL AND COMMERCIAL LAWN APPLICATIONS BY COMMERCIAL APPLICATORS)**

DM Fosphite is a systemic fungicide, which is used for the control of Pythium blight, Summer Complex (Bermuda and others), Dollar Spot, Grey Leaf Spot, Brown Patch, Anthracnose and damping-off diseases to turf grasses on sod farms and lawns, commercial turf production, golf courses (including tees and greens), parks and commercial landscapes. Begin preventative applications when conditions first favor disease and repeat at described intervals. Use higher rates under severe disease conditions.

Disease	Application Method	Rate	Application Program
Pythium (Yellow Turf) Rhizoctonia Sclerotinia Helminthosporium Alternaria Anthracnose ( <i>Colletotrichum graminicola</i> )	Foliar Spray	3-6 fluid ounces/1,000 square feet; (30-60 milliliters/100 square meters)  or  3 to 6 quarts/100 gallons of water/acre; (7 to 15 liters/1000 liters of water/hectare)	Apply amount in minimum of 2 gal. of water per 1,000 sq. ft. Apply at 10 to 21 day intervals as needed. Do not mow or irrigate treated areas until sprays have completely dried.  For sod farms and large areas: Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed.
	Irrigation	Overhead: 2-3 quarts/acre; (5-7 liters/hectare) in at least 100 gallons of water (1000 liters) of water	Apply with normal irrigation schedule. Apply at 10 to 21 day intervals and repeat as needed.

\* Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock and poultry.

**HOUSEHOLD PRODUCTS: FOR NON-COMMERCIAL APPLICATORS (NOT SUBJECT TO WORKER PROTECTION STANDARD)**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Not registered for homeowner use in the state of New York.

1 pint (0.473 liters) makes 24 gallons of solution at maximum treatment rate.

**DM FOSPHITE**  
**EPA REG. NO. 34704-924**

1 quart (0.946 liters) makes 48 gallons of solution at maximum treatment rate.  
 1 gallon (3.785 liters) makes 192 gallons of solution at maximum treatment rate.  
 2½ gallons (9.463 liters) makes 480 gallons of solution at maximum treatment rate.  
 5 gallons (18.925 liters) makes 960 gallons of solution at maximum treatment rate.

**GENERAL APPLICATION INSTRUCTIONS**

DM Fosphite is applied by various application methods, including foliar sprays, soil drench, soil incorporation, and bare root dip. For foliar sprays, apply DM Fosphite with sufficient water for adequate coverage of foliage, according to crop and growth stage. To insure good coverage, spray to wet.

**COMPATIBILITY**

Mixing DM Fosphite with certain surfactants, foliar fertilizers or other pesticides may cause crop injury. DM Fosphite is a slightly acidic buffer solution. Avoid mixing DM Fosphite with strongly acidic or alkaline materials. Do not tank mix without first testing the mixture's compatibility nor apply it without assessing its safety to the crop (phytotoxicity). It is not possible to test all material combinations and environmental tank mix combinations. Always conduct a test with the intended tank mix on a small area prior to any large scale applications. To determine if a combination is phytotoxic to a specific crop, spray a small area of foliage and fruit, and then evaluate 3-7 days later for visual symptoms.

Test the compatibility of spray adjuvants (i.e. stickers, spreaders, wetting agents) with DM Fosphite before use. Do not use acidifying type compatibility agents. To determine the compatibility of DM Fosphite with other products, use a jar compatibility test. Add the correct proportions of each product and the appropriate quantity of water to clean container, thoroughly mix, then let stand for 3-5 minutes. If the mixture remains in solution or can be remixed readily, the products are considered compatible.

**MIXING INSTRUCTIONS**

1. Fill the spray tank with ½ of the required volume of water.
2. Add the amount of DM Fosphite as indicated under the Rate portion of the table below.
3. Continue to fill the tank with water to the desired volume while agitating.
4. Continue agitation when applying.

This product may be applied up to the day of harvest.

**INSTRUCTIONS FOR APPLICATIONS**  
**GARDEN CROPS, ORNAMENTALS, BEDDING PLANTS**  
**AND TREES (CONIFERS, OAKS, ETC.) & TURF**

DM Fosphite fungicide is used for control of *Pythium*, Downy Mildew, *Phytophthora* (such as Sudden Oak Death, *Phytophthora ramorum*) and various other diseases (see list below) of ORNAMENTAL PLANTS AND TREES. Plants, which can be treated with DM Fosphite, include:

**GARDEN CROPS**

DM Fosphite fungicide is used for control of Downy Mildew, Powdery Mildew, *Pythium*, *Phytophthora* and various other diseases (such as Purple Blotch, Late Blight, Blue Mold, Soft Rot, Root Rot, Brown Rot, Foot Rot, Crown/Collar Rot, Canker, Fruit/Foliar Rot, Fire Blight, Heart Rot, Pink Rot, *Pythium* Leak, Red Stele, Leather Rot, etc.) of garden crops. Apply DM Fosphite to plants including (but not limited to): Almond, Apple, Apricot, Artichoke, Asparagus, Avocado, Baby Lettuce, Banana, Beet, Blackberry, Blueberry, Broadbean, Cane Berries, Carrot, Cherry, Chestnut, Chilies, Cinchona, Cinnamon, Citrus, Cole Crops, Corn, Cranberry, Crucifers, Cucurbits, Dates, Dry Beans, Eggplant, Fresh beans, Garlic, Grapes, Kiwi, Lettuce, Lima Beans, Macadamia, Melons, Mint, Nectarine, Nut Crops, Nutmeg, Okra, Olive, Onion, Papaya, Peach, Peanut, Pear, Peas, Peppers, Pineapple, Plum, Pome Fruit, Potato, Prune, Raspberry, Rhubarb, Root crops, Rosemary, Spinach, Squash, Strawberry, String Beans, Stone Fruit, Sugarbeet, Sunflower, Tomato, Walnut.

**BEDDING PLANTS, TREES & ORNAMENTALS**

Ageratum, Algerian Ivy, Aglaonema, Alyssum, Anthurium, Aphelandra, Arborvitae, Artemisia, Ash, Aster, Azaleas, Baby's Breath, Begonia, Bougainvillea, Boxwood, Caladium, Carnation, Cattleya skinneri, Ceanothus, Chrysanthemum, Coleus, Columbine, Cotoneaster, Cissus, Daisy, Delphinium, Dieffenbachia, Dogwood, Easter Lily, English Ivy, Ficus, Firs, Foxglove, Gaillardia, Geranium, Gloxinia, Hibiscus, Impatiens, Japanese Holly, Juniper, Leather-leaf Fern, Magnolia, Maple, Marigold, Monterey Pine, Oaks, Pansy, Petunia, Peperomia, Philodendron, Phlox, Photinia, Pines, Pinks, Pittosporum, Pieris, Poinsettia, Pothos, Primrose, Prostrate Rosemary, Rhododendron, Roses, Salvia, Schefflera, Sedum, Sempervivum, Snapdragon, Spathiphyllum, Taxus media, Spruces, Sycamore, Verbena, Vinca, White Cedar, White Pine, Zygocactus.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the use rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Applicatio. Method	Rate	Application Program
<i>Phytophthora</i> <i>Fusarium</i> <i>Pythium</i> <i>Rhizoctonia</i> Fire Blight ( <i>Erwinia amylovora</i> ) Anthracnose	Foliar Spray	1/3 fluid ounce to 2/3 fluid ounce/gallon of water. (2.5-5 milliliters/liter of water)	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply DM Fosphite at intervals less than 3 days.
	Soil Drench	3-6 fluid ounces/25 gallons of water or 1/8 to 1/4 fluid ounce per gallon of water (1 to 2 liters/1000 liters of water)	Apply 1-2 qts. of diluted solution per sq. ft. Adjust volume as needed to thoroughly wet media. Apply at 2-3 week intervals and repeat as needed. Do not apply DM Fosphite at intervals less than 3 days.
	Soil Incorporation	1/2 to 1 pint/ 1/2 cubic yard. (150-300 milliliters/ 1/2 cubic meter).	Mix 1/2 to 1 pint of DM Fosphite with enough water to wet one cubic yard of soil; mix immediately before potting. If conditions are favorable for disease development, make additional applications as a foliar spray or soil drench.
	Root Dip	2/3 fluid ounce/gallon of water (5 milliliters/liter of water).	Dip transplants into diluted solution of DM Fosphite to thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.
Downy Mildew Powdery Mildew	Foliar Spray	1/3 fluid ounce to 2/3 fluid ounce/gallon of water (2.5-5 milliliters/liter of water).	Apply diluted solution to thoroughly wet foliage. Apply at 2-3 week intervals and repeat as needed. Do not apply DM Fosphite at intervals less than 3 days.

**TURF**

DM Fosphite is a systemic fungicide, which is used for the control of *Pythium* blight, Summer Complex (Bermuda and others), Dollar Spot, Grey Leaf Spot, Brown Patch, Anthracnose and damping-off diseases to turf grasses. Begin preventative applications when conditions first favor disease and repeat at directed intervals. Use higher rates under severe disease conditions.

Disease	Application Method	Rate	Application Program
<i>Pythium</i> (Yellow Turf) <i>Rhizoctonia</i> <i>Sclerotinia</i> <i>Helminthosporium</i> <i>Alternaria</i> Anthracnose ( <i>Colletotrichum graminicola</i> )	Foliar Spray	3-6 fluid ounces/ 1,000 sq. ft. (30-60 milliliters/ 100 square meters).	Apply amount in minimum of 2 gal. of water per 1,000 sq. ft. Apply at 7 to 21 day intervals as needed. Do not mow or irrigate treated areas until sprays have completely dried.
Late Blight ( <i>Phytophthora infestans</i> ) Pink Rot ( <i>Phytophthora erythroseptica</i> )	Post-harvest Treatment of Potatoes: Spray, rinse or inject into humidification water prior to or during storage.	Dilute DM Fosphite at 1:20 ratio with application water. 6.4-12.8 fluid ounces/ 0.5 gallon of water / 1000 tubers; (95-190 milliliter/liter of water)	Inject concentrate into water used in humidification or process water used for post-harvest storage, rinses and associated tanks, flumes and lines.

**STORAGE AND DISPOSAL**

**PESTICIDE STORAGE:** Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry place.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinstate). After emptying and cleaning, it may be allowable to temporarily hold residue or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC)

13/13

**DM-FOSPHITE**  
**EPA REG. NO. 34704-924**

**Storage & Disposal cont'd.:**

at [www.acrecycle.org](http://www.acrecycle.org).

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.  
*(For packages up to 5 gallons:)* **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

*(For packages greater than 5 gallons and less than 56 gallons:)* **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

*(For packages greater than 56 gallons:)* To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.

*(Red text in italics will not be printed.)*

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**  
BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS PRODUCT IS SOLD AS IS TO THE EXTENT ALLOWED BY APPLICABLE LAW. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND, TO THE EXTENT REQUIRED BY APPLICABLE LAW, WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, 7251 WEST 4TH STREET, GREELEY, CO 80634.

TO THE EXTENT ALLOWED BY APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT ALLOWED BY APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

FORMULATED FOR



P.O. BOX 1286, GREELEY, COLORADO 80632-1286