

34704-931

09/22/2009

1/10

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



SEP 22 2009

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

John Tice
Loveland Products, Inc.
7251 W 4th Street
P.O. Box 1286
Greeley, CO 80632

Dear Mr. Tice,

Subject: Amendment to change primary brand name to "Wrangler Insecticide," revised warranty statement and increase cotton application rates
Wrangler Insecticide
EPA Registration No. 34704-931

The labeling referred to above submitted in connection with the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable.

Please submit one final printed copy of the labeling before releasing the product for shipment. A stamped copy of the labeling is enclosed for your records. If you have any questions regarding this label, please contact Autumn Metzger at (703) 305-5314 or metzger.autumn@epa.gov.

Sincerely,

Venus Eagle
Product Manager 01
Insecticide-Rodenticide Branch
Registration Division (7505P)

Wrangler™

INSECTICIDE

ACCEPTED

SEP 22 2009

2/10

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:

EPA. Reg. No: 34704-931

For protection of cotton from certain insects.

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolindinimine 40.7%

INERT INGREDIENTS: 59.3%

TOTAL 100.0%

Contains 4 pounds of active ingredient per gallon.

Shake well before using.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.

Note to Physician: No specific antidote is available. Treat the patient symptomatically.

EPA REG. NO. 34704-931

EPA EST. NO. 34704-MS-001

NET CONTENTS 1 GAL. (3.78 L)

IHT

EXP D 06/09

**WRANGLER™ INSECTICIDE
EPA REG. NO. 34704-931**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, absorbed through skin, or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical resistant to this product are listed below. More options can be obtained by following the instructions for Category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: Long sleeved shirt and long pants, chemical resistant gloves made of any waterproof material such as, nitrile rubber, butyl rubber, neoprene rubber, barrier laminate, polyethylene, polyvinyl chloride (PVC) or viton, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170-240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove Personal Protective Equipment immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

OBSERVE THE FOLLOWING PRECUATIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WRANGLER™ INSECTICIDE
EPA REG. NO. 34704-931

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR part 170.

This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls,
- Chemical-resistant gloves made of any waterproof material such as, nitrile rubber, butyl rubber, neoprene rubber, barrier laminate, polyethylene, polyvinyl chloride (PVC) or viton, and
- Shoes plus socks.

Spray Drift Management

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

Mixing and Loading

To avoid potential contamination of groundwater, the use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading areas and potential surface to groundwater conduits such as field sumps, uncased well head, sinkholes or field drains.

For Aerial Applications

The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Importance of Droplet Size

An important factor influencing drift is droplet size. Small droplets (<150-200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

**WRANGLER™ INSECTICIDE
EPA REG. NO. 34704-931**

Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions

Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

No-Spray Zone Requirements for Foliar Applications

Do not apply by ground within 25 feet, or by air within 150 feet of lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries and commercial fish farm ponds.

Runoff Management

Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip.

When used on erodible soils, use best management practices for minimizing runoff. Consult your local Natural Resources Conservation Service for recommendations in your use area.

Endangered Species Notice

Under the Endangered Species Act, It is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent or Pesticide State Lead Agency for information concerning endangered species in your area.

Resistance Management

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

WRANGLER™ Insecticide contains a Group 4A insecticide called imidacloprid. Insect biotypes with acquired or inherent tolerance to Group 4A products may eventually dominate the insect population if Group 4A products are used repeatedly as the predominant method of control for targeted species. This may eventually result in partial or total loss of control of those species by WRANGLER Insecticide and to other Group 4A products.

The active ingredient in WRANGLER Insecticide is a member of neonicotinoid chemical group. Avoid using a block of more than three consecutive applications of WRANGLER Insecticide and/or other Group 4A products having the same or similar mode of action. Following a neonicotinoid block of treatments, Loveland Products, Inc. strongly encourages the rotation to a block of applications with effective products from a different mode of action before using additional applications of neonicotinoid products. Using a block of rotation or windowed approach, along with other IPM practices,

6/10

**WRANGLER™ INSECTICIDE
EPA REG. NO. 34704-931**

is considered an effective use strategy for preventing or delaying an insect pest's ability to develop resistance to this class of chemistry.

Do not make foliar applications of WRANGLER Insecticide or other Group 4A products on crops previously treated with a long-residual, soil-applied product from the neonicotinoid chemical class.

Other Group 4A neonicotinoid products used as foliar treatments include: Actara® , Assail® , CALYPSO® , Centric® , Intruder™ , LEVERAGE™ and PROVADO® . Other 4A Group neonicotinoid products used as soil treatment include: ADMIRE® and Platinum® .

Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org/>.

USE INSTRUCTIONS

Restrictions:

Do not apply WRANGLER Insecticide through any type of irrigation system.
Do not apply WRANGLER Insecticide in enclosed structures such as greenhouses or plant houses.

MIXING INSTRUCTIONS

Minimum spray volumes are 10 gallons/acre by ground application and 5 gallons/acre through aerial equipment.

To prepare the application mixture, add half of the required amount of water to the spray tank and with agitation add WRANGLER Insecticide. Complete filling tank with balance of water needed. Maintain sufficient agitation during both mixing and application. WRANGLER Insecticide may also be used with other pesticides and/or fertilizer solutions. Please see Compatibility Note below. When tank mixtures of WRANGLER Insecticide and other pesticides are involved, prepare the tank mixture as instructed above and follow Mixing Order below.

Mixing Order

When pesticide mixtures are needed, add wettable powders first, WRANGLER Insecticide or other flowables second, and emulsifiable concentrates last. Ensure good agitation as each component is added. Do not add an additional component until the previous is thoroughly mixed. If a fertilizer solution is added, a fertilizer-pesticide compatibility agent may be needed. Maintain constant agitation during both mixing and application to ensure uniformity of spray mixture.

Compatibility Note

Test compatibility of the intended tank mixture before adding WRANGLER Insecticide to the spray or mix tank. Add proportionate amounts of each ingredient in the appropriate order, to a pint or quart jar, cap, shake for 5 minutes, and let set for 5 minutes. Do not use if poor mixing or formation of precipitates that do not readily re-disperse. This indicates an incompatible mixture. For further information, contact your local Loveland Products, Inc. representative.

ROTATIONAL CROPS

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for imidacloprid, as soon as practical following the last application. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed. Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

WRANGLER™ INSECTICIDE
EPA REG. NO. 34704-931

IMMEDIATE PLANT-BACK

All crops on this label plus the following crops:

Barley	Head & stem Brassica	Root vegetables
Canola	vegetables	Sorghum
Corn (field, pop and sweet)	Leafy vegetables	Strawberry
Corn vegetables	Leafy petiole vegetables	Sugar beet
Cotton	Legume vegetables	Tobacco
Cucurbit vegetables	(except soybean, dry)	Tuberous vegetables
Fruiting vegetables	Potato	Wheat
Globe artichoke	Rape Seed	

30-DAY PLANT-BACK

Cereals (including buckwheat, millet, oats, rice, rye, and triticale)
Safflower
Soybean, Dry

12-MONTH PLANT BACK

All Other Crops

APPLICATION INFORMATION — WRANGLER INSECTICIDE

Apply WRANGLER Insecticide with properly calibrated ground or aerial application equipment. Apply specified rate per acre as a directed or broadcast spray to infested area at earliest threshold for target pest, as population begins to develop. Thorough uniform coverage of all plant parts is required to achieve optimum control. Scout fields and retreat if needed.

The lower rates can be used early season when pest pressures are low or when tank-mixing with other effective products registered for target insect control. Degree of control or suppression of additional labeled pests will be determined, in part, by the stage of pest development at application and infestation level of those pests. WRANGLER Insecticide provides optimal performance against early instar and early nymphal stages of insects as well as bollworm/budworm eggs. Applications made with less than 5 gallons per acre may result in slower activity and/or less overall control from a single application than an application made with higher gallonages. Use an organosilicone-based spray adjuvant for applications targeting aphids and whiteflies.

Regardless of formulation or method of application, apply no more than 0.5 lb active ingredient of imidacloprid per acre per season, including seed treatment, soil and foliar uses.

WRANGLER™ INSECTICIDE
EPA REG. NO. 34704-931

COTTON

Pests Controlled	Rate Fluid Ounces/Acre
Cotton aphid	0.9 - 1.8
Cotton fleahopper	
Bandedwinged whitefly	
Plant bugs (excludes <i>Lygus hesperus</i>)	
Green stink bug	
Southern green stink bug	
Bollworm/Budworm (ovicidal effect)	

Pests Suppressed	Rate Fluid Ounces/Acre
<i>Lygus</i> bug (<i>Lygus hesperus</i>)	1.35 - 1.8
Whiteflies (other than bandedwinged whitefly)	

Restrictions

Pre-Harvest (PHI): **14 days**
 Minimum interval between applications: **7 days**
 Maximum WRANGLER Insecticide allowed per season: **8.9 fluid ounces/Acre** (0.31 lb. AI/A)
 Maximum number of WRANGLER Insecticide applications per season: **5**

Do not graze treated fields after any application of WRANGLER Insecticide.

Tank Mix Information

Pests Controlled (In addition to pests listed above)	WRANGLER Insecticide Rate Fl. oz./A	Bidrin® 8 Rate Fl. oz./A
For early season control of: Thrips	0.9 - 1.35	1.6 - 3.2
For mid to late season control of: Plant bugs Stink bugs (including Brown stink bug) Grasshoppers Saltmarsh caterpillar Cotton leafperforator	0.9 - 1.35	4.0 - 8.0

Restrictions (in addition to Restrictions listed above)

Refer to the Bidrin 8 product label for specific use rates; observe all restrictions and precautions that appear on the label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place, out of direct sunlight, and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

9/10

WRANGLER™ INSECTICIDE
EPA REG. NO. 34704-931

Storage & Disposal cont'd.:

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org. If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For packages up to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 56 gallons: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.

10/10

**WRANGLER™ INSECTICIDE
EPA REG. NO. 34704-931**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Admire, Calypso and Provado are registered trademarks and Leverage is a trademark of Bayer. Actara, Centric and Platinum are registered trademarks of a Syngenta Group Company. Assail is a registered trademark and Intruder is a trademark of Nippon Soda Company, LTD. Bidrin is a registered trademark of AMVAC Chemical Corporation. Wrangler™ is a trademark of Loveland Products, Inc. Bidrin and Leverage are Restricted Use Pesticides.

FORMULATED FOR



**P.O. BOX 1286, GREELEY, COLORADO 80632-1286
1-888-574-2878**