

34704-931

05-03-2011

$\frac{1}{33}$



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

MAY 03 2011

Dr. Premjit Halarnkar
Loveland Products, Inc.
P.O. Box 1286
Greeley, Colorado 80632-1286

Subject: Revised Label Use Instructions Section
Wrangler Insecticide
EPA Reg. No. 34704-931
Your Submission date, April 15, 2011

Dear Dr. Halarnkar:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act is acceptable. A stamped copy of the label is enclosed for your records. Submit two (2) copies of your final printed labeling before you release the product for shipment. If there are questions call Dani Daniel at 703 305-5409.

Sincerely,

A handwritten signature in cursive script that reads "Venus Eagle".

Venus Eagle
Product Manager (01)
Insecticide-Rodenticide Branch
Registration Division (7504P)

Enclosure:

2
33



Wrangler[®]

INSECTICIDE

ACTIVE INGREDIENT:

Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine 40.7%

OTHER INGREDIENTS 59.3%

TOTAL 100.0%

Contains 4 pounds of Imidacloprid per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SHAKE WELL BEFORE USING

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</p> <p>FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.</p> <p>Note to Physician: No specific antidote is available. Treat the patient symptomatically.</p>	

ACCEPTED

MAY 03 2011

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under:

EPA REG. NO. 34704-931

EPA EST. NO. 34704-MS-001

NET CONTENTS 1 GAL. (3.78 L)

EPA. Reg. No: 34704-931

EXP 04/11 Irrigation

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed, absorbed through skin, or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical resistant to this product are listed below. More options can be obtained by following the instructions for Category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: Long sleeved shirt and long pants, chemical resistant gloves made of any waterproof material such as, nitrile rubber, butyl rubber, neoprene rubber, barrier laminate, polyethylene, polyvinyl chloride (PVC) or viton, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170-240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove Personal Protective Equipment immediately after handling this product.
- Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls,
- Chemical-resistant gloves made of any waterproof material such as, nitrile rubber, butyl rubber, neoprene rubber, barrier laminate, polyethylene, polyvinyl chloride (PVC) or viton, and
- Shoes plus socks.

Spray Drift Management

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

Mixing and Loading

To avoid potential contamination of groundwater, the use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading areas and potential surface to groundwater conduits such as field sumps, uncased well head, sinkholes or field drains.

For Aerial Applications

Mount the spray boom on the aircraft so as to minimize drift caused by wing tip vortices. Use the minimum practical boom length, do not exceed 75% of the wing span or rotor diameter.

Release spray at the lowest possible height consistent with good pest control and flight safety. Do not make applications more than 10 feet above the crop canopy.

Importance of Droplet Size

An important factor influencing drift is droplet size. Small droplets (<150-200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, make applications to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure.

Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions

Do not make aerial or ground applications during temperature inversions. Drift potential is high during temper-

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

ature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

No-Spray Zone Requirements for Foliar Applications

Do not apply by ground within 25 feet, or by air within 150 feet of lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries and commercial fish farm ponds.

Airblast (Air Assist) Specific Applications for Tree Crops and Vineyards

Airblast sprayer carry droplets into the canopy of trees/vineyards via a radially, or laterally directed air stream. The following specific drift management practices should be followed:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy.
- Block off upward pointed nozzles when there is no overhanging canopy.
- Use only enough air volume to penetrate the canopy and provide good coverage.
- Do not allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows).
- Only spray inward, toward the orchard or vineyard, for application to the outside rows.

Runoff Management

Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip.

When used on erodible soils, use best management practices for minimizing runoff. Consult your local Natural Resources Conservation Service for recommendations in your use area.

Endangered Species Notice

Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent or Pesticide State Lead Agency for information concerning endangered species in your area.

Resistance Management

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

WRANGLER® Insecticide contains a Group 4A insecticide called imidacloprid. Insect biotypes with acquired or inherent tolerance to Group 4A products may eventually dominate the insect population if Group 4A products are used repeatedly as the predominant method of control for targeted species. This may eventually result in partial or total loss of control of those species by WRANGLER Insecticide and to other Group 4A products.

The active ingredient in WRANGLER Insecticide is a member of neonicotinoid chemical group. Avoid using a block of more than three consecutive applications of WRANGLER Insecticide and/or other Group 4A products having the same or similar mode of action. Following a neonicotinoid block of treatments, Loveland Products, Inc. strongly encourages the rotation to a block of applications with effective products from a different mode of action before using additional applications of neonicotinoid products. Using a block of rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect pest's ability to develop resistance to this class of chemistry.

Do not make foliar applications of WRANGLER Insecticide or other Group 4A products on crops previously treated with a long-residual, soil-applied product from the neonicotinoid chemical class.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

Other Group 4A neonicotinoid products used as foliar treatments include: Actara®, Assail®, CALYPSO®, Centric®, Intruder®, LEVERAGE® and PROVADO®. Other 4A Group neonicotinoid products used as soil treatment include: ADMIRE® and Platinum®.

Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org/>.

MIXING INSTRUCTIONS

Minimum spray volumes are 10 gallons/acre by ground application and 5 gallons/acre through aerial equipment. To prepare the application mixture, add half of the required amount of water to the spray tank and with agitation add WRANGLER Insecticide. Complete filling tank with balance of water needed. Maintain sufficient agitation during both mixing and application. WRANGLER Insecticide may also be used with other pesticides and/or fertilizer solutions. Please see Compatibility Note below. When tank mixtures of WRANGLER Insecticide and other pesticides are involved, prepare the tank mixture as instructed above and follow Mixing Order below.

Mixing Order

When pesticide mixtures are needed, add wettable powders first, WRANGLER Insecticide or other flowables second, and emulsifiable concentrates last. Ensure good agitation as each component is added. Do not add an additional component until the previous is thoroughly mixed. If a fertilizer solution is added, a fertilizer-pesticide compatibility agent may be needed. Maintain constant agitation during both mixing and application to ensure uniformity of spray mixture.

Compatibility Note

Test compatibility of the intended tank mixture before adding WRANGLER Insecticide to the spray or mix tank. Add proportionate amounts of each ingredient in the appropriate order, to a pint or quart jar, cap, shake for 5 minutes, and let set for 5 minutes. Do not use if poor mixing or formation of precipitates that do not readily re-disperse. This indicates an incompatible mixture. For further information, contact your local Loveland Products, Inc. representative.

CHEMIGATION

Types of Irrigation Systems: Foliar chemigation applications of Wrangler Insecticide may be made to crops through overhead sprinkler systems if specified in crop-specific application sections. Soil chemigation application of Wrangler Insecticide may only be made to crops through chemigation as specified in crop-specified application sections and only through low-pressure systems specifically for a given crop. Do not apply Wrangler Insecticide through any other type of irrigation system.

Make foliar chemigation applications of Wrangler Insecticide as concentrated as possible. Retention of Wrangler Insecticide on target site of insect infestation is necessary for optimum activity. Do not use chemigation of Wrangler Insecticide in water volumes exceeding 0.10 inches per acre. See crop-specific application sections of the label for more information.

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have any questions about calibration, contact Cooperative Extension Service specialists, equipment manufacturers, or other experts.

**WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931**

Chemigation Monitoring: A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift: Do not apply when wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water from Public Water Systems: Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, back flow preventer (RPZ), or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shuts down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

ROTATIONAL CROPS*

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval must be observed.

IMMEDIATE PLANT-BACK:

All crops on this label plus the following crops not on this label: barley, canola, Christmas trees, corn (field, sweet and pop), cranberry, Globe artichoke, grape, mustard seed, okra, potato, rapeseed, strawberry, sorghum, sunflower, tobacco, watercress, wheat and all crops from the following Crop Groups as recognized and defined by EPA. Crops contained within a particular crop group are subject to change. For information related to specific crops, please contact your Loveland Products, Inc. representative or refer to EPA website (www.epa.gov) for latest crop groups.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

ROOT VEGETABLES - Crops of Crop Group 1
 LEAFY GREEN VEGETABLES - Crops of Crop Group 4
 HEAD and STEM BRASSICA VEGETABLES - Crops of Crop Group 5
 LEGUME VEGETABLES - Crops of Crop Group 6 including: Edible Podded plus Dried plus Succulent Shelled,
 Peas and Beans
 FRUITING VEGETABLES - Crops of Crop Group 8
 CUCURBIT VEGETABLES - Crops of Crop Group 9
 CITRUS - Crops of Crop Group 10
 POME FRUIT - Crops of Crop Group 11
 STONE FRUIT - Crops of Crop Group 12
 BUSHBERRY and CANEBERRY - Crops of Crop Group 13-07
 HERBS - Crops of Crop Group 19A
 TROPICAL FRUIT – Including: Acerola, Atemoya, Avocado, Birida, Black sapote, Canistel, Cherimoya, Custard
 apple, Feijoa, Llama, Jaboticaba, Guava, Longan, Lychee, Mamey sapote, Mango, Papaya, Passion fruit,
 Persimmon, Pulasan, Rambuten, Sapodilla, Soursop, Spanish lime, Star apple, Starfruit, Sugar apple, Wax
 jambu

30-DAY PLANT-BACK:

Cereals (including buckwheat, millet, oats, rice, rye, and triticale), soybeans and safflower

10-MONTH PLANT-BACK:

Onion and bulb vegetables

12-MONTH PLANT-BACK:

All Other Crops

*Cover crops for soil building or erosion control may be planted any time; but do not graze or harvest for food or feed.

APPLICATION INFORMATION — WRANGLER INSECTICIDE

Apply WRANGLER Insecticide with properly calibrated ground or aerial application equipment. Apply specified rate per acre as a directed or broadcast spray to infested area at earliest threshold for target pest, as population begins to develop. Thorough uniform coverage of all plant parts is required to achieve optimum control. Scout fields and retreat if needed.

The lower rates can be used early season when pest pressures are low or when tank-mixing with other effective products registered for target insect control. Degree of control or suppression of additional labeled pests will be determined, in part, by the stage of pest development at application and infestation level of those pests. WRANGLER Insecticide provides optimal performance against early instar and early nymphal stages of insects as well as bollworm/budworm eggs. Applications made with less than 5 gallons per acre may result in slower activity and/or less overall control from a single application than an application made with higher gallonages. Use an organosilicone-based spray adjuvant for applications targeting aphids and whiteflies.

Regardless of formulation or method of application, apply no more than 0.5 lb active ingredient of imidacloprid per acre per season, including seed treatment, soil and foliar uses.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

GLOBE ARTICHOKE*

Pests Controlled	Rate fluid ounces/Acre
Aphids	8.0 – 16.0
Leafhoppers	

Restrictions:

Pre-Harvest Interval (PHI): 7 days

Maximum Wrangler Insecticide amount allowed per season: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

Applications

Apply specified dosage in the following method:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
2. In-furrow spray at planting directed on or below seed.

*Use not permitted in California unless otherwise directed by supplemental labeling.

HERBS

Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Bumet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dillweed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

Pests Controlled	Rate fluid ounces/Acre
Aphids	8.0 – 12.0
Flea beetles	
Leafhoppers	
Whiteflies	

Pests/Diseases Suppressed

Thrips (foliage-feeding thrips only)	8.0 – 12.0
--------------------------------------	------------

Restrictions:

Pre-Harvest Interval (PHI): **14 days**

Maximum Wrangler Insecticide allowed per crop season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

Instructions

Apply specified dosage in one of the following methods:

1. In-furrow spray during planting directed on or below seed;
2. In-furrow spray or transplant-water drench during setting or transplanting;
3. Shanked-into or below eventual seed-line;
4. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, Loveland Products, Inc. strongly recommends that only small areas or numbers of plants of each be treated and evaluated prior to commercial use.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

FIELD CROPS
Application Instructions – Wrangler Insecticide

COTTON - Soil Treatment

Pests Controlled	Rate fluid ounces/1000 row-feet	Rate fluid ounces/Acre
Cotton aphid	0.65	8.5 – 10.5 (Depending on row-spacing)
Plant bugs		
Thrips		
Whiteflies		

Restrictions

Maximum Wrangler Insecticide allowed per crop season: **10.5 fluid ounces/Acre** (0.33 lb AI/Acre)

Regardless of formulation or method of application, apply no more than 0.5 lb active ingredient of Wrangler Insecticide, Provado, Trimax or Leverage per acre per season, including seed treatment as Gaucho®, soil and foliar uses. Do not apply more than a total of 6 applications of the active ingredient per season. Do not graze treated fields after any application of Wrangler Insecticide. Please see Resistance Management section of this label.

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. In-furrow spray during planting directed on or below seed;
2. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting;
3. Chemigation into root-zone through low-pressure drip or trickle irrigation.

COTTON - Foliar Treatment

Pests Controlled	Rate fluid ounces/Acre
Cotton aphid	1.0 – 2.0
Cotton fleahopper	
Bandedwinged whitefly	
Plant bugs (excludes <i>Lygus hesperus</i>)	
Green stink bug	
Southern green stink bug	
Bollworm/Budworm (ovicidal effect)	

Pests Suppressed

Lygus bug (<i>Lygus hesperus</i>)	1.5 – 2.0
Whiteflies (other than bandedwinged whitefly)	

Restrictions

Pre-Harvest (PHI): **14 days**

Minimum interval between applications: **7 days**

Maximum WRANGLER Insecticide allowed per season: **10 fluid ounces/Acre** (0.31 lb. AI/A)

Maximum number of WRANGLER Insecticide applications per season: **5**

Do not graze treated fields after any application of WRANGLER Insecticide.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

Tank Mix Information

Pests Controlled (In addition to pests listed above)	WRANGLER Insecticide Rate Fl. oz./A	Bidrin® 8* Rate Fl. oz./A
For early season control of: Thrips	1.0 – 1.5	1.6 – 3.2
For mid to late season control of: Plant bugs Stink bugs (including Brown stink bug) Grasshoppers Saltmarsh caterpillar Cotton leafperforator	1.0 – 1.5	4.0 – 8.0
Restrictions (in addition to Restrictions listed above) *Refer to the Bidrin 8 product label for specific use rates; follow all restrictions and precautions that appear on the label.		

PEANUT*

Pests Controlled	Rate fluid ounces/Acre
Aphids Leafhoppers Whiteflies	8.0 – 12.0
Pest Suppressed Thrips	8.0 – 12.0

Restrictions:

Pre-Harvest Interval (PHI): 14 days

Maximum Wrangler Insecticide allowed per season: **12.0 fluid ounces/Acre** (0.38 lb ai/Acre)

Applications:

Apply specified dosage in one of the following methods:

1. In-furrow spray during planting directed on or below seed;
2. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Notes

Increases in Tomato spotted wilt virus (TSWV) incidence have been observed with applications of Wrangler Insecticide on multiple varieties of peanut. This may also be the case with other tospoviruses, or other viruses transmitted by various thrips species or perhaps, other pests. Prior to applying Wrangler Insecticide to peanuts, Loveland Products, Inc. recommends consultation with the State, Cooperative Extension Service, or Loveland Products, Inc. representative, for recommendations. Growers are advised to weigh insect control benefits against potential increase in viral disease levels. In areas where TSWV or other tospovirus are endemic, growers are encouraged to use virus resistant varieties and consult the University of Georgia, Tomato spotted wilt virus index, before applying Wrangler Insecticide.

*Use not permitted in California unless otherwise directed by supplemental labeling.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

POTATO

Pests Controlled	Rate	Rate
	fluid ounces/1000 row-feet	fluid ounces/Acre
Aphids		
Colorado potato beetle		
Flea beetles	0.45 – 0.65	6.5 – 10.0
Leafhoppers		
Potato psyllid		
Pests/Diseases Suppressed		
Symptoms of:		
Potato leaf roll virus (PLRV)		
Potato yellows	0.45 – 0.65	6.5 – 10.0
Net necrosis (PLRV)		
Wireworms (with in-furrow spray at-planting)		

Restrictions

Maximum Wrangler Insecticide allowed per crop season: **10.0 fluid ounces/Acre** (0.31 lb AI/Acre)

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. In-furrow spray during planting directed on seed pieces or seed potatoes;
 2. Subsurface side-dress on both sides on the row covered with 3 or more inches of soil;
 3. Narrow band spray at ground cracking directly over the row during hilling covered with 3 or more inches of soil;
 4. Narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting.
- For effective pest control or suppression, Wrangler Insecticide applications must be placed below soil-surface and in contact with seed piece or within root-zone. For potatoes grown on highly permeable soils with shallow water table, at-plant applications of Wrangler Insecticide may be made in a 2 to 4 inch band (width of planter shoe opening) and completely covered.

POTATO

(Seed Piece Treatment)

Pests Controlled	Rate	Rate
	fluid ounces/100 lbs. seed	fluid ounces/Acre**
Aphids		
Colorado potato beetle		
Flea beetles	0.2 – 0.4	4.0 – 8.0
Leafhoppers		
Potato psyllid		
Wireworms (seed-piece protection)		
Pests/Diseases Suppressed		
Symptoms of:		
Potato leaf roll virus (PLRV)		
Potato yellows	0.4	8.0
Net necrosis (PLRV)		

Restrictions

Maximum Wrangler Insecticide allowed per crop season: **10.0 fluid ounces/Acre** (0.31 lb AI/Acre)

Do not use treated seed-pieces for food, feed, or fodder. Do not apply any subsequent application of Wrangler Insecticide (in-furrow), Gaucho, Leverage or Provado following a Wrangler Insecticide seed-piece treatment.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

Potato
(Seed Piece Treatment) cont'd.:

Instructions

Apply specified dosage as a diluted spray onto seed-pieces using a shielded spray system. Dilute with 3 parts water, or less, to 1 part Wrangler Insecticide. Agitate or stir spray solution as needed. Fungicidal or inert absorbent dusts may be applied after Wrangler Insecticide application. Apply only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Plant seed-pieces as soon as possible after treating avoiding prolonged exposure of Wrangler Insecticide treated seed-pieces to sunlight and in accordance with the directions of your local Extension specialist.

Consult your local Loveland Products, Inc. representative or crop protection product dealer for information relevant to your area.

**Based on a seeding rate of 2000 lbs/acre.

TOBACCO

Pests Controlled	Rate fluid ounces/1000 plants (as seedling tray drench)	Rate fluid ounces/1000 plants (in-furrow or transplant-water)
Aphids		
Flea beetles	0.5	0.7
Mole crickets		
Whiteflies	0.7 – 1.4	0.9 – 1.4
Wireworms		
Pests/Diseases Suppressed		
Cutworms		
Symptoms of: Tomato spotted wilt virus (TSWV)	0.7 – 1.4	0.9 – 1.4

Restrictions

Pre-Harvest Interval (PHI): **14 days**

Maximum Wrangler Insecticide allowed per crop season: **16.0 fluid ounces/Acre** (0.50 lb AI/Acre)

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. Uniform, broadcast foliar spray to seedlings in trays (tray drench) not more than 7 days prior to transplanting followed immediately by overhead irrigation to wash Wrangler Insecticide from foliage into potting media. Failure to wash Wrangler Insecticide from foliage may result in reduction in pest control. Transplants must be handled carefully during setting to avoid dislodging treated potted media from roots.
2. In-furrow spray or transplant-water drench during setting.
3. Chemigation into root-zone through low-pressure drip, trickle, micro sprinkler or equivalent equipment.

Important Note: Proper tray drench applications of Wrangler Insecticide have been shown to be the most efficacious method of application. However, the specified rate of Wrangler Insecticide may be applied as a combination of the tray drench in the planthouse and/or transplant-water drench in field. Adverse growing conditions may cause a delay in uptake of Wrangler Insecticide into the plant and a delay in control.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

VEGETABLE and SMALL FRUIT CROPS
Application Directions – Wrangler Insecticide

Restrictions

Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

CUCURBIT-VEGETABLES

Including: Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourd (edible; includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* spp. (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melo* including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, and Winter melon), Pumpkin, Squash (includes summer squash types such as: butternut squash, calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash and spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

Field application instructions. See details below for additional planthouse instructions.

Pests Controlled

Rate
fluid ounces/Acre

Aphids

Cucumber beetles

Leafhoppers

8.0 – 12.0

Thrips (foliage-feeding thrips only)

Whiteflies

Pests/Diseases Suppressed

Bacterial wilt (as vectored by various cucumber beetles)

Leaf silvering resulting from whitefly feeding

8.0 – 12.0

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum Wrangler Insecticide allowed per application: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. In-furrow spray directed on or below seed;
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1-1/2" with sufficient irrigation within 24 hours of application;
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;
5. Post-seeding drench, transplant-water drench, or hill drench;
6. Subsurface side-dress on both sides of each row. Wrangler Insecticide must be incorporated into root-zone.

Planthouse Application Instructions*

Pests Controlled

Rate
fluid ounces/1000 Plants

Aphids

Whiteflies

0.05

Restrictions

Maximum amount of Wrangler Insecticide applied in the planthouse: **0.05 fluid ounces** (0.00156 lb AI)/1000 plants.

Maximum number Wrangler Insecticide applications in planthouse: **1**

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

Cucurbit Vegetables cont'd:

Instructions:

Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners:

1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash Wrangler Insecticide from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash Wrangler Insecticide from foliage may result in reduced pest control;
2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.

The application made in the planthouse will only provide short-term protection and is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection. Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Transplants must be handled carefully during setting to avoid dislodging treated potting media from roots.

Not all varieties of cucurbit vegetables have been tested for tolerance to Wrangler Insecticide applied to seedling flats. Therefore, treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.

*Use not permitted in California unless otherwise directed by supplemental labeling.

GREENHOUSE VEGETABLES*

(Mature plants in production greenhouses)

Cucumber, Tomato, only

Pests Controlled

Rate
fluid ounces/1000 plants

Aphids

Whiteflies

0.7

Restrictions

Pre-Harvest Interval (PHI): **0 days**

Maximum number of Wrangler Insecticide applications per crop season: **1**

Instructions

Apply specified dosage in a minimum of 16 gallons of water for tomatoes and 21 gallons of water for cucumbers using soil drenches, micro-irrigation, drip irrigation, or hand-held or motorized calibrated irrigation equipment. Do not apply to immature plants since phytotoxicity may occur.

Apply when infestation pressure surpasses threshold and beneficials are not able to maintain pest populations below damage thresholds. Repellency of bumble bee pollinators and negative effects on some beneficials (*Orius* spp.) can occur when Wrangler Insecticide is applied.

Many varieties of vegetables have been tested for tolerance to Wrangler Insecticide and show good safety. However, certain varieties may show more sensitivity to Wrangler Insecticide. Therefore, treat a few plants before treating the whole greenhouse.

*Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

16
33

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

FRUITING VEGETABLES¹

Including: Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento and sweet) Tomato, Pepinos, Tomatillo

Field application instructions. See details below for additional planthouse instructions.

Pests Controlled

	Rate fluid ounces/Acre
Aphids	
Colorado potato beetle	Okra and Pepper
Flea beetles	8.0 – 16.0
Leafhoppers	
Thrips (foliage-feeding thrips, only)	Other Crops
Whiteflies	8.0 – 12.0

Pests/Diseases Suppressed

Symptoms of:	Okra and Pepper
Tomato mottle virus	8.0 – 16.0
Tomato spotted wilt virus	Other Crops
Tomato yellow leaf curl virus	8.0 – 12.0

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum Wrangler Insecticide allowed on pepper and okra crops per application: **16.0 fluid ounces/Acre** (0.50 lb AI/Acre)

Maximum Wrangler Insecticide allowed on other fruiting crops per application: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. In-furrow spray directed on or below seed;
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1-1/2" with sufficient irrigation within 24 hours of application;
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;
5. Post-seeding drench, transplant-water drench, or hill drench;
6. Subsurface side-dress on both sides of each row. Wrangler Insecticide must be incorporated into root-zone.

Planthouse Application Instructions²

Pests Controlled

	Rate fluid ounces/1000 plants
Aphids	
Whiteflies	0.05

Restrictions

Maximum amount of Wrangler Insecticide applied in the planthouse: **0.05 fluid ounces** (0.00156 lb AI)/1000 plants.

Maximum number Wrangler Insecticide applications in planthouse: **1**

Instructions

Apply specified dosage to seedlings in trays in the planthouse, targeting soil media (tray drench), not more than 7 days prior to transplanting, in one of the following manners:

1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash Wrangler Insecticide from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash Wrangler Insecticide from foliage may result in reduced pest control;

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

Fruiting Vegetables¹ cont'd.:

2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.

The application made in the planthouse will only provide short-term protection and is not intended as a substitution for a field application. An additional field application must be made within 2 weeks following transplanting to provide continuous protection.

Applications of higher rates or increased number of applications in planthouse may result in significant plant injury. Transplants must be handled carefully during setting to avoid dislodging treated potted media from roots.

Not all varieties of fruiting vegetables have been tested for tolerance to Wrangler Insecticide applied to seedling flats. Therefore treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse.

¹ Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

² Use not permitted in CA unless otherwise directed by supplemental labeling.

HEAD and STEM BRASSICA VEGETABLES

Including: Broccoli, Broccoli raab (*rapini*), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (*gai lon*) broccoli, Chinese (*bok choy*) cabbage, Chinese (*napa*) cabbage, Chinese mustard (*gai choy*) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens, Turnip tops (leaves)

AND

LEAFY VEGETABLES

Including: Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Cilantro, Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Raddicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian Spinach)), Watercress (commercial production only, applications must not be made to native cress growing in streams or other bodies of water), Watercress (upland)

Pests Controlled

Rate
fluid ounces/Acre
(on 36 Inch rows)

Aphids

Leafhoppers

Thrips (foliage feeding thrips only)

Whiteflies

5.0 – 12.0

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum Wrangler Insecticide allowed per application: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. In-furrow spray directed on or below seed;
3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1-1/2" with sufficient irrigation within 24 hours of application;
4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;
5. Post-seeding drench, transplant-water drench, or hill drench;
6. Subsurface side-dress on both sides of each row. Wrangler Insecticide must be incorporated into root-zone.
7. Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

LEAFY PETIOLE VEGETABLES

Including: Cardoon, Celery, Celtuce, Chinese celery (fresh leaves and stalk only), Florence fennel (including sweet anise, sweet fennel, Finocchio), Rhubarb, Swiss chard

Pests Controlled	Rate Fluid ounces/Acre
Aphids	
Leafhoppers	5.0 – 12.0
Thrips (foliage feeding thrips only)	
Whiteflies	

Restrictions

Pre-Harvest Interval (PHI): **45 days**
Maximum Wrangler Insecticide allowed per application: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

Instructions

- Apply specified dosage of Wrangler Insecticide in one of the following methods:
1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
 2. In-furrow spray directed on or below seed;
 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1-1/2" with sufficient irrigation within 24 hours of application;
 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting;
 5. Post-seeding drench, transplant-water drench, or hill drench;
 6. Subsurface side-dress on both sides of each row. Wrangler Insecticide must be incorporated into root zone.
 7. Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

LEGUME VEGETABLES except soybean, dry

Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean

- Bean** (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin)
Bean (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)
Bean (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)
Pea (*Pisum* spp., includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)
Other Beans and Peas [Broad bean (fava), Chickpea (garbanzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean]

Pests Controlled	Rate fluid ounces/Acre
Aphids	
Leafhoppers	
Thrips (foliage feeding thrips, only)	8.0 – 12.0
Whiteflies	

Pests/Diseases Suppressed

Symptoms of:

Bean common mosaic virus (BCMV)	
Bean golden mosaic virus (BGMV)	8.0 – 12.0
Beet curly top hybrigeminivirus (BCTV)	

Restrictions

Pre-Harvest Interval (PHI): **21 days**
Maximum Wrangler Insecticide allowed per crop season: **12.0 fluid ounces/Acre** (0.38 AI/Acre)

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

Legume Vegetables except soybean, dry cont'd.:

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
2. In-furrow-spray at planting directed on- or below seed;
3. In a narrow (2" or less) surface band over seed-line during planting incorporated to a depth of 1 to 1-1/2" with sufficient irrigation within 24 hours following application;
4. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting;
5. As a post-seeding drench, transplant drench, or hill drench.
6. Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

ROOT VEGETABLES*

Including: Beet (garden) ^{1/}, Burdock (edible) ^{1/}, Carrot ^{1/}, Celeriac ^{1/}, Chervil (turnip-rooted) ^{1/}, Chicory ^{1/}, Ginseng, Horseradish, Parsley (turnip-rooted), Parsnip ^{1/}, Radish ^{1/}, Oriental radish (diakon) ^{1/}, Rutabaga ^{1/}, Salsify (oyster plant), Salsify (black) ^{1/}, Salsify (Spanish), Skirret and Turnip ^{1/}.

Pests Controlled

Rate
fluid ounces/1000 row-feet

Rate
fluid ounces/Acre

Aphids

Flea beetles

0.35 – 0.85

5.0 – 12.0

Leafhoppers

Whiteflies

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum Wrangler Insecticide allowed per crop season: **12.0 fluid ounces/ Acre** (0.38 lb AI/Acre)

Maximum Wrangler Insecticide applications per crop season: **1**

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. Chemigation into root-zone through low pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. In-furrow spray (rate specified per 1000 row-feet) or, shanked-in 1 to 2 inches below seed depth during planting;
3. In a narrow (2 inches or less) band directly (1 to 2 inches) below the eventual seed row in a bedding operation 14 or fewer days before planting.

Important Note: The rate applied affects the length of control. Use higher rates where infestations occur later in crop development, or where pest pressure is continuous. Wrangler Insecticide rates less than 0.7 fluid ounces/ 1000 row-feet will not provide adequate residual pest control. Wrangler Insecticide treated crops grown on very high organic matter soils (muck) may also require additional pest management control.

^{1/} Tops or greens from these crops may be utilized for food or feed.

*Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

20
33

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

TUBEROUS and CORM VEGETABLES*

Including: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Cassava (bitter & sweet) ^{1/}, Chayote (root), Chufa, Dasheen (taro) ^{1/}, Ginger, Leren, Sweetpotato, Tanier (cocoyam) ^{1/}, Turmeric, Yam bean (jicama, manioc pea), Yam (true) ^{1/} (For application instructions on potato, see Field Crops section)

Pests Controlled	Rate fluid ounces/1000 row-feet	Rate fluid ounces/Acre
Aphids		
Flea beetles	0.35 – 0.85	5.0 – 12.0
Leafhoppers		
Thrips (foliage feeding thrips only)		
Whiteflies		

Restrictions

Pre-Harvest Interval (PHI) from planting application: **3 days** (leaves); **125 days** (corms)
Maximum Wrangler Insecticide allowed per crop season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)
Maximum Wrangler Insecticide applications per crop season: **1**

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:
1. In-furrow spray (rate specified per 1000 row-feet) over planting material (hulis) or shanked-in 1 to 2 inches below hulis depth at planting;
2. Side-dress not more than 0.3 fluid ounces/1000 row-feet no later than 45 days after planting. Observe same PHI as above.

Important Note: The rate applied affects the length of control. Use higher rates where infestations occur later in crop development, or where pest pressure is continuous. Wrangler Insecticide rates less than 0.35 fluid ounces/1000 row-feet may not provide adequate residual pest control. Wrangler Insecticide treated crops grown on very high organic matter soils (muck) may also require additional pest management control.

^{1/}Tops or greens from these crops may be utilized for food or feed.
*Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

STRAWBERRY^{1/}
Annual And Perennial Crops

Pests Controlled	Rate fluid ounces/Acre
Aphids	
Whiteflies	12.0 – 16.0

Restrictions

Pre-Harvest Interval (PHI): **14 days**
Maximum Wrangler Insecticide allowed per crop season: **16.0 fluid ounces/Acre** (0.50 lb AI/Acre)

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:
1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment after plants are established or on perennial crops in early spring prior to bud opening;
2. As a plant material or plant hole treatment just prior to, or during transplanting.

The rate applied affects the length of control. Use higher rates where infestations may occur later in crop development or where pest exposure is continuous.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

STRAWBERRY^{1/} / cont'd.:

Post-harvest Use on Perennial Crops

Pests Controlled

Rate
fluid ounces/Acre

White-grub-complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle, Oriental beetle)	8.0 – 12.0
---	------------

Restrictions

Pre-Harvest Interval (PHI): **14 days**

Maximum Wrangler Insecticide allowed per season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

Instructions

Apply a single application **post harvest to coincide with renovation of strawberry fields** and during active egg-laying period of beetles. Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. As a ground spray via boom or backpack sprayer in a minimum of 20 gallons of water per acre;
2. As a row-band spray using an adjusted amount of product based on the treated row band area in proportion to the amount required per full acre. The bandwidth should be equivalent to the width of the anticipated fruiting bed;
3. As a chemigation application with 600 to 1000 gallons of water followed by 0.10 to 0.25 inches irrigation.

Important: All soil-surface applications must be followed by 0.25 inches of rainfall or overhead irrigation water per acre within 2 hours of application. Failure to adequately incorporate Wrangler Insecticide into egg-deposition zone may result in decreased activity of beetle grubs.

^{1/}Do not use both application methods on the same crop in the same season.

SUGAR BEET*

(for use only in CA)

Pests Controlled

Rate
fluid ounces/Acre

Aphids	3.0 – 6.0
Leafhoppers	
Whiteflies	
Flea beetles	

Pests/Diseases Suppressed

Symptoms of:

Western yellows/Beet curly top hybrigeminivirus (BCTV)	3.0 – 6.0
--	-----------

Restrictions

Maximum Wrangler Insecticide allowed per crop season: **6.0 fluid ounces/Acre** (0.18 lb AI/Acre)

Maximum imidacloprid allowed per season: **0.18 lb AI/Acre** (from any formulation) on any row spacing

Instructions

Apply specified dosage of Wrangler Insecticide in the following method:

1. Apply specified dosage in sufficient carrier volume to insure uniform application. Apply directly below each seed furrow either during the bedding operation immediately prior to planting or at the time of planting.

The low rate may be applied to aid establishment of stands in whitefly areas, or for early season control of the other pests listed.

*Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

Wrangler Insecticide Conversion Chart for Linear Application

RATE: fluid ounces/Acre	Rate: fluid ounces/1000 row-feet Based on average row spacing (in inches)							
	10	15	20	25	30	35	40	45
5	0.0475	0.07125	0.095	0.11875	0.1425	0.16625	0.19	0.21375
6	0.057	0.0855	0.114	0.1425	0.171	0.1995	0.228	0.2565
7	0.0665	0.09975	0.133	0.16625	0.1995	0.23275	0.266	0.29925
8	0.076	0.114	0.152	0.19	0.228	0.266	0.304	0.342
9	0.0855	0.12825	0.171	0.21375	0.2565	0.29925	0.342	0.38475
10	0.095	0.145	0.19	0.24	0.285	0.335	0.38	0.43
12	0.115	0.17	0.23	0.285	0.345	0.4	0.46	0.515
14	0.135	0.2	0.27	0.335	0.4	0.47	0.535	0.605
16	0.155	0.23	0.305	0.385	0.46	0.535	0.61	0.69
18	0.17	0.26	0.345	0.43	0.515	0.605	0.69	0.775
20	0.19	0.285	0.38	0.48	0.575	0.67	0.765	0.86
22	0.21	0.315	0.42	0.525	0.63	0.735	0.84	0.945
24	0.23	0.345	0.46	0.575	0.69	0.805	0.92	1.035
26	0.25	0.375	0.495	0.62	0.745	0.87	0.995	1.12
28	0.27	0.4	0.535	0.67	0.805	0.935	1.07	1.205
30	0.285	0.43	0.575	0.715	0.86	1.005	1.145	1.29
32	0.305	0.46	0.61	0.76	0.92	1.07	1.225	1.375

Important Note: The Wrangler Insecticide rate applied affects the length of control and to a considerable extent, the degree of control or effect. Row-spacing X Wrangler Insecticide rate combinations in italics may not provide adequate residual pest control and are not suitable for long-term, residual control. Use higher labeled rates where infestations may occur later in crop development or where pest pressure is continuous. Loveland Products, Inc. offers no warranty for use of Wrangler Insecticide at rates below 0.35 fluid ounces/1000 row-feet.

SOYBEAN

Pests Controlled

Rate
fluid ounces/Acre

- Aphids
- Bean leaf beetle
- Cucumber beetles/Rootworm adults
- Japanese beetle (adults)
- Leafhoppers
- Whiteflies

1.5

Restrictions:

Pre-Harvest Interval (PHI): **7 days**

Minimum interval between applications: **7 days**

Maximum Wrangler Insecticide amount allowed per crop season: **4.5 fluid ounces/Acre** (0.14 lb AI/Acre)

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

TREE, BUSH and VINE CROPS
Application Directions – Wrangler Insecticide

BANANA AND PLANTAIN

Pests Controlled	Rate fluid ounces/Acre
Aphids	8.0 – 16.0
Leafhoppers	
Pests/Diseases Suppressed	
Scales	8.0 – 16.0

Restrictions

Pre-Harvest Interval (PHI): **0 day**

Maximum Wrangler Insecticide allowed per crop season: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

Instructions

Apply specified dosage of this product in the following method:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

BUSHBERRY

Including: Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Ligonberry, Salal

Pests Controlled	Rate fluid ounces/Acre
Japanese beetle (adults, feeding on foliage)	
White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	8.0 – 16.0

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Maximum Wrangler Insecticide allowed per season: **16.0 fluid ounces/Acre** (0.50 lb AI/Acre)

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. 18-inch band on each side of the row followed with 0.25 inches of irrigation immediately after application.

For optimal grub control, apply Wrangler Insecticide to control 1st or 2nd instar larvae. Application may be made post-bloom up to 7 days prior to harvest, or post-harvest until October 1st. For optimum control of Japanese beetle larvae, make applications from June 1 to July 15. Do not apply during bloom.

Application to grass covered rows, row middles, drive lanes, headlands, and other grassy areas in and around the berry field will control resident grub populations. Applications directed to the root-zone will help protect berry plant roots from grub feeding.

Apply Wrangler Insecticide to moist soil. If necessary, apply one hour of irrigation water immediately before application of Wrangler Insecticide. To facilitate movement of Wrangler Insecticide into the soil and root-zone, 1/2 to 1 inch of irrigation water or rainfall must be applied or received within 24 hours of application.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

CANEBERRY

Including

Blackberry (*Rubus eubatus*, including bingleberry, black satin berry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, Loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, youngberry, and varieties and/or hybrids of these) Raspberry (black and red, *Rubus occidentalis*, *Rubus strigosus*, *Rubus idaeus*)

Pests Controlled	Rate fluid ounces/Acre
Aphids	
Leafhoppers	
Whiteflies	8.0 – 16.0
Rednecked cane borer	12.0 – 16.0
Pest Suppressed	
Thrips (foliage feeding thrips only)	8.0 – 16.0

Restrictions:

Pre-Harvest Interval (PHI): **7 days**

Maximum Wrangler Insecticide allowed per season: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

Do not apply pre-bloom or during bloom or when bees are actively foraging.

Applications:

Apply specified dosage in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
2. Basal, soil drench in a minimum of 500 gallons solution per acre.

CITRUS (Containerized) – Soil Treatment

Including: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tagor). Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, White sapote (*Casimiroa* spp.), and other cultivars and/or hybrids of these.

Pests Controlled	Rate mL/ft³ container media
Aphids	
Asian citrus psyllid	
Black fly	
Citrus leafminer	0.375
Leafhoppers/Sharpshooters	
Mealybugs	
Scales	
Whiteflies	
Citrus root weevil (larval complex)	0.625 – 1.25
Pests/Diseases Suppressed	
Citrus thrips (foliage feeding thrips only)	1.25

Instructions

Determine volume of container and calculate dosage necessary to treat container. Apply calculated dosage of Wrangler Insecticide per container as a soil drench or through low-pressure drip or trickle irrigation water. Use sufficient carrier volume to ensure thorough uniform distribution throughout the media without loss of gravitational water from the container. For optimal results, treatment should be made at planting prior to insect infestation. Retreat if necessary. For control of larvae of the citrus root weevil complex, application should be made prior to neonate larvae entering potting media. Utilize higher dosage for heavy infestations.

25
33

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

CITRUS (Field)

Including: Calamondin, Citrus, Citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, White sapote (*Casimiroa* spp.), and other cultivars and/or hybrids of these.

Pests Controlled

Rate
fluid ounces/Acre

Aphids	
Asian citrus psyllid	
Black fly	
Citrus leafminer	8.0 – 16.0
Leafhoppers/Sharpshooters	
Mealybugs	
Scales	
Termites (FL only)	
Whiteflies	

Pests/Diseases Suppressed

Citrus nematode	
Symptoms of:	
Citrus tristeza virus CTV through vector control	
Citrus yellows	16.0
Thrips (foliage feeding thrips only)	

Restrictions

Pre-Harvest Interval (PHI): **0 day**
Maximum Wrangler Insecticide allowed per season **16.0 fluid ounces/Acre** (0.50 lb AI/Acre)

Instructions

- Apply specified dosage of Wrangler Insecticide in one of the following methods:
1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For optimum results, apply to newly planted trees or those previously trained to drip, trickle or micro-sprinkler irrigation. To break soil surface tension, lightly pre-wet soil prior to applications of Wrangler Insecticide. Chemigation application can be made separate to normal irrigation but followed by 10 to 20 minutes of additional watering to move Wrangler Insecticide into root-zone. Allow 24 hours before initiating subsequent irrigations;
 2. Soil surface band spray on both sides of the tree. Overlap bands at the base of the tree to create a continuous band within the drip-line area of the tree, to be followed immediately with light sprinkler irrigation sufficient to move the product into the upper portion of the root-zone. This method is suitable for very coarse soils with 0.75% organic matter or less;
 3. Drench to base of tree not exceeding one-quart total solution per tree immediately around trunk of tree and extending outward covering the entire fibrous root system of the tree. Only suitable for trees up to 8 feet tall;
 4. For control of existing termite infestations, apply specified dosage in 1 to 4 quarts of total solution volume, depending on size of tree, as a drench application to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk.
 5. For suppression of citrus nematode, apply specified dosage through low-pressure chemigation or soil surface band spray only, ensuring complete coverage of the root system and utilizing application directions stated above for the respective application method. Repeated and regular use of Wrangler Insecticide over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response.

26
33

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

COFFEE

Pests Controlled	Rate fluid ounces/Acre
Aphids	8.0 – 16.0
Leafhoppers	
Leafminer	

Pests/Diseases Suppressed

Scales	8.0 – 16.0
--------	------------

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Maximum Wrangler Insecticide allowed per crop season: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

Do not apply pre-bloom or during bloom or when bees are actively foraging.

Instructions

Apply specified dosage in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation;
3. Basal, soil drench in sufficient water to insure incorporation into the root-zone followed by irrigation.

CRANBERRY

Pests Controlled	Rate fluid ounces/Acre
Rootgrubs (Scarabaeidae)	
Rootworms (Chrysomelidae)	8.0 – 16.0

Restrictions

Pre-Harvest Interval (PHI): **30 days**

Maximum Wrangler Insecticide allowed per season: **16.0 fluid ounces/Acre** (0.50 lb AI/Acre)

Do not apply during bloom.

Instructions

Apply Wrangler Insecticide to moist soil. Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. As a soil spray (ground application) directed to the root and crown area using a minimum of 20 gal of water per acre;
2. As a chemigation application with 600 to 1000 gal water.

Immediately upon application, Wrangler Insecticide must be incorporated into root-zone by 0.1 - 0.3 inches water/Acre, either with the chemigation application or through irrigation/rainfall if not applied through chemigation. Inadequate incorporation within 24 hours of application may result in reduced control.

Rootgrubs and Rootworms

Best control may be achieved when application is made post-bloom immediately after bees are removed. Applications should target early instar larvae.

Wrangler Insecticide has not been tested for crop response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of the Wrangler Insecticide and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

GRAPE

Including: American bunch grape, Muscadine grape and Vinifera grape

Pests Controlled

Rate
fluid ounces/Acre

European fruit lecanium

Mealybugs

Leafhoppers/Sharpshooters

8.0 – 16.0

*Phylloxera** spp.

Pests/Diseases Suppressed

Grapeleaf skeletonizer

Nematodes

Pierce's disease

12.0 – 16.0

Restrictions

Pre-Harvest Interval (PHI): **30 days**

Maximum Wrangler Insecticide allowed per season: **16.0 fluid ounces/Acre** (0.50 lb AI/Acre)

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation;
3. Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation.
4. For suppression of nematodes, apply 14 fluid ounces in a single application or two 7 fluid ounce applications on a 30 to 45 day interval. Apply only by 1) chemigation into root-zone through above ground low-pressure drip, trickle, micro-sprinkler, or equivalent equipment; or 2) French plow technique, followed immediately by sufficient irrigation to move the product into the entire root-zone of the plant. Repeated and regular use of Wrangler Insecticide over several consecutive growing seasons provides the greatest degree of nematode suppression and yields the greatest plant response.

For optimal results, make application(s) between bud-break and the pea-berry stage. A total of 14 fluid ounces/Acre is recommended under any of the following conditions:

1. Where vigorous vine growth is expected;
2. In warmer growing areas;
3. Where mealybug and European fruit lecanium populations are expected to be heavy;
4. Where vine populations exceed 600 per acre, or;
5. For suppression of nematodes.

*Repeated and regular use of Wrangler Insecticide over several, consecutive growing seasons controls existing *Phylloxera* infestations over time or prevents *Phylloxera* from becoming established.

HOP

Pests Controlled

Rate
fluid ounces/Acre

Aphids

9.6

Restrictions

Pre-Harvest Interval (PHI): **60 days**

Maximum Wrangler Insecticide allowed per season: **9.6 fluid ounces/Acre** (0.3 lb AI/Acre)

Use not permitted in California unless otherwise directed by supplemental labeling.

Instructions

Apply specified dosage of Wrangler Insecticide in one of the following methods:

1. Chemigation into root-zone through low-pressure drop, trickle, micro-sprinkler or equivalent equipment;
2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation;
3. Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation.

28
33

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

POME FRUIT

Including: Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

Pests Controlled	Rate fluid ounces/Acre
Aphids (including woolly apple aphid)	8.0 – 12.0
Leafhoppers	

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum Wrangler Insecticide allowed per season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

Do not apply pre-bloom or during bloom or when bees are actively foraging.

Instructions

Apply specified dosage of Wrangler Insecticide in the following method:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

POMEGRANATE

Pests Controlled	Rate fluid ounces/Acre
Aphids	8.0 – 16.0
Leafhoppers/Sharpshooters	
Whiteflies	

Restrictions

Pre-Harvest Interval (PHI): **0 day**

Maximum Wrangler Insecticide allowed per crop season: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

Do not apply pre-bloom or during bloom or when bees are actively foraging.

Instructions

Apply specified dosage of this product in the following method:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

STONE FRUIT

Including: Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson, and Japanese), Plumcot, Prune (fresh and dried)

In-field, Soil Application

Pests Controlled	Rate fluid ounces/Acre
Aphids (including woolly apple aphid)	8.0 – 12.0
Leafhoppers	

Restrictions

Pre-Harvest Interval (PHI): **21 days**

Maximum Wrangler Insecticide allowed per season: **12.0 fluid ounces/Acre** (0.38 lb AI/Acre)

Do not apply pre-bloom or during bloom or when bees are actively foraging.

Instructions

Apply specified dosage of Wrangler Insecticide in the following method:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

Pre-plant, Root Dip Application

Pests Controlled	Rate fluid ounces/10 gallons root-dip solution
Black peach aphid (infesting roots)	1.0

Mix Wrangler Insecticide at **1.0 fluid ounces** per 10 gallons of water. Thoroughly wet bare-root transplant to slightly above the graft union by soaking roots in the Wrangler Insecticide solution for up to 5 minutes. Allow solution to dry on roots and transplant trees as soon as possible following treatment.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

TROPICAL FRUIT

Including: Acerola, Atemoya, Avocado, Birida, Black sapote, Canistel, Cherimoya, Custard apple, Feijoa, Guava, Jaboticaba, Llama, Longan, Lychee, Marney sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursap, Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu

Pests Controlled	Rate fluid ounces/Acre
Aphids	
Leafhoppers	12.0 – 16.0
Whiteflies	
Pests/Diseases Suppressed	
Scales	16.0

Restrictions

Pre-Harvest Interval (PHI): **6 days**

Maximum Wrangler Insecticide allowed per application: **16.0 fluid ounces/Acre** (0.50 lb AI/A)

Do not apply pre-bloom or during bloom or when bees are actively foraging.

Instructions

Apply specified dosage of Wrangler Insecticide in the following method:

1. Chemigation through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.

TREE NUTS - Soil Treatment

Including: Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (black and English)

Pests Controlled	Rate fluid ounces/Acre
Aphids	8.0 – 16.0
Leafhoppers/Sharpshooters	
Mealybugs	
Spittlebugs	
Termites	
Whiteflies	

Pests/Diseases Suppressed

Pecan scab (from reduction in honeydew deposition)	12.0 – 16.0
Thrips (foliage-feeding thrips only)	16.0

Restrictions

Pre-Harvest Interval (PHI): **7 days**

Maximum Wrangler Insecticide allowed per crop season: **16.0 fluid ounces/Acre** (0.50 lb AI/Acre)

Do not apply pre-bloom or during bloom or when bees are actively foraging.

Instructions

Apply specified dosage prior to or at onset of pest infestation in one of the following methods:

1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
Pre-wet soil prior to applications of this product and allow soil to dry following application and prior to subsequent irrigation;
2. Emitter or spot application in a minimum of 4 fluid ounces of mixture per emitter site;
3. Shank or subsurface side-dress, injected to a depth just above or just within the root zone and between the trunk and drip line of the tree canopy. Product should be applied in a minimum of 10 gallons per acre using multiple shanks on both sides of trees. Ensure product placement is below sod or orchard floor debris. Irrigation covering entire treated area should follow within 48 hours to promote uptake by root system.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

Tree Nuts - Soil Treatment cont'd.:

4. For control of termites, apply specified dosage to slightly moist soil as a high-volume drench to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk. Utilize sufficient carrier volume to penetrate the soil to a depth of 18 to 24 inches to obtain optimum control. Allow soil to dry following treatment and prior to applying any irrigation.

Use the higher rates when applied by shank or subsurface side-dress, used on larger trees, soils are high in clay content, high plant populations exist, and/or where extended control is desired. Under some conditions, control may not occur for 14 or more days or until two (2) irrigations have been made. Applications made later in the season may result in reduced efficacy.

TREE NUTS - Foliar Treatment

Including: Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (black and English)

Pests Controlled	Rate Fl. oz./A
Aphids (except black pecan aphid)	1.5 to 3.0
Leafhoppers/Sharpshooters	
Phylloxera spp. (leaf infestations)	
Spittlebugs	
Whiteflies	
Black pecan aphid	3.0
Mealybugs	
San Jose scale	

Restrictions:

Pre-Harvest Interval (PHI): **7 days**
 Minimum interval between applications: **6 days**
 Maximum WRANGLER Insecticide allowed per season: **10.4 fluid ounces/Acre** (0.36 lb AI/A)
 Minimum application volume (water); 50 GPA – ground application, 25 GPA – aerial application.
 Do not apply within 10 days prior to bloom or during bloom or when bees are actively foraging.

Applications:

Applications for control of San Jose scale should be timed according to crawler stage, treating each successive generation. Two applications on a 10 to 14-day interval may be required to achieve control.

CHRISTMAS TREE

Pests Controlled	Rate fluid ounces/Acre
White grub complex (damage from grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)	8.0 – 16.0

Restrictions

Maximum Wrangler Insecticide allowed per crop season: **16.0 fluid ounces/Acre** (0.5 lb AI/Acre)

Instructions

Soil incorporation and movement of this product to the root-zone is required for activity. This product can be incorporated most readily when applied to moist soil. Apply specified dosage in one of the following methods:
 1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment;
 2. 18-inch band on each side of the row (small trees) to full broadcast application (large trees) followed by rainfall or 0.25 to 1 inch of irrigation within 12 hours after application.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

Christmas Tree cont'd.:

For optimal grub control, apply this product during adult flight activity, or up to mid-July, when first instar larvae are present.

POPLAR/COTTONWOOD

(Includes members of the genus *Populus* grown for pulp or timber)

Pests Controlled

Rate
fluid ounces/Acre

Aphids

Cottonwood leaf beetle

8.0 – 16.0

Pests/Diseases Suppressed

Phylloxera popularia

8.0 – 16.0

Restrictions

Maximum Wrangler Insecticide allowed at-plant per crop season: **16.0 fluid ounces/Acre** (0.50 lb AI/Acre)

Do not apply pre-bloom or during bloom or when bees are actively foraging.

Instructions

Apply specified dosage of Wrangler Insecticide in the following method:

1. Chemigation through low-pressure drip irrigation.

For Cottonwood leaf beetle, protection against damage will occur when application is made early, when the beetles first begin feeding. Larger trees may require earlier treatment as a result of slower uptake.

2. For narrow-row, cutting orchards/nurseries used for plant propagation, shank into root-zone followed by adequate irrigation to promote uptake. (Adequate irrigation depends on soil moisture level at application. Under dry conditions, use 0.25 inches/Acre).

For *Phylloxera*, apply early in the year, from break of dormancy through May.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org. If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

Storage & Disposal cont'd.:

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 56 gallons: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

WRANGLER® INSECTICIDE
EPA REG. NO. 34704-931

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Actara, Centric and Platinum are registered trademarks of a Syngenta Group Company.
Assail and Intruder are registered trademarks of Nippon Soda Company, LTD.
Calypso, Gaucho, Leverage and Provado are registered trademarks and Trimax is a trademark of Bayer.
Wrangler is a registered trademark of Loveland Products, Inc.

Leverage is a Restricted Use Pesticide.