



34704-941

08-03-2006

13

	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460	EPA Registration Number: 34704-941	Date of Issuance: AUG 3 2006
		NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)	
Name and Address of Registrant (include ZIP Code): Loveland Products PO Box 1286 Greeley, CO 80632-1286		Name of Pesticide Product: Dyna-Shield Fludioxonil	
<small>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA registration number.</small>			
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3 (c) (7) (A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit/cite all data required for the registration/ reregistration of your product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Make the following label changes: <ol style="list-style-type: none"> a. Page one: Change the EPA Registration Number to 34704-941. b. Page one: Ingredient Statement: Move the CAS number for the active ingredient to the line below "Other Ingredients" and refer it to the active ingredient name by means of an asterisk "Fludioxonil *" <p style="text-align: right;">(Continued)</p>			
Signature of Approving Official:  Mary Waller, Product Manager 21 Fungicide Branch Registration Division (7505P)		Date: 8/3/2006	

Continued from first page

EPA Registration. Number 34704-941

c. **Page one: Ingredient Statement:** Line up the words "Fludioxonil", "Other Ingredients" and "Total" so that they line up with "Active Ingredient" along the left margin.

d. **Page one: Ingredient Statement:** Line up the percentages so that the decimal points fall along the same vertical line.

e. **Page two: Under "Directions for Use":**

Change the sentence "Dyna-Shield Fludioxonil should be used only in accordance with recommendations on this label or in separately labeled Loveland Products, Inc. supplemental labeling recommendations for this product." to "Dyna-Shield Fludioxonil must be used only in accordance with recommendations on this label or in EPA-approved supplemental labeling."

Then move this sentence so that it immediately precedes the Agricultural Use Requirements Box.

f. **Page five: Under "Mixing Procedures":**

Reformat this paragraph from:

"This seed has been treated with fludioxonil fungicide. Do not use for feed, food, or oil purposes. Store away from feed and foodstuffs. Forage may not be grazed until 30 days after planting."

Use an EPA-approved dye or colorant that imparts an unnatural color to the seed.
to:

"This seed has been treated with fludioxonil fungicide. Do not use for feed, food, or oil purposes. Store away from feed and foodstuffs. Forage may not be grazed until 30 days after planting. Use an EPA-approved dye or colorant that imparts an unnatural color to the seed."

g. **Page nine: Under "Note for Potatoes Intended for Seed Production:"** change "Amistar at 0.13 fl. oz./1000 ft of row" to "Amistar at 0.13 oz./1000 ft of row".

g. **Page nine: Warranty language**

Change "product=s" to "product's"

Change "The buyer or user of this product assumes all such inherent risks." to "To the extent allowed by state law, the buyer or user of this product assumes all such inherent risks."

3. Submit one (1) copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. If you have any questions, contact Lisa Jones of my staff at (703) 308-9424 or at jones.lisa@epa.gov.

Sincerely,

Mary L. Waller

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosures: Stamped copy of label

4
13

Dyna-Shield™ Fludioxonil

A seed treatment product for protection against certain seed and seedling diseases of crop plants.

ACTIVE INGREDIENT

Fludioxonil (CAS No. 131341-86-1)	40.3%
OTHER INGREDIENTS:	59.7%
Total:	100.0%

Dyna-Shield Fludioxonil contains 4 lbs. active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional Precautionary Statements and Directions For Use inside booklet.

EPA Reg. No. 34704-

EPA Est. No. _____

Net Contents: _____

FIRST AID	
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If in eyes	Hold eye open and rinse slowly. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-301-7976.	

Packaged for:



P.O. Box 1286
Greeley, CO 80632

**ACCEPTED
with COMMENTS
in EPA Letter Dated**

AUG 3 2006

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

34704-
941

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Causes moderate eye irritation. Harmful if absorbed through the skin or inhaled. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart, and
- shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers used closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsates.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Dyna-Shield Fludioxonil should be used only in accordance with recommendations on this label or in separately published Loveland Products, Inc. supplemental labeling recommendations for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls,
- chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton and
- shoes plus socks.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Loveland Products, Inc., no claims are made to guarantee the germination of carry-over seed or propagating material for all crop seed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE: Store in a cool, dry secure place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact you State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional office for guidance in proper disposal methods.

CONTAINER DISPOSAL: Triple rinse (or equivalent), then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or alternatives allowed by state or local authorities.

FOR BULK AND MINIBULK CONTAINERS

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER PRECAUTIONS: Before refilling, inspect thoroughly for damage, such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices. Refill only with Dyna-Shield Fludioxonil. The contents of this container cannot be completely removed by cleaning. Refilling with materials other than Dyna-Shield Fludioxonil will result in contamination and may weaken container. After filling and before transporting, check for leaks. Do not refill or transport damaged or leaking container.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.

GENERAL INFORMATION

Dyna-Shield Fludioxonil is a seed treatment fungicide which provides protection against certain soil-borne and seed-borne diseases of crop plants. Dyna-Shield Fludioxonil is active against *Fusarium*, *Rhizoctonia*, *Helminthosporium*, and weakly pathogenic fungi such as *Aspergillus* and *Penicillium*. When rate ranges are given, use higher rate when disease pressure is expected to be severe.

MIXING PROCEDURES

Apply Dyna-Shield Fludioxonil as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the recommended amount of Dyna-Shield Fludioxonil into the required amount of water for the slurry treater and dilution rate to be used. Follow the manufacturer application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Allow seed to dry before bagging.

Federal law requires that bags containing treated seed shall be labeled with the following information:

"This seed has been treated with fludioxonil fungicide. Do not use for feed, food, or oil purposes. Store away from feed and foodstuffs. Forage may not be grazed until 30 days after planting."

Use an EPA-approved dye or colorant that imparts an unnatural color to the seed. Dyna-Shield Fludioxonil does not control disease caused by *Pythium* spp. or *Phytophthora* spp. if these disease are expected to be a problem, apply Dyna-Shield Fludioxonil tank mixed with Dyna-Shield Mefenoxam. If Dyna-Shield Mefenoxam is not registered on that crop, apply Dyna-Shield Fludioxonil as a tank mix with another labeled seed treatment product that has activity against these diseases. Consult the label for appropriate use rate and follow all use instructions for the products. Follow the label directions for the most restrictive or label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing.

BRASSICA (COLE) LEAFY VEGETABLE GROUP

Broccoli (including Chinese and raab), Brussels sprouts, cabbage (including Chinese bok choy, Chinese napa, and Chinese mustard), cauliflower, cavalo broccola, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, and rape greens. For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 – 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

BULB VEGETABLES

Garlic (including great headed), leek, onion (dry bulb and green), Welch onion, and shallot. For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 – 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed or propagating material.

CEREAL GRAINS (INCLUDING FORAGE AND SILAGE)

Barley, buckwheat, corn (field corn, popcorn, and sweet corn), millet (pearl and proso), oats, rice, rye, sorghum, teosinte, triticale, wheat, and wild rice). **Corn (Field Corn and Popcorn):** For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.034 – 0.068 fl. oz. of Dyna-Shield Fludioxonil per 80,000 kernel count unit of seed or 0.006 to 0.012 mg active ingredient per kernel. Dyna-Shield Fludioxonil contains 14.17 grams active ingredient per ounce.

Rate of Dyna-Shield™ Fludioxonil* mg active per kernel	Rate of Dyna-Shield™ Fludioxonil fl. oz. per 80,000 kernel count unit of seed
0.006 mg	0.034 fl. oz.
0.012 mg	0.068 fl. oz.

* Refer to the General Information section for additional guidance on rate selection.

Apply Dyna-Shield Mefenoxam seed treatment in combination with Dyna-Shield Fludioxonil for protection against *Pythium* spp. Apply Dyna-Shield Mefenoxam at

9
13

0.0425 fl. oz. per 100 lbs. seed or up to 0.004 mg active ingredient per kernel tank mixed with the recommended rates of Dyna-Shield Fludioxonil. Or apply Dyna-Shield Mefenoxam at 0.085 fl. oz. per 100 lbs. of seed or up to 0.008 mg active ingredient per kernel, tank mixed with the recommended rates of Dyna-Shield Fludioxonil. Follow application and other instructions found on the Dyna-Shield Mefenoxam product label.

Grain Sorghum: For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.08 - 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

Apply Dyna-Shield Mefenoxam in combination with Dyna-Shield Fludioxonil for protection against damping-off due to *Pythium* spp. Apply Dyna-Shield Mefenoxam at 0.32 - 0.64 fl. oz. per 100 lbs. of seed tank mixed with recommended rates of Dyna-Shield Fludioxonil.

For systemic downy mildew control on cultivars which possess sorghum downy mildew resistance, apply Dyna-Shield Mefenoxam at 0.64 fl. oz. per 100 lbs. of seed as a tank mix with recommended rates of Dyna-Shield Fludioxonil. To those cultivars which do not possess resistance to a particular race of sorghum downy mildew grown in areas with recent history of severe disease, apply 1.28 fl. oz. of Dyna-Shield Mefenoxam tank mixed with recommended rates of Dyna-Shield Fludioxonil. Follow application and other instructions found on the Dyna-Shield Mefenoxam label.

NOTE: Seed treated with the 1.28 fl. oz. of Dyna-Shield Mefenoxam should be planted and not stored or carried over to the following year. Only high quality sorghum seed ($\geq 90\%$ germ) should be treated with Dyna-Shield products.

Sweet Corn: For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.08 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

Apply Dyna-Shield Mefenoxam seed treatment product in combination with Dyna-Shield Fludioxonil for protection against damping-off due to *Pythium* spp. Apply Dyna-Shield Mefenoxam at 0.32 fl. oz. per 100 lbs. of seed tank mixed with the recommended rates of Dyna-Shield Fludioxonil. For protection against systemic downy mildew of sweet corn, apply Dyna-Shield Mefenoxam at 1.28 fl. oz. per 100 lbs. of seed tank mixed with recommended rates of Dyna-Shield Fludioxonil. Follow application and other instructions found on the Dyna-Shield Mefenoxam product label.

For all other cereal crops: For protection against seed-borne and soil-borne fungi which cause seed decay, damping-off, and seedling blights, apply 0.08 - 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

Important: (1) Do not use treated seed for feed, food, or oil. (2) Forage may not be grazed until 30 days after planting.

CUCURBIT VEGETABLES

Chayote, Chinese waxgourd, citron melon, cucumber, gherkin, gourd edible (includes hyotan, cucuzza, hechima, Chinese okra), momordica, all muskmelon, pumpkin, all squash, and watermelon.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 - 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

12
13

FLAX, COTTON, CRAMBE, JAJOBA, LESQUERELLA, RAPE, SAFFLOWER, SUNFLOWER

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 – 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

FOLIAGE OF LEGUME VEGETABLES

Any cultivar of bean (*Phaseolus* spp.), pea (*Pisum* spp.), and soybean.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 – 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

FRUITING VEGETABLES EXCEPT CUCURBITS

Eggplant, groundcherry, pepino, pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), tomatillo, and tomato.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 – 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

GRASS FORAGE, FODDER AND HAY

All grasses (including turfgrass, bermudagrass, bluegrass, and fescue or bromegrass), Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off and seedling blight, apply 0.08 – 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed or propagating root or tuber material.

Important: Forage may not be grazed until 30 days after planting.

HERBS AND SPICES

All spice, angelica, anise, anise (star), annatto, balm, basil, borage, burnet, chamomile, caper buds, caraway, caraway black, cardamom, cassia bark, cassia buds, catnip, celery seed, chervil, chive, Chinese chive, cinnamon, clary, clove buds, coriander (Chinese parsley or cilantro), costmary, culantro (seed), cumin, curry (leaf), dill seed and dill weed, common fennel and Florence fennel, fenugreek, grains of paradise, horehound, hyssop, juniper berry, lavender, lemongrass, lovage, lovage seed, mace, marigold, marjoram, mustard (seed), nasturtium, nutmeg, parsley (dried), pennyroyal, pepper (black and white), poppy seed, rosemary, rue, saffron, sage, savory (summer and winter), sweet bay, tansy, tarragon, thyme, vanilla, wintergreen, woodruff, wormwood.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 – 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed or propagating root or tuber material.

LEAFY VEGETABLES EXCEPT BRASSICA

Amaranth (leafy, Chinese spinach, tampala), arugula, cardoon, celery (including Chinese), celtuce, chervil, chrysanthemum (edible-leaved and garland), corn salad, cress

(garden and up-land), dandelion, dock, endive, fennel (finocchio), lettuce (head and leaf), orach, parsley, purslane (garden and winter), radicchio, rhubarb, spinach (including New Zealand and vine), and swiss chard.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off and seedling blight, apply 0.08 - 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

LEAVES OF ROOT AND TUBER VEGETABLES

Beet (garden and sugar), burdock (edible), carrot, cassava (bitter and sweet), chervil (turnip rooted), chicory, dasheen (taro), parsnip, radish (including oriental), rutabaga, salsify (black), sweet potato, taniel, turnip and true yam.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 - 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed or propagating root or tuber material.

LEGUME VEGETABLES (SUCCULENT OR DRY)

Bean (*Lupinus* spp. - includes grain lupin, sweet lupin, and white lupin), bean (*Phaseolus* spp. - includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, and wax bean), Bean (*Vigna* spp. - includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, and yardlong bean), broad bean, chickpea, guar, jackbean, lablab bean, lentils, pea (dwarf, edible-pod, English, field, garden, green, snow, sugar snap), pigeon pea, soybean, soybean (immature), and sword bean.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight apply 0.08 - 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed or propagating root or tuber material.

NONGRASS ANIMAL FEEDS (FORAGE, FODDER, STRAW, AND HAY)

Alfalfa, bean (velvet), clover, kudzu, lespedeza, lupin, sainfoin, trefoil, vetch (including crown vetch and milk vetch).

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 - 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

Important: Forage may not be grazed until 30 days after planting.

PEANUTS

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 - 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed.

ROOT AND TUBER VEGETABLES

Arracacha, arrowroot, artichoke (Chinese and Jerusalem), beet (garden and sugar), burdock (edible), canna (edible), carrot, cassava (bitter and sweet), celery root, chayote, chervil, chicory, chufa, dasheen, ginger, ginseng, horseradish, leren, parsley, parsnip, potato (0.08 fl. oz. only), radish (including oriental), rutabaga, salsify (oyster, black, and

12
13

spanish), skirret, sweet potato, fanier, turmeric, turnip, yam bean (jicama, maonic pea), and true yam.

For protection against seed-borne and soil-borne fungi which cause decay, damping-off, and seedling blight, apply 0.08 - 0.16 fl. oz. of Dyna-Shield Fludioxonil per 100 lbs. of seed or propagating root or tuber material.

Note for Application to Potato: Dyna-Shield Fludioxonil applications to potato seed pieces must be made only through specific types of equipment. Please contact your Loveland Products, Inc. seed treatment representative to obtain information on proper equipment for application.

Note for Potatoes Intended for Seed Production: As a resistance management strategy, if Dyna-Shield Fludioxonil is used on potatoes intended for seed production, a labeled rate of mancozeb seed treatment dust must be applied to seed tubers after the Dyna-Shield Fludioxonil treatment. If the mancozeb dust is not used, an in-furrow application of Abound® at 0.40 fl. oz. /1000 ft. of row or Amistar® at 0.13 fl. oz. /1000 ft. of row must be applied over the Dyna-Shield Fludioxonil treated seed tubers.

Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing.

SEED TREATMENT - EXPORT USE

For treatment of the seed for export only to countries requiring treatment with Dyna-Shield Fludioxonil and Dyna-Shield Mefenoxam before importation. User should verify actual registered rate in country of destination and/or use. Seed treated in accordance with the export use pattern must be labeled as follows: TREATED SEED: FOR EXPORT ONLY - NOT FOR DOMESTIC SALE OR USE.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product

13
13

is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS PRODUCT IS SOLD AS IS TO THE EXTENT ALLOWED BY APPLICABLE LAW. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND, TO THE EXTENT REQUIRED BY APPLICABLE LAW, WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, 7251 WEST 4TH STREET, GREELEY, CO 80634.

TO THE EXTENT ALLOWED BY APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT ALLOWED BY APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Amistar® and Abound® are trademarks of a Syngenta Group Company

Dyna-Shield™ is a registered trademark of Loveland Products, Inc.