
 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number: 34704-988	Date of Issuance: 12-11-07
	Term of Issuance: Conditional	
Name of Pesticide Product: Dimension* 0.19% Plus Fertilizer		
Name and Address of Registrant (include ZIP Code): Loveland Products, Inc. P.O. Box 1286 Greeley, CO 80632-1286		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>		
<p>This product is conditionally registered in accordance with FIFRA section 3(c)(7)(a) provided you agree in writing to:</p> <ol style="list-style-type: none"> 1. Add the phrase "EPA Registration No. 34704-988" to the label before you release the product for shipment. 2. To the Environmental Hazards add "rinseate" after "washwaters". 3. To the Storage and Disposal Section change "Storage" to "Pesticide Storage" and "Disposal" to "Pesticide Disposal". 4. Submit to this Agency a copy of the purchasing order or bill of sales showing the source of the product names on the Formulator's Exemption Statement dated September 6, 2007 and with which this product is to be manufactured: prior to manufacturing this pesticide product. 5. Submit one copy of the final printed labeling before you release this product for shipment. 6. Submit and or cite all data required for the registration of this product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for re-registration of this product under FIFRA, section 4. <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance</p>		

2/8

with FIFRA section 6e. Your release for shipment of the product under the enclosed stamped copy of the label constitutes acceptance of these conditions.

Signature of Approving Official:

Date:



12-11-07

Erik Kraft, Acting Product Manager 23
Herbicide Branch, Registration Division (7505P)

You will submit one copy of your final printed labeling before you release the product for shipment. A stamped copy of labeling is enclosed for your records. If you have any questions please contact Erik Kraft at 703-308-9358.



Dimension* 0.19% Plus Fertilizer

Turf and Ornamental Herbicide

For season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf, including golf course fairways, roughs and tee boxes.

- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York, this product may be applied only by commercial applicators at no more than 263 lb (0.5 lb active ingredient) per acre (6.0 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

ACTIVE INGREDIENT

dithiopyr: 3,5-pyridinedicarboethioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester 0.19%

INERT INGREDIENTS 99.81%
TOTAL 100.00%

This product contains 0.095 pound of the active ingredient dithiopyr per 50 pound bag. Product protected by U.S. Patent No. 4,692,184. Other patents pending.

*Trademark of Dow Agrosciences LLC

Notice: Read the entire label. Use only according to label directions. Before using this product, read CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY. If terms are unacceptable, return at once unopened. Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Guaranteed Fertilizer Analysis: XX-XX-XX

Water-Insoluble Nitrogen	XX%
Total Nitrogen (N)	XX%
Ammoniacal Nitrogen	XX%
Urea Nitrogen	XX%
Available Phosphoric Acid (P ₂ O ₅)	XX%
Soluble Potash (K ₂ O)	XX%
Calcium (Ca)	XX%
Magnesium (Mg)	XX%
Copper (Cu)	XX%
Iron (Fe)	XX%
Manganese (Mn)	XX%
Zinc (Zn)	XX%

† This product may be formulated with one or more of the following fertilizer materials: urea, methylene urea, sulfur coated urea, ammoniated phosphates, sulfate of potash, muriate of potash, copper oxide, iron oxide, manganese oxide, zinc oxide or limestone.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If in eyes:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled:	• Move person to fresh air if person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL:
1-800-301-7976.

ACCEPTED
with COMMENTS
in EPA Letter Dated

12-11-07

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

31704-987

EPA REG. NO. 34704-XXX
EPA EST. NO. 34704-MS-001
NET WEIGHT 50 LBS. (22.68 KG)

09/07

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes Eye Irritation • Harmful if Inhaled
Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal. **STORAGE:** Store this product only in its original container in a dry, cool, secured storage area. **DISPOSAL:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures. **EMPTY CONTAINER DISPOSAL:** Do not reuse empty bag. Dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **SPILL:** In case of spill, sweep up material and dispose of material according to "product disposal" directions listed above.

USE DIRECTIONS FOR TURF

This turf and ornamental herbicide provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals" elsewhere in this label.

Use Precautions for Turf

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. Initial application of this product should be made only after turfgrasses have developed a vigorous root system, a uniform stand, and have received at least two mowings following seeding, sodding, or sprigging. Use of this product on turfgrasses that are not well-established, or weakened by stress from weather, pests, disease, chemical, or mechanical influences, may increase the chances of turf injury.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

Cool-Season Grasses
bentgrass, creeping†
bluegrass, Kentucky
fescue, fine††
fescue, tall
ryegrass, perennial

Agrostis palustris
Poa pratensis
Festuca rubra
Festuca arundinacea
Lolium perenne

4/8

SIGNATURE BRA FERTILIZER DIMENSION* 0.19% PLUS FERTILIZER EPA REG. NO. 34704-XXX

Warm-Season Grasses

Bahiagrass
Bermudagrass
Buffalograss††
Carpentergrass
Centipedeagrass
Kikuyagrass
St. Augustinegrass
Zoysiagrass

Paspalum notatum
Cynodon dactylon
Buchloe dactyloides
Axonopus affinis
Eremochloa ophiuroides
Pennisetum clandestinum
Stenolaphrum secundatum
Zoysia japonica

† Use of this product on certain varieties of creeping bentgrass, such as 'cohansey', 'carmen', 'seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. Do not apply this product to colonial bentgrass (*Agrostis tenuis*).

†† Use of this product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Ailanta', 'banner', 'beauty', 'bilgart', 'CF-2', 'enjoy', 'HF-93', 'highlight', 'ivalo', 'Jamestown', 'koker', 'majenta', 'mary', 'pennlawn', 'tamara', 'tatjana', 'wadorf', and 'waldina'. Not all varieties of fine fescue have been tested.

††† Do not use this product on seeding buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Note: It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response of turfgrass varieties not listed on this label on a limited area, at recommended use rates, prior to initiating large-scale use.

Precautions for Reseeding, Overseeding, or Sprigging

- Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 18 weeks after a split application program totaling 6.0 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses.
- When reseeding or overseeding, follow established cultural practices for seedbed preparation, seeding, irrigation and fertilization.
- For best overseeding results, use mechanical or power seeding equipment (silt seeders) designed to provide good contact between soil and seed.

Precautions for Control of Crabgrass

Early postemergence weed control with this product is limited to crabgrass. Postemergence application effectively controls crabgrass, but only if application is made prior to the 3-leaf stage of growth or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

Other Use Precautions

- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not allow people or pets to enter treated area until dust has settled.
- Do not graze livestock or feed forage cut from areas treated with this product.

Application Directions for Turf

Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by ½ inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
- Use of split (half-rate) applications spaced 5 - 10 weeks apart may provide improved weed control.

Maximum Use Rates (Applies to all Turfgrass Uses)

- Do not apply more than 263 lb (0.5 lb active ingredient) per acre (6.0 lb per 1000 sq ft) per application or more than 790 lb (1.5 lb active ingredient) per acre (18.0 lb per 1000 sq ft) per year if using split or sequential applications.
- In the state of New York, this product may be applied only by commercial applicators at no more than 263 lb (0.5 lb active ingredient) per acre (6.0 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

Control of Crabgrass

Preemergence and Early Postemergence Control

- This product provides "preemergence" control of crabgrass in established turfgrasses (including large, smooth, and southern species) when applied prior to germination and emergence of crabgrass.
- This product also provides "early postemergence" control of crabgrass if applied prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. The practical benefit of early postemergence crabgrass control is a 2 - 4 week longer period of time (window of application) during which crabgrass control can be achieved.

Application Frequency Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall as described in the following use programs.

Spring Applications.

For applications made in the spring or early summer, use an appropriate application rate corresponding to one of the three control programs listed in Table 1. Specific rate recommendations are based on the user's location (Region), the mowing height of the turfgrass, and whether the application timing is considered to be preemergence or early postemergence. The duration of residual weed control provided by this product is directly related to application rate, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where turfgrass is cut relatively high (e.g., homeowner lawns). This program provides 3 - 5 months of preemergence crabgrass control. This program will also provide early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g., homeowner lawns).

Use Program 2 for preemergence control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways), and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4 - 6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g., residential lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut relatively low (e.g., golf fairways) and (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4 - 6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively low (e.g., golf fairways).

Table 1 Recommended Use Rates

Region	Application Rates	Program 1	Program 2	Program 3
All states, except NY, and parts of states not included in transition, south, coastal south or west.	lb/acre lb/1000 sq ft lb a/acre	100 2.3 0.19	132 3.0 0.25	200 4.6 0.38
Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, & RI	lb/acre lb/1000 sq ft lb a/acre	132 3.0 0.25	200 4.6 0.38	263 6.0 0.5
South: AL, AR, GA, LA, MS, NC, NM, OK, SC, TN, & TX	lb/acre lb/1000 sq ft lb a/acre	200 4.6 0.38	132 + 132 3.0 + 3.0† 0.25 + 0.25	163 + 163 3.7 + 3.7† 0.31 + 0.31
Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX	lb/acre lb/1000 sq ft lb a/acre	132 + 132 3.0 + 3.0† 0.25 + 0.25	163 + 163 3.7 + 3.7† 0.31 + 0.31	200 + 200 4.6 + 4.6† 0.38 + 0.38
West: AZ, CA, & NV. In this climatically diverse region, use the higher rates in local areas with longer crabgrass seasons.	lb/acre lb/1000 sq ft lb a/acre	100 + 132 2.3 + 3.0 0.19 + 0.25	132 + 200 3.0 + 4.6 0.25 + 0.38	132 + 132 3.0 + 3.0† 0.25 + 0.25

† Split applications may be made 5 - 10 weeks apart.

Use of Split or Sequential Applications: Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. Split applications may be made with the rates in Table 1 being split between two applications made 5-10 weeks apart. Where longer periods of control are required, sequential applications may be made in late summer or fall following a spring application or in the spring following a fall application, provided the maximum application rate per year is not exceeded. Refer to the "Maximum Use Rates" section above and separate rate restrictions for the state of New York.

Fall Applications.

This product can also be applied in the late summer through early fall (late August through November) at use rates listed in Table 1 for "Program 3" to provide control of crabgrass through the early part of the next spring. Fall application of this product may be followed by an appropriately timed application in the following spring for season-long control of crabgrass.

Weeds Controlled or Suppressed

Spring Applications

When applied preemergence (prior to germination) for control of crabgrass in the spring, this product will also control or suppress the following weeds at application rates given in the Table 1 above:

5/8

**SIGNATURE BRA FERTILIZER
DIMENSION* 0.19% PLUS FERTILIZER
EPA REG. NO. 34704-XXX**

Grasses

barley
barnyardgrass
bluegrass, annual
brome
crabgrass, large
crabgrass, smooth
crabgrass, Southern
crowfootgrass
dallisgrass (seeding)
goosegrass
foxtail, green
foxtail, yellow
kikuyugrass†
oats, wild
ryegrass
(annual & perennial)
sandbur
smutgrass

Hordeum spp.
Echinochloa crus-galli
Poa annua
Bromus spp.
Digitaria sanguinalis
Digitaria ischaemum
Digitaria ciliaris
Dactyloctenium aegyptium
Paspalum dilatatum
Eleusine indica
Setaria viridis
Setaria pumila
Pennisetum clandestinum
Avena fatua
Lolium spp.

Cenchrus spp.
Sporobolus indicus

Broadleaf Weeds

bittercrass†
carpetweed
chickweed†
geranium, Carolina†
henbit
knotweed, prostrate
lespedeza, common†
marestail
medic, black
mustard
oxalis, buttercup
pineappleweed†
pigweed, redroot
parsley-piert†
purslane, common
rocket, London
shepherdspurge
speedwell, corn†
spurge, garden
spurge, prostrate
spurge, spotted
woodsorrel, creeping
woodsorrel, yellow

Cardamine spp.
Mollugo verticillata
Stellaria spp.
Geranium carolinianum
Lamium spp.
Polygonum aviculare
Lespedeza striata
Coryza canadensis
Medicago lupulina
Brassica spp.
Oxalis pes-caprae
Matricaria matricarioides
Amaranthus retroflexus
Achermilla arvensis
Portulaca oleracea
Sisymbrium irio
Capsella bursa-pastoris
Veronica arvensis
Euphorbia hirta
Euphorbia humifusa
Euphorbia maculata
Oxalis corniculata
Oxalis stricta

†Suppression only

Fall Applications

When applied preemergence (prior to germination) in late summer through early fall (late August through November) this product will control or suppress the following weeds in addition to providing preemergence control of crabgrass in the following spring.

bluegrass, annual
bittercrass
chickweed
henbit
geranium, Carolina
parsley-piert
pineappleweed
shepherdspurge

Poa annua
Cardamine spp.
Stellaria spp.
Lamium spp.
Geranium carolinianum
Achermilla arvensis
Matricaria matricarioides
Capsella bursa-pastoris

Application Instructions and Equipment

This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application.

Check equipment frequently to verify calibration and proper functioning.

USE DIRECTIONS FOR LANDSCAPE ORNAMENTALS

This product provides preemergence control or suppression of listed annual grass and broadleaf weeds in plantings of ornamental plants listed on this label. This product may be applied in ornamental gardens, parks, golf courses and residential areas where ornamental plants are grown for aesthetic purposes. Apply this product with a properly calibrated spreader that will provide

uniform particle distribution. Follow instructions in the "Application Instructions and Equipment" section.

Use Rates

This product may be applied as a single application or split application. Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing seasons. To make a split application, divide the rate in the following table into separate applications made 5-10 weeks apart. Sequential applications may be made in the late summer following a spring application or in the spring after a fall application, provided maximum application rates per year given below are not exceeded.

Pounds of Product/Acre	Pounds of Product/1000 sq ft	Pounds of Product/100 sq ft†
263	6.0	0.6 (9.6 oz)

†Rate based on an area of 100 sq ft for treatment of small areas.

Maximum Use Rates

- Do not apply more than 263 lb (0.5 lb active ingredient) per acre (6.0 lb per 1000 sq ft) per application or more than 790 lb (1.5 lb active ingredient) per acre (18.0 lb per 1000 sq ft) per year if using split or sequential applications.
- In the state of New York, this product may be applied only by commercial applicators at no more than 263 lb (0.5 lb active ingredient) per acre (6.0 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

Application Timing and Recommendations

This product is effective as a preemergence herbicide, but will not control established weeds. Applications to mulched areas or bare ground must be made prior to weed seed germination. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Existing weedy vegetation may be controlled by hand weeding, cultivation or using postemergence herbicides prior to application.

Tips for Improved Weed Control

- The herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by ½ inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
- Avoid disturbance of treated areas. Loss of weed control may result if the treated soil surface is disturbed by soil mixing or tillage.

Use Precautions

- Apply this product only to established ornamentals.
- Do not apply this product directly to bare roots of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress resulting from drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides, or damage due to insects, nematodes or disease.
- To avoid foliar damage to ornamentals, irrigate immediately after application to remove any herbicide granules adhering to foliage.
- Do not apply under conditions that would affect uniformity of application or distribution on the soil surface. Uneven product distribution will result in uneven weed control. Application under windy conditions can result in uneven distribution or cause herbicide granules to drift from the intended treatment area.
- Do not graze livestock or feed forage cut from areas treated with this product.

Weeds Controlled or Suppressed

Used as directed, this product will control crabgrass and control or suppress other listed grass and broadleaf weeds when applied prior to their germination. Refer to "Use Directions for Turf" section for a complete listing of weeds controlled or suppressed. This product will not control established broadleaf weeds or grasses, except for crabgrass in early stages of development. The area to be treated should be free of weeds prior to application.

Tolerant Ornamentals

The ornamentals listed below have shown tolerance to this product when applied according to directions in this label. However, this product has not been tested on all ornamental species, all cultivars of species on this label, or under all possible growing conditions. For species or cultivars not listed, it is recommended that a few plants in a limited area be treated and observed for tolerance under local growing conditions prior to large-scale use.

Common Name	Botanical Name	Tolerant Cultivars
Abelia, Dwarf	<i>Abelia X grandiflora</i>	Nana
Ajuga	<i>Ajuga reptans</i>	Bronze
	<i>Ajuga genevensis</i>	Bronze Beauty
Almond, Flowering	<i>Prunus glandulosa</i>	
Apple, paradise†	<i>Malus pumila</i>	
Arborvitae	<i>Thuja occidentalis</i>	Nigra Pyramidalis Smaragh Techny Woodwardii
Arborvitae, Golden	<i>Thuja orientalis</i>	

6/8

SIGNATURE BRA FERTILIZER
DIMENSION* 0.19% PLUS FERTILIZER
EPA REG. NO. 34704-XXX

Common Name	Botanical Name	Tolerant Cultivars
Aster, Chinese	<i>Callistephus chinensis</i>	Dwarf Queen
Ash, Green	<i>Fraxinus pennsylvanica</i>	
Ash, Mountain	<i>Sorbus aucuparia</i>	
Ash, White	<i>Fraxinus americana</i>	
Azalea	<i>Rhododendron</i> spp.	Brilliant Buccaneer Carror Chimes (Belgian) Elsie Lee Exbury Fashion Hardizer Beauty Hershey Red Higasa Hincrimson Holland (Hybrid) Marion Lee Northern Lights Orange Cup Orchid Lights Snow Southern Charm
Azalea, Flame	<i>Rhododendron calendulaceum</i>	
Azalea, Kirishima	<i>Rhododendron obtusum</i>	
Bamboo, Heavenly	<i>Nandina domestica</i>	
Barberry	<i>Berberis thunbergii</i>	Aurea Dwarf Pigmy Green Kobold Pygmy Red Rose Glow Atropurpurea
Barberry, Purple	<i>Berberis vulgaris atropurpurea</i>	
Banket Flower	<i>Gaillardia grandiflora</i>	Massachusetts
Bearberry (common)	<i>Arctostaphylos uva-ursi</i>	
Beebalm	<i>Monarda didyma</i>	
Begonia	<i>Begonia</i> spp.	
Birch, River	<i>Betula nigra</i>	
Blackeyed Susan	<i>Rudbeckia hirta</i>	Goldstrum
Blanket Flower	<i>Gaillardia</i> spp.	
Blueberry	<i>Vaccinium</i> spp.	Bluecrop Blue Jay Jersey North Blue Northland
Bottlebrush	<i>Callistemon citrinus</i>	
Boxwood, Japanese	<i>Buxus sinica</i>	Japonica
Boxwood, Weller	<i>Buxus sempervirens</i>	
Broom	<i>Cytisus</i> spp.	
Silkyleaf woadwaxen	<i>Genista pilosa</i>	Moonlight
Bugle, Carpet	<i>Ajuga reptans</i>	
Camellia	<i>Camellia japonica</i>	Debutante Mathotiana Supreme Chansonette
Candy Tuft	<i>Camellia sasanqua</i>	
Carex, Variegated	<i>Iberis</i> spp.	Snow White
Cedar, Red	<i>Carex</i> spp.	
Celostia	<i>Juniperus virginiana</i>	
Celastrum	<i>Celosia</i> spp.	
Centaurea	<i>Centaurea montana</i>	
Cockscomb, Plumosa	<i>Celosia cristata</i>	Scarlet Plumosa
Coleus	<i>Coleus blumei</i>	Red Kewpie
Columbine	<i>Aquilegia</i> spp.	
Copper leaf	<i>Acalypha wilkesiana</i>	
Coreopsis	<i>Coreopsis</i> spp.	
Corn Flower	<i>Centaurea</i> spp.	Moonbeam
Cranberry Cotoneaster	<i>Cotoneaster apiculatus</i>	
Coyotebrush	<i>Baccharis pilularis</i>	
Cycads	<i>Cycas revoluta</i>	
Cypress, Bald	<i>Taxodium distichum</i>	
Cypress, Italian	<i>Cupressus sempervirens</i>	Glaucia Gracilis False
Cypress, Japanese	<i>Chamaecyparis obtusa</i>	
Cypress, Leyland	<i>Cupressocyparis leylandii</i>	
Daffodil	<i>Narcissus</i> spp.	King Alfred Aztec Gold Bright Yellow (Hybrid) Single Gold (Evergreen) Wilsonis Yellow
Daylily	<i>Hemerocallis</i> spp.	
Dianthus (Sweet William)	<i>Dianthus</i> spp.	
Delphinium	<i>Delphinium</i> spp.	Magic fountain
Dogwood, flowering	<i>Cornus florida</i>	
Dogwood, Redosier	<i>Cornus sericea</i>	Flavaramaea
Douglas Fir	<i>Pseudotsuga menziesii</i>	
Dusty Miller	<i>Senecio cineraria</i>	

Common Name	Botanical Name	Tolerant Cultivars
Elm, Drake	<i>Ulmus</i>	
Euonymus	<i>Euonymus fortunei</i>	Argenteo-variegata Aureo-marginata Colorata Emerald Gaiety Emerald 'n Gold Gold Edge Gold Princess Silver King Tricolor Vegetus
Fan Palm, European	<i>Chamaerops humilis</i>	
Fan Palm, Mexican	<i>Washingtonia robusta</i>	
Fern (various)	<i>Asparagus</i> spp.	Rainbow Nitida
Fescue	<i>Festuca glauca</i>	
Fetterbush	<i>Leucothoe fontanesiana</i>	
Ficus	<i>Ficus retusa</i>	
Fir Fraser	<i>Abies fraseri</i>	Arnold Dwarf Bronxensis Dwarf Lynwood Gold Meadowlark Weeping
Forsythia	<i>Forsythia X intermedia</i>	
Fuchsia	<i>Spring Glory</i>	
Gallium	<i>Fuchsia</i> spp.	
Gardenia	<i>Galium odoratum</i>	
Geranium	<i>Gardenia jasminoides</i>	Mystery Radicans
Gum	<i>Palargonium X hortorum</i>	
Hawthorn	<i>Eucalyptus citriodora</i>	
	<i>Crataegus</i> spp.	Cockspur White Crimson Cloud Enchantress Jack Evans Washington White Mediterranean Pink
Heather, Twisted	<i>Erica cinerea</i>	
Hemlock, Canada	<i>Tsuga canadensis</i>	
Hibiscus	<i>Hibiscus</i> spp.	Blue Bird Brilliant Hula Girl Blue Boy Blue Girl Burdott China Girl Compacta Forsteri Hellerie Japanese Northern Beauty Needlepoint Nellie R. Stevens Savannah
Holly	<i>Ilex</i> spp.	
	<i>Ilex X meserveae</i>	
Holly, Chinese	<i>Ilex X attenuate</i>	
Holly, Japanese	<i>Ilex cornuta</i>	
Holly, Yaupon	<i>Ilex crenata</i>	
Honeysuckle	<i>Ilex vomitoria</i>	
	<i>Lonicera japonica</i>	Claveys Dwarf Halliana Tatarian Canadian White Zebelli Red Hosta Albo Marginata
Hosta	<i>Hosta</i> spp.	
Ice Plant	<i>Hosta lancifolia</i>	
Impatiens	<i>Carpobrotus edulis</i>	
Iris	<i>Impatiens</i> spp.	
	<i>Iris</i> spp.	Dwarf Blue Wedgewood Bulgaria Thorndale
Ivy, English	<i>Hedera helix</i>	
Jasmine, Asian	<i>Trachelospermum asiaticum</i>	
Juniper	<i>Juniperus</i> spp.	Arcadia Armstrong Bar Harbor Blue Chip Blue Pacific Blue Rug Blue Star Broadmoor Buffalo Calgary Carpet Emerald Sea Emerald Spreader Andorra Compacta Fruitland Green Gold Tip Hetzi Hughes Manhattan Blue Parsoni Pfitzeriana Plumosa Prince of Wales Procumbens Dwarf
	<i>Juniperus horizontalis</i>	
	<i>Juniperus conferta</i>	
	<i>Juniperus squamata</i>	
	<i>Juniperus sabina</i>	
	<i>Juniperus conferta</i>	
	<i>Juniperus horizontalis</i>	
	<i>Juniperus chinensis</i>	
	<i>Juniperus hetzii</i>	
	<i>Juniperus horizontalis</i>	
	<i>Juniperus virginiana</i>	
	<i>Juniperus davurica</i>	
	<i>Juniperus x media</i>	
	<i>Juniperus horizontalis</i>	
	<i>Juniperus procumbens</i>	

SIGNATURE BRAND FERTILIZER
DIMENSION* 0.19% PLUS FERTILIZER
EPA REG. NO. 34704-XXX

Common Name	Botanical Name	Tolerant Cultivars
Juniper cont'd.:	<i>Juniperus sargentii</i>	San Jose Sargent Blue Sargent Green Scandia Scopulorum Moonglow Scopulorum Skyrocket Spartan Tamariscifolia Weberi Youngstown Yukon Belle
King Palm	<i>Archontophoenix cunninghamiana</i>	
Laurel, Australian	<i>Pittosporum tobira</i>	
Laurel, Mountain	<i>Kalmia latifolia</i>	
Leucothoe	<i>Leucothoe fontanesiana</i>	
Ligustrum, Japanese	<i>Ligustrum japonicum</i>	
Lily, African	<i>Agapanthus africanus</i>	Albus Peter Pan
Lily, African Blue	<i>Pieris japonica</i>	Mt. Fire
Lily of the Valley	<i>Lilium muscarum</i>	Evergreen Giant
Lilyturf		Lilac Beauty Majestic Monroe White Variegata
Liriope, Green	<i>Liriope spicata</i>	
Maple, Japanese	<i>Acer japonicum</i>	
Maple, Norway	<i>Acer platanoides</i>	
Maple, Red†	<i>Acer rubrum</i>	
Maple, Silver†	<i>Acer saccharinum</i>	
Maple Sugar†	<i>Acer saccharum</i>	
Marigold	<i>Tagetes patula</i>	Honeycomb Variegata Wheeler Dwarf Golden Snowflake Double White
Mock Orange†	<i>Philadelphus</i> spp.	Sunnyside Faurei Langer Muskogee Standard Pink
Mondo Grass	<i>Ophiopogon japonicus</i>	
Moss Rose	<i>Portulaca grandiflora</i>	
Myrtle, Crape	<i>Lagerstroemia indica</i>	
Myrtle, Wax	<i>Myrica cerifera</i>	
Nandina	<i>Nandina domestica</i>	Compacta Nana
Narcissus	<i>Narcissus</i> spp.	
Oak, Laurel	<i>Quercus laurifolia</i>	
Oak, Pin	<i>Quercus palustris</i>	
Oak, Red	<i>Quercus rubra</i>	
Oak, Southern	<i>Quercus virginiana</i>	
Oak, Willow	<i>Quercus phellos</i>	
Oleander	<i>Nerium oleander</i>	Hardy Red Petite Pink Sister Agnes Wirtlig
Osteospermum	<i>Osteospermum fruticosum</i>	
Pachysandra	<i>Pachysandra terminalis</i>	
Palm, Bangalaw	<i>Archontophoenix cunninghamiana</i>	
Pampas Grass	<i>Cortaderia selloana</i>	
Pansy	<i>Viola x wittrockiana</i>	
Paper Flower	<i>Bougainvillea glabra</i>	Barbara Karst
Peach†	<i>Prunus persica</i>	
Periwinkle, Dwarf	<i>Vinca minor</i>	
Petunia	<i>Petunia X hybrida</i>	Picotti
Photinia, Red Tip	<i>Photinia X fraseri</i>	
Pieris	<i>Pieris japonica</i>	
Pine, Australian	<i>Pinus nigra</i>	
Pine, Japanese Black	<i>Pinus thunbergiana</i>	
Pine, Loblolly	<i>Pinus taeda</i>	
Pine, Longleaf	<i>Pinus palustris</i>	
Pine, Mugo	<i>Pinus mugo</i>	
Pine, Scotch	<i>Pinus sylvestris</i>	
Pine, Slash	<i>Pinus elliotii</i>	
Pine, Swiss Mt.	<i>Pinus mugo</i>	
Pine, Virginia	<i>Pinus virginiana</i>	
Pine, Eastern White	<i>Pinus strobus</i>	
Pineapple, Guava†	<i>Feijoa sellowiana</i>	
Pieris	<i>Pieris taiwanensis</i>	
Pittosporum, Japanese	<i>Pittosporum tobira</i>	Abbotswood
Potentilla	<i>Potentilla nepalensis</i>	
Bush cinquefoil	<i>Potentilla fruticosa</i>	
Privet, Japanese	<i>Ligustrum japonicum</i>	Golden Vicary Regal Texanum Wax Yellow Tipped

Common Name	Botanical Name	Tolerant Cultivars
Privet, Glossy	<i>Ligustrum lucidum</i>	Gnome Lalandei Victory
Pyreantha	<i>Pyreantha koidzumii</i>	
Queen Palm	<i>Arecastrum ramanzottianum</i>	
Quince, Japanese†		
Rhododendron	<i>Rhododendron</i> spp.	Album Cunningham White Fashion Hardy PJM Purple Gem Silvery Pink
Rhododendron, Carolina	<i>Rhododendron carolinianum</i>	
Rhododendron, Catawba	<i>Rhododendron catawbiense</i>	
Rockcress	<i>Arabis</i> spp.	Snowcap
Rhodie Max (Rosebay)	<i>Rhododendron maximum</i>	
Rose†	<i>Rosa banksiae</i>	Lutea
Rosemary†	<i>Rosmarinus officinalis</i>	
Rosemary, Bog	<i>Andromeda polifolia</i>	Nana
Salvia	<i>Salvia farinacea</i>	Rhea
Sedum	<i>Sedum spurium</i>	Dragon Blood Red Red Carpet Yellow Vancouver Gold
Silkleaf woodwaxen	<i>Genista plicosa</i>	
Snapdragon	<i>Antirrhinum</i> spp.	
Sourwood	<i>Oxydendrum arboreum</i>	
False Spiraea	<i>Astilbe X arendsii</i>	
Spiraea	<i>Spiraea</i> spp.	Fanall Anthony Waterer Red Dochica Froebel Pink Goldenflame Red Snowmound White Van Houtte White
Spiraea, Garland	<i>Spiraea</i> spp.	
Spruce, Black Hills	<i>Picea glauca</i>	
Spruce, Colorado Blue	<i>Picea pungens</i>	
Spruce, Norway	<i>Picea abies</i>	
Spruce, White	<i>Picea glauca</i>	Conica
Sweetflag, Grassy-Leaved	<i>Acorus gramineus</i>	
Sweetgum	<i>Liquidambar styraciflua</i>	
Sycamore	<i>Platanus occidentalis</i>	
Tree Fern (Tiki Fern)	<i>Asparagus virgatus</i>	
Trumpet Flower, Evening	<i>Gelsemium sempervirens</i>	
Tulip	<i>Tulip</i> spp.	Apeldoorn
Verbena, Shrub	<i>Lantana sellowiana</i>	
Vervain	<i>Verbena</i> spp.	St. Paul
Viburnum	<i>Viburnum</i> spp.	American Cranberry Bush Arrowwood Common Snowball European Cranberry Bush Linden Mohican Wright
Vinca (Periwinkle)	<i>Vinca minor</i>	
Windmill Palm	<i>Trachycarpus fortunei</i>	
Xylosma	<i>Xylosma congestum</i>	
Yarrow	<i>Achillea</i> spp.	
Yasupon	<i>Ilex vomitoria</i>	Dwarf
Yew	<i>Taxus cuspidata</i> <i>Taxus X media</i>	Densiformis

†Ornamental species only: Do not use on food producing trees and ornamentals.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY BEFORE BUYING OR USING THIS PRODUCT. read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is

8/8

**SIGNATURE BRA FERTILIZER
DIMENSION* 0.19% PLUS FERTILIZER
EPA REG. NO. 34704-XXX**

reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL, THIS PRODUCT IS SOLD AS IS TO THE EXTENT ALLOWED BY APPLICABLE LAW. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL, BUYER OR USER MUST SEND, TO THE EXTENT REQUIRED BY APPLICABLE LAW, WRITTEN NOTICE OF SUCH CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, 7251 WEST 4TH STREET, GREELEY, CO 80634.

TO THE EXTENT ALLOWED BY APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT TO THE EXTENT ALLOWED BY APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

FORMULATED FOR



P.O. BOX 1286, GREELEY, COLORADO 80632-1286