



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 26, 2016

Dr. Chris Mason
Loveland Products Inc.
P.O. Box 1286
Greeley, CO 80632-1286

Subject: Notification per PRN 98-10 – Add referral statement to look inside the label booklet for Precautionary Statements, First Aid, Directions for Use and Storage and Disposal statements
Product Name: Whiteout 2,4-D
EPA Registration Number: 34704-1032
Application Date: 02/12/2016
Decision Number: 514332

Dear Dr. Mason:


The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 34704-1032
Decision No. 514332

If you have any questions, you may contact Karen Samek at (703) 347-8825 or via email at samek.karen@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long horizontal line extending from the top of the "g" in "Montague".

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs



WHITEOUT 2,4-D

GROUP 4 HERBICIDE

NOTIFICATION

34704-1032

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

05/26/2016

CONTAINS NO SOLVENT

ACTIVE INGREDIENT:

2-ethylhexyl ester of 2,4-dichlorophenoxyacetic acid* **60.80%**

OTHER INGREDIENTS: **39.20%**

TOTAL 100.00%

*2,4-dichlorophenoxyacetic acid equivalent 40.32% or 3.7 pounds per gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-866-944-8565 for emergency medical treatment information.	

For Additional Precautionary Statements, Complete First Aid, Directions for Use, Storage and Disposal and Other Use Information, See Inside This Label Booklet.

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300.

EPA REG. NO. 34704-1032

EPA EST. NO. 34704-MT-001

NET CONTENTS 2.5 GAL (9.46 L)

100115 V2D 01B16

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: Harmful if swallowed

Causes eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT/PPE:

Some materials that are chemical-resistant to this product are butyl rubber, natural rubber, neoprene or nitrile rubber.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes and socks,
- Chemical resistant gloves when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate,
- Chemical resistant apron when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

ENGINEERING CONTROLS STATEMENTS

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash-waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution must be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this product only as directed on label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

WHITEOUT® 2,4-D
EPA REG. NO. 34704-1032

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls,
- Chemical resistant gloves made of any waterproof material,
- Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or nurseries.

Do not enter or allow people (or pets) to enter the treated areas until sprays have dried.

PRODUCT INFORMATION

Performance of Whiteout® 2,4-D may be affected by local conditions, crop varieties, and application method. User should consult local extension service, agricultural experiment or university weed specialists, and state regulatory agencies for recommendations in your area.

Best results are obtained when product is applied to young succulent weeds that are actively growing. Application rates lower than specified will be satisfactory on susceptible annual weeds. For perennial weeds and conditions such as the very dry areas of the western states, where control is difficult, use the higher specified rates. When Whiteout 2,4-D is used for weed control in crops, the growth stage of the crop must be considered.

Some plants and weeds, especially woody varieties, are difficult to control and may require repeat applications. Apply 2.0 to 5.0 gallons of total spray by air or 5.0 to 25.0 gallons by ground equipment unless otherwise directed in this label or on Federally approved supplemental labeling for this product. In either case, use the same amount of Whiteout 2,4-D specified per acre. When an adjuvant is to be used with this product, Loveland Products, Inc. recommends using LI 700®. Aerial application must be used only when there is no danger of drift to susceptible crops. Many states have regulations concerning aerial application of 2,4-D formulations. Consult local regulatory authorities before making applications. Although Whiteout 2,4-D is a low volatile formulation, at temperatures above 90 °F vapors may damage susceptible plants nearby.

Product must not be used in greenhouses. Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination and all plant growth.

Spray equipment used to apply 2,4-D must not be used for any other purpose until thoroughly cleaned with a suitable chemical cleaner.

Spray Preparation: Add the specified amount of product to approximately one-half the volume of water to be used for spraying. Agitate well, and then add the remainder of the water. Continue agitation during application until spray tank is empty.

USE IN LIQUID NITROGEN FERTILIZER: Whiteout 2,4-D is specially formulated to be combined with liquid nitrogen fertilizer suitable for foliar application on corn, grass, pastures, or small grains in one operation. Use Whiteout 2,4-D according to directions on this label for those crops. Use liquid fertilizer at rates recommended by supplier or extension service specialist. Mix Whiteout 2,4-D and fertilizer according to the following instructions:

Add half the recommended amount of liquid fertilizer to the spray tank. Add Whiteout 2,4-D while agitating the tank. Add the remainder of the liquid fertilizer while continuing to agitate. Application should be made immediately, maintaining agitation until tank is empty. **DO NOT APPLY DURING COLD (NEAR FREEZING) WEATHER.** Spray mixture may not be stored.

SPRAY DRIFT

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. ground, aerial, airblast) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) within 250 feet downwind. If applying a medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial equipment and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

For aerial equipment, the boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made in a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

For ground boom application, do not apply with a nozzle height greater than 4 feet above the crop canopy.

APPLICATION INSTRUCTIONS

Spray volume: Apply with calibrated air or ground equipment using sufficient spray volume to provide adequate coverage of target weeds or as otherwise directed in specific use directions. For broadcast application, apply the specified rate of this product in a spray volume of 2.0 or more gallons per acre by air and 10.0 or more gallons per acre for ground equipment. Use low-pressure sprays to minimize drift. Where states have regulations, that specify minimum spray volumes, they must be observed. In general, spray volume should be increased as crop canopy, height and weed density increase in order to obtain adequate spray coverage.

Application Rates: Generally, lower rates in specified rate ranges will be satisfactory for more sensitive weeds species, when weeds are small, and when environmental conditions are favorable for rapid growth. Use higher rates in the specified rate range for less

WHITEOUT® 2,4-D
EPA REG. NO. 34704-1032

sensitive species and under less favorable growing conditions. Deep-rooted perennial weeds such as Canada thistle and field bindweed and many woody plants usually require repeated applications for effective control.

Spot Treatments

To prevent misapplication apply spot treatments with a calibrated boom or with hand sprayers using a fixed spray volume per 1000 square feet as indicated below.

Hand-Held Sprayers: Hand-held sprayers may be used for spot applications of Whiteout 2,4-D. Care must be taken to apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based on the application rate for an area of 1000 square feet. Mix the amount of Whiteout 2,4-D (fluid ounce or milliliter) corresponding to the desired broadcast rate in 1.0 to 3.0 gallons of spray. To calculate the amount of Whiteout 2,4-D required for larger areas, multiply the table value (fluid ounce or milliliter) by the thousands of square feet to be treated: An area of 1000 square feet is approximately 10.5 x 10.5 yards (strides) in size.

Rate Conversion Table for Spot Treatment:

Label Broadcast Rate (Pt/A)							
0.5	0.6	0.75	1.0	2.0	3.0	4.0	8.0
Equivalent Amount of Whiteout 2,4-D per 1000 Sq Ft							
0.2 fl oz (5.5 ml)	0.25 fl oz (7.3 ml)	0.33 fl oz (8.3 ml)	0.375 fl oz (11.0 ml)	0.75 fl oz (22.0 ml)	1.0 fl oz (33.0 ml)	1.5 fl oz (44.0 ml)	3.0 fl oz (88.0 ml)

Conversion factors: 1.0 pint =16.0 fluid ounces; 1.0 fluid ounce = 29.6 milliliters

Band Application: Whiteout 2,4-D may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated acre.

$$\text{Band rate per treated acre} = \frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast rate per acre}$$

$$\text{Band volume per treated acre} = \frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast volume per acre}$$

WHERE TO USE

Whiteout 2,4-D is used to control broad-leaved weeds in cereal crops, corn, sorghum, weeds and brush in rangeland, pastures, rights-of-way, ornamental turf, grass seed crops, fallow land, forest management and similar non-crop uses.

Annual or Biennial Weeds

Beggarticks	Lambsquarters, common	Rape, wild
Bittercress, smallflowered (2)	Lettuce, prickly (1) (2)	Rocket, yellow
Bitterweed	Lettuce, wild	Salsify, common
Broomweed, common (1)	Lupines	Salsify, western (1)
Burdock, common	Mallow, little	Shepherd's-purse
Buttercup, smallflowered (1) (2)	Mallow, Venice (1)	Sicklepod
Carpetweed	Marshelder	Smartweed (annual species) (1) (2)
Cinquefoil, common (2)	Morningglory, annual	Sneezeweed, bitter
Cinquefoil, rough (2)	Morningglory, ivy	Sowthistle, annual
Cocklebur, common	Morningglory, woolly	Sowthistle, spiny
Coffeeweed	Mousetail (2)	Spanishneedles
Copperleaf, Virginia	Mustards (except Blue mustard)	Sunflower
Croton, Texas	Parsnip, wild	Sweetclover
Croton, woolly	Pennycress (fanweed)	Tansymustard
Flixweed	Pepperweeds (<i>Lepidium</i> spp.) (1) (2)	Thistle, bull
Galinsoga	Pigweeds (<i>Amaranthus</i> spp.)	Thistle, musk
Geranium, Carolina (2)	Poorjoe	Thistle, Russian (tumbleweed)
Hemp, wild	Primrose, common	Velvetleaf
Horseweed (marestail) (2)	Purslane, common (2)	Vetches
Jewelweed	Pusley, Florida	
Jimsonweed	Radish, wild	
Knotweed (1)	Ragweed, common	
Kochia	Ragweed, giant	

WHITEOUT® 2,4-D
EPA REG. NO. 34704-1032

Perennial Weeds

Alfalfa (1)	Chicory	Ironweed, western (2)
Artichoke, Jerusalem (1)	Clover, red (1) (2)	Ivy, ground (1)
Aster, many-flower (1)	Coffeeweed	Nettles (including stinging) (1)
Austrian fieldcress	Cress, hoary (1)	Onion, wild (1)
Bindweed (hedge, field and European) (1) (2)	Dandelion	Pennywort
Blue lettuce	Docks (1)	Plantains
Blueweed, Texas	Dogbanes (1)	Ragwort, tansy
Broomweed	Eveningprimrose, cutleaf (2)	Sowthistle, perennial
Bullnettle (1) (2)	Garlic, wild	Thistle, Canada (1) (2)
Carrot, wild	Goldenrod	Vervains (1)
Catnip	Hawkweed, orange	Wormwood
	Healall	

- (1) Difficult-to-Control Weeds: These weeds are only partially controlled and may require repeat applications and/or use of the higher listed rate of this product even under ideal conditions of application.
(2) This product may not be used to control this weed species in the state of California.

CROPS
CEREAL GRAINS (Barley, Millet, Oats, Rye, Triticale, Wheat)
(Not under seeded with Legumes)

Crop/Application Timing	Whiteout 2,4-D (Pt/A)	Specific Use Directions
Barley, Millet, Rye, Triticale, Wheat Annual and biennial broadleaf weeds Perennial broadleaf weeds	0.5 to 2.0† 1.0 to 2.0†	Apply after crop is fully tillered, but before boot stage of growth (usually 4 to 8 inches tall) and weeds are small. Do not apply before tillering or from early boot through the milk stage of growth.
Oats (Spring Seeded) (Fall Seeded Southern)	0.5 0.75 to 1.25†	Apply after crop is fully tillered, but before boot stage of growth (usually 4 to 8 inches tall) and weeds are small. Do not apply before tillering or from early boot through the milk stage of growth. Do not apply during or immediately following cold weather.
Preharvest application (All cereals)	1.0	Apply using air or ground equipment to control weeds that could interfere with harvest, or to suppress perennial weeds. Apply when grain is in dough stage. Do not apply from early boot through the milk stage of growth.

† Use the lower rate in the rate range if small annual or biennial weeds are the major problem; use the higher rate if perennial weeds or biennial weeds are present that are considered to be hard-to-kill as determined by local experience. Higher rates increase the risk of crop injury and should be used only where weed control justifies such risk. Do not apply Whiteout 2,4-D at the crop seedling stage of growth. Consult state agricultural experiment station or extension service weed specialists for recommendations or suggestions to fit local conditions.

Cereal Grain Restrictions:

Postemergence:

- Make no more than 1 application per crop cycle.
- Do not apply more than 2.0 pints per acre per application.

Preharvest:

- Make no more than 1 application per crop cycle.
- Do not apply more than 1.0 pint per acre per application.
- Pre-Harvest Interval is 14 days.
- Do not graze lactating dairy cattle in treated areas for 14 days after application. Remove meat animals from freshly treated areas 7 days before slaughter.
- Do not harvest hay from treated grain fields.

Whiteout 2,4-D contains 0.46 pound acid equivalent of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 1.75 pounds of acid equivalent per acre per year.

WHITEOUT® 2,4-D
EPA REG. NO. 34704-1032

CORN (Field Corn, Popcorn and Sweet Corn)

Application Timing/Stage of Growth	Whiteout 2,4-D (Pt/A)	Specific Use Directions
Preplant (Burndown) Preemergence (Field corn, popcorn, and sweet corn)	1.0 to 2.0	General: For best results, growth conditions should be favorable for active weed growth. Use high rate in rate range for less susceptible weeds, cover crops such as alfalfa, weeds in advanced stages of development, or under less favorable growth conditions. Preplant: Apply 7 to 14 days before planting corn to control emerged broadleaf weed seedlings or existing cover crops. Preemergence: Apply any time after planting, but before corn emergence to control broadleaf weed seedlings or existing cover crops. Do not use on light sandy soils.
Postemergence (Field corn, popcorn, and sweet corn) Annual broadleaf weeds Crop up to 8 inches tall Crop 8 inches tall to tasseling (directed spray only) Perennial broadleaf weeds	0.5 to 1.0 1.0 1.0	Apply when weeds are small and corn is less than 8 inches tall (to top of canopy). If corn is more than 8 inches tall, use drop nozzles to keep spray off foliage. Treat perennial weeds when they are in bud to bloom stage. Do not harvest sweet corn before 45 days following application. Do not apply from tasseling to hard dough stage. Note: Corn treated with 2,4-D may become temporarily brittle. Wind or cultivation may cause stem breakage during the period of time that corn is brittle. Sweet corn: To minimize potential for crop injury, use only lowest rate in rate range.
Preharvest (Field corn and popcorn only)	Up to 3.0	Apply after corn is in hard dough (or denting) stage. Do not make preharvest applications to sweet corn.

Corn Restrictions:

- Do not forage or feed corn fodder to livestock for 7 days following application.

Preplant or Preemergence:

- Do not make more than 1 application per crop cycle.
- Do not apply more than 2.0 pints per acre per application.

Postemergence:

- Do not make more than 1 application per crop cycle.
- Do not apply more than 1.0 pint per acre per application.

Preharvest (field and popcorn ONLY):

- Do not make more than 1 application per crop cycle.
- Do not apply more than 3.0 pints per acre per application.
- Minimum spray interval between applications is 30 days.
- Pre-Harvest interval is 7 days.

Whiteout 2,4-D contains 0.46 pound acid equivalent of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 3.0 pounds of acid equivalent per acre per year.

SORGHUM (Grain Sorghum (Milo) and Forage Sorghum)

Application Timing/Stage of Growth	Whiteout 2,4-D (Pt/A)	Specific Use Directions
Postemergence† Crop 6 to 8 inches tall Crop 8 to 15 inches tall (directed spray only)	0.5 to 1.0† 0.75 to 1.0	Apply when sorghum is 6 to 15 inches tall. If sorghum is more than 8 inches tall (top of canopy), use drop nozzles to keep spray off of foliage. Do not use with oil or other adjuvants. Do not treat during boot, flowering or dough stage.

†Temporary crop injury can be expected under conditions of high soil moisture and high air temperatures. If it is necessary to apply Whiteout 2,4-D under these conditions, use no more than 0.6 pint per acre.

Sorghum Restrictions:

Postemergence:

- Do not make more than 1 application per crop cycle
- Do not apply more than 1.0 pint per acre per application.
- Do not permit meat or dairy animals to consume treated crop as fodder or forage for 30 days following application.

WHITEOUT® 2,4-D
EPA REG. NO. 34704-1032

Sorghum cont'd.:

- Preharvest Interval is 30 days.

Whiteout 2,4-D contains 0.46 pound acid equivalent of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 0.5 pound of acid equivalent derived from ester forms of 2,4-D per acre per year.

SOYBEANS — For Use in Crop Residue Management Systems
(Pre-plant Burndown Application Only)

Application Timing/Stage of Growth	Whiteout 2,4-D (Pt/A)	Number of Applications	Specific Use Directions
Preplant (Burndown) (OPTION 1)	0.75 to 1.0	Limited to 2 preplant applications/crop cycle	Apply not less than 7 days before planting soybeans. See Use Precautions and Restrictions below.
Preplant (Burndown) (OPTION 2)	1.0 to 2.0	Limited to 1 preplant application/crop cycle	Apply not less than 15 days before planting soybeans. See Use Precautions and Restrictions below.

Use Directions: Use Whiteout 2,4-D to control emerged broadleaf weeds or existing cover crops. For best results, apply when weeds are small and actively growing. Use the higher rate in the respective rate range for larger weeds and when perennials are present. Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may be added to spray mixtures to increase the herbicidal effectiveness on certain weeds. Read and follow all directions and precautions on this label and on the label of each product added to the spray mixture.

Restrictions and Limitations for Use in Soybeans

- Choose either Option 1 OR Option 2. These 2 options may not be used together in the same crop cycle, as the maximum application rate would be exceeded.
 - Do not apply this product prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield.
 - Do not use on low organic sandy soils (less than 1.0%).
 - Do not apply this product when weather conditions such as temperature, air inversions, or wind favor drift from treated areas to susceptible plants.
 - Do not mow or cultivate weeds prior to treating with this product as poor control may result.
 - Do not feed treated hay, forage, or fodder or graze treated soybeans to livestock. Do not feed or graze treated cover crops to livestock.
 - Do not replant fields treated with this product in the same growing season with crops other than those labeled for 2,4-D use.
- Whiteout 2,4-D contains 0.46 pounds acid equivalent of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 1.0 pound of acid equivalent per acre per year.

ORNAMENTAL TURF (Excluding Grasses Grown For Seed or Sod Farms)
(Includes lawns, golf courses, cemeteries and parks)

Use Requirements for Ornamental Turf Areas: When this product is applied to ornamental turf areas, follow PPE and reentry instructions in the “Non-agricultural Use Requirements” section of this label.

Treatment Site Application Timing	Whiteout 2,4-D (Pt/A)	Specific Use Directions
Ornamental Turf (Postemergence) Seedling grass (5-leaf stage or later)	0.75 to 1.0	Apply when weeds are small and actively growing. For best results, apply when soil moisture is adequate for active weed growth.
Well-established grasses	2.0 to 3.0	Deep-rooted perennial weeds such as Bindweed and Canada thistle may require repeat applications.
Biennial and perennial broadleaf weeds lished	3.0	Do not apply to newly seeded grasses until well established (5-leaf stage or later) and then use a maximum of 1 pt/A. Cool season grasses are tolerant of higher rates.

Ornamental Turf Restrictions:

- Do not apply more than 3.0 pints per acre per application.
- Do not make more than 2 broadcast applications per treatment site per year.

Whiteout 2,4-D contains 0.46 pound acid equivalent of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 3.0 pounds of acid equivalent per acre per year excluding spot treatments.

WHITEOUT® 2,4-D
EPA REG. NO. 34704-1032

GRASSES GROWN FOR SEED or SOD FARMS

Agricultural Use Requirements: When used in grass grown for seed or sod farms, follow PPE and reentry instructions in the "Agricultural Use Requirements" section of this label.

Treatment Site Application Timing	Whiteout 2,4-D (pt/acre)	Specific Use Directions
Grasses Grown for Seed (Postemergence Use) Seedling grass (5-leaf stage or later)	0.75 to 1.0	Apply when weeds are small and actively growing. For best results, apply when soil moisture is adequate for active weed growth.
Well-established grasses Sod Farms (Postemergence)	1.0 to 4.0 2.0 to 4.0	Do not apply to newly seeded grasses until well established (5-leaf stage or later) and then use a maximum of 1.0 pt/A. Cool season grasses are tolerant of higher rates. Do not apply to grass in the early boot through milk stage if seed production is desired. When grass is well established, higher rates of up to 4.0 pt/A may be applied for control of hard-to-kill annual or perennial weeds. Deep-rooted perennials such as Bindweed and Canada thistle may require repeat applications. Avoid mowing sod farms for 1 to 2 days before or after application. Delay irrigation until the day following application.

Grass Seed Crop Restrictions:

- Do not graze dairy cattle within 7 days of application. Do not apply this product within 30 days of cutting grass for hay. Remove meat animals from treated areas 7 days prior to slaughter.
- Do not apply more than 4.0 pints per acre per application.
- Do not make more than 2 applications per year.
- Minimum spray interval between applications is 21 days.
- Do not apply more than 8.0 pints per acre per year.
- If grass is to be cut for hay, Agricultural Use requirements for the Worker Protection Standard are applicable.

Whiteout 2,4-D contains 0.46 pound acid equivalent of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of acid equivalent per acre per year.

FALLOWLAND and CROP STUBBLE

Fallowland is considered to be idle cropland, postharvest to crops or between crops.

Type of weeds	Whiteout 2,4-D (Pt/A)	Specific Use Directions
Annual broadleaf weeds	1.0 to 2.0	Use a lower rate in the rate range when weeds are small (2 to 3 inches tall) and actively growing. Use a higher rate in the rate range when weeds are larger and under less favorable growth conditions.
Biennial broadleaf weeds	2.0 to 4.0	Apply when musk thistles or other biennial species are in the seedling to rosette stage and before development of flower stalks. The lower rate can be used in the spring during the rosette stage. Use the highest rate in the fall or after flower stalks have developed.
Perennial broadleaf weeds	2.0 to 4.0	Apply when perennial weeds are in bud to early bloom stage or while in good vegetative growth.
Wild garlic and onion in crop stubble	4.0	Apply to new regrowth of wild garlic or onion that occurs in the fall after harvest of other crops.

Fallow Land Restrictions:

- Plant only labeled crops within 29 days following last application.
- Do not apply more than 4.0 pints per acre per application.
- Do not make more than 2 applications per year.
- Minimum spray interval between applications is 30 days.

Whiteout 2,4-D contains 0.46 pound acid equivalent of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of acid equivalent per acre per year.

WHITEOUT® 2,4-D
EPA REG. NO. 34704-1032

RANGELAND, ESTABLISHED GRASS PASTURES
(Including Perennial Grasslands Not In Agricultural Production
Such As Conservation Reserve Program Acres)

Target Weeds or Woody Plants	Whiteout 2,4-D (Pt/A)	Specific Use Directions
Annual broadleaf weeds Biennial and perennial broadleaf weeds	2.0 2.0 to 4.0	For best results, apply when weeds are small and growing actively before the bud stage. Apply when Musk thistles or other biennial species are in the seedling to rosette stage and before flower stalks appear. Refer to the "Weeds Controlled" section for a listing of susceptible weed species and weeds that may be only partially controlled and require repeat applications and/or use of higher listed rates, even under ideal conditions of application.
Spot Treatment to control broadleaf weeds	See Instructions for "Spot Treatment" under "Application Instructions"	Note: To control broadleaf weeds in small areas with a hand sprayer, use an application rate equivalent to the listed broadcast rate and spray to thoroughly wet all foliage. See rate conversion table and instructions for "Spot Treatment" and use of hand-held sprayers under "Application Instructions."
Tree Injection Application	See instructions for tree injection application in "Forestry Uses" section.	
Wild garlic and wild onion	4.0	Make three applications (fall-spring-fall or spring-fall-spring) starting in late fall or early spring.
Broadleaf weed control in newly sprigged coastal bermudagrass	2.0 to 4.0	Applications may be made either preemergence or postemergence, Follow "Specific Use Directions" for annual, biennial and perennial broadleaf weed control, above.
Southern wild rose broadcast application	up to 4.0	Broadcast: Apply in spray volume of 5.0 or more gal/A by aircraft or 10.0 or more gal/A by ground equipment. Spot treatment: Apply when foliage is well developed. Thorough coverage is required. Use 8.0 pt of Whiteout 2,4-D plus 4.0 to 8.0 fl oz of an agricultural surfactant/100 gal of water. Two or more treatments may be required. Do not exceed 4.0 pt/A/application.
Spot treatment	8.0 pt/100 gal of spray	
CRP Acres		For program lands such as CRP, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

Pasture & Rangeland Restrictions:

- Do not apply more than 4.0 pints per acre per application.
- Do not apply more than 4.0 pints per acre per application for spot treatments.
- Do not make more than 2 applications per year.
- Minimum spray interval between applications is 30 days.
- Pre-Harvest Interval for cut forage for hay is 7 days.

• If grass is to be cut for hay, Agricultural Use requirements for the Worker Protection Standard are applicable.

Whiteout 2,4-D contains 0.46 pound acid equivalent of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of acid equivalent per acre per year.

WHITEOUT® 2,4-D
EPA REG. NO. 34704-1032

NON-CROPLAND AREAS

Such as fencerows, hedgerows, roadsides, drainage ditches, rights-of-way, utility power lines, railroads, airports, and other non-crop areas.

Treatment Site Method of Application	Whiteout 2,4-D (Pt/A)	Specific Use Directions
Annual broadleaf weeds	2.0 to 4.0	Apply when annual weeds are small and growing actively before the bud stage.
Biennial and perennial broadleaf weeds and susceptible woody plants	4.0 to 8.0	Biennial and perennial weeds should be rosette to bud stage, but not flowering at the time of application. For difficult to control perennial broadleaf weeds and woody species, tank mix up to 4.0 qt of Whiteout 2,4-D plus 1.0 to 8.0 pt of 3.0 lb/gal Triclopyr herbicide/A. For ground application: (High volume) apply a total spray volume of 100 to 400 gal/A; (low volume) apply a total spray volume of 10.0 to 100 gal/A. For helicopter: Apply a total spray volume of 5.0 to 30.0 gal/A.
Spot Treatment to control broadleaf weeds	See Instructions for "Spot Treatment" under "Application Instructions"	Note: To control broadleaf weeds in small areas with a hand sprayer, use an application rate equivalent to the listed broadcast rate and spray to thoroughly wet all foliage. See rate conversion table and instructions for "Spot Treatment" and use of hand-held sprayer under "Application Instructions."
Tree injection application	See instructions for tree injection application in "Forestry Uses" section.	
Southern wild rose broadcast application	up to 4.0	Broadcast: Apply in a spray volume of 5.0 or more gal/A by aircraft or 10.0 or more gal/A by ground equipment.
Spot treatment	8.0 pt/100 gal of spray	Apply when foliage is well developed. Thorough coverage is required. Use 8.0 pt of Whiteout 2,4-D plus 4.0 to 8.0 fl oz of an agricultural surfactant/100 gal of water. Two or more treatments may be required.

Weed Control Restrictions and Limitations for Use in Non-crop Areas:

Postemergence (annual & perennial weeds):

- Do not make more than 2 applications per year.
- Do not apply more than 4.0 pints per acre per application.
- Minimum spray interval between applications is 30 days.

Postemergence (woody plants):

- Do not make more than 1 application per year.
- Do not apply more than 8.0 pints per acre per application.
- Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Whiteout 2,4-D contains 0.46 pound acid equivalent of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of acid equivalent per acre per year.

FORESTRY USES

Forest site preparation, forest roadsides, brush control, established conifer release, including Christmas trees and reforestation areas.

Treatment Site Method of Application	Whiteout 2,4-D (Pt/A)	Specific Use Directions
Annual weeds	2.0 to 4.0 pt/A	Apply when weeds are small and growing actively before the bud stage. Apply when, biennial and perennial species are in the seedling to rosette stage and before flower stalks appear. For difficult to control perennial broadleaf weeds and woody species, use up to 4.0 qt of Whiteout 2,4-D and 1.0 to 4.0 qt of 3.0 lb/gal Triclopyr herbicide/A.
Biennial and perennial broadleaf weeds and susceptible woody plants	4.0 to 8.0 pt/A	For conifer release, make application in early spring before budbreak of conifers when weeds are small and actively growing.

WHITEOUT® 2,4-D
EPA REG. NO. 34704-1032

Forestry Uses cont'd.:

Treatment Site Method of Application	Whiteout 2,4-D (Pt/A)	Specific Use Directions
Spot Treatment to control broadleaf weeds	See instructions for "Spot Treatment" under "Application Instructions"	Note: To control broadleaf weeds in small areas with a hand sprayer, use an application rate equivalent to the listed broadcast rate and spray to thoroughly wet all foliage. See rate conversion table and instructions for "Spot Treatment" and use of hand-held sprayer under "Application Instructions."
Conifer Release; species such as white pine, ponderosa pine, jack pine, red pine, black spruce, white spruce, red spruce, and balsam fir	3.0 to 6.0 pt/A	To control competing hardwood species such as alder, aspen, birch, hazel, and willow, apply from mid to late summer when growth of conifer trees has hardened off and woody plants are still actively growing. Apply with ground or air equipment, using sufficient spray volume to ensure complete coverage. Because this treatment may cause occasional conifer injury, do not apply if such injury cannot be tolerated.
Directed Spray; Conifer Plantations including pine	8.0 pt/100 gal	Apply when brush or weeds are actively growing by directing the spray so as to avoid contact with conifer foliage and injurious amounts of spray. Apply in water carrier in a spray volume of 10.0 to 100 gal/A.
Surface of Cut Stumps (May also be used in rangeland, pastures, and noncropland) Frill and Girdle (May also be used in rangeland, pastures, and noncropland)	1.75 fl oz/gal of water	Apply as soon as possible after cutting trees. Thoroughly soak the entire stump with 2,4-D mixture including cut surface, bark and exposed roots. Cut frills (overlapping V-shaped notches cut downward through the bark in a continuous ring around the base of the tree) using an axe or other suitable tool. Saturate the freshly cut frills with the 2,4-D mixture.
Tree Injection Application (May also be used in rangeland, pastures, and noncropland)	(1.0 to 2.0 ml/ injection site)	To control and prevent resprouting of unwanted hardwood trees such as elm, hickory, oak, and sweetgum in forests and other non-crop areas, apply by injecting at a rate of 1.0 ml of undiluted Whiteout 2,4-D/inch of trunk diameter as measured at breast height (DBH), approximately 4-1/2 ft above the ground. Injection sites, however, must be as close to the root collar as possible and the injection bit must penetrate the inner bark. Applications may be made throughout the year, but for best results apply between May 15 and October 15. Maples should not be treated during the spring sap flow. For hard to control species such as ash, maple, and dogwood use 2.0 ml of undiluted Whiteout 2,4-D/ injection. Site injections are limited to one injection application per year. (max 2ml of 4 lbs ae formulation per injection site). Note: No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.

Forest Management Restrictions:

Broadcast Application:

- Do not make more than 1 application per year.
- Do not apply more than 8.0 pints per acre per application.

Injection Application:

- Do not make more than 1 application per year.
- Do not apply more than 2.0 milliliters per injection site.

Whiteout 2,4-D contains 0.46 pound acid equivalent of 2,4-D per pint. When tank mixing with products that contain 2,4-D, do not exceed a combined total of 4.0 pounds of acid equivalent per acre per year.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides or fungicides. Do not store near heat or open flame. Re-close all partially used containers by thoroughly tightening screw cap. Absorb any spill with suitable clay absorbent and dispose of as indicated under "Pesticide Disposal."

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org. If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For packages up to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 56 gallons: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY,

WHITEOUT® 2,4-D
EPA REG. NO. 34704-1032

EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Whiteout is a registered trademark of Loveland Products, Inc.

FORMULATED FOR
LOVELAND PRODUCTS, INC.
P.O. BOX 1286, GREELEY, COLORADO 80632-1286