

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Edith Emory Loveland Products, Inc. P.O. Box 1286 Greeley, CO 80632-1286 NOV - 3 2011

Subject:

First Choice Sanction Fungicide

EPA Registration No. 34704-1040

Your resubmission received on: October 26, 2011

Decision No. 446266

Dear Ms. Emory:

The label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable and the Agency acknowledges the new primary brand name of First Choice Sanction Fungicide. One copy of the label stamped "Accepted" is enclosed for your records. Please submit one copy of the final printed label before the product is released for shipment. If you have any questions regarding this correspondence, please contact Tracy Keigwin of my staff by phone at 703-305-6605 or via e-mail at keigwin.tracy@epa.gov.

Sincerely,

Mary L. Waller Mary L. Waller

Product Manager (21)


Fungicide Branch

Registration Division (7505P)

Enclosure: "Accepted" label



FIRST CHOICE® SANCTION® FUNGICIDE



For the control of powdery mildew on apples, grapes, stone fruits, strawberries, hops, lettuce, vegetables and other crops including ornamentals.

READ ENTIRE CONTAINER LABEL BEFORE USING THIS PRODUCT

ACTIVE INGREDIENTS:				
* Mono Potassium Phosphate				31.1%
** Sulfur			٠	31.1%
OTHER INGREDIENTS:				37.8%
	TOT	AL		100.0%

* Contains 4.1 lbs mono potassium phosphate per gallon

** Contains 4.1 lbs sulfur per gallon

CAUTION—PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

If Inhaled:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL: 1-866-944-8565

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

EPA REG. NO. 34704-1040 EPA EST. NO. 86868-CA-001 NET CONTENTS 2½ GALS NET WEIGHT 13.1 LBS/GAL

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye Irritation. Avoid contact with eyes, skin or clothing. Wear safety glasses or goggles when handling undiluted product. Once product is diluted, eye protection is no longer required. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE) Applicators and other handlers must wear:

Long-sleeved shirt and long pants

- Chemical resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

· Chemical-resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and was PPE separately from other laundry.

Engineering Controls Statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

· Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

· Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

 Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters. Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, or other undesirable results may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. PPE required for the early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material.
- · Shoes plus socks
- Protective eyewear
- · Chemical-resistant headgear for overhead exposure

PRODUCT INSTRUCTIONS

General: First Choice® Sanction® Fungicide is a foliar applied fungicide for control of powdery mildew. It is designed for foliar application and may be applied alone or in tank mixes containing pesticides. However, when use of an unfamiliar mix is made, a compatibility test is always recommended. First Choice Sanction Fungicide can be applied by ground or air. Foliage must be thoroughly covered with spray. Dense leaf canopies can prevent adequate spray coverage. To facilitate necessary coverage a super spreading agent such as Hi-Wett® or Widespread® Max is suggested. Powdery mildew control is best when applications of First Choice Sanction Fungicide are made within three days of a conidia spore release.

Mixing Directions: Fill the tank approximately 1/2 full with water before adding First Choice Sanction Fungicide. Agitate thoroughly while adding remaining water. Do not allow mixture to stand. Failure to maintain agitation will cause First Choice Sanction Fungicide to settle and may necessitate manual stirring to redisperse. When used in combination with pesticides, add pesticides to the tank last. Sulfur in any form is a corrosive material. To reduce the effect, equipment should be flushed daily.

Application Information: Apply the specified rate of First Choice Sanction Fungicide listed in the table when directed. When disease pressure is low, use low per acre rates early in the season. The per acre rate must be increased as disease pressure increases.

1. <u>Air Application Water Volume:</u> Orchards - apply in no less than 10 gallons of water per acre; All other applications - Apply in 5 to 20 gallons of water per acre.

2. <u>Ground Application (Concentrate) Water Volume:</u> Orchards - Apply in 20 to 100 gallons of water; All other applications - Apply in no less than 5 gallons of water per acre.

3. <u>Ground Application (Dilute) Water Volume: Orchards</u> - Apply in 100 to 800 gallons of water per acre; <u>All other applications</u> - Apply in 20 to 60 gallons of water per acre.

Compatibility: First Choice Sanction Fungicide is compatible with most pesticides and can be applied in existing spray programs. When using a chemical mixture that has not been used before, always try a small sample rate before application, or check compatibility by doing a jar test. Due to its acidic nature, adding First Choice Sanction Fungicide to the spray solution can lower the pH of the finished spray solution. Adhere to pesticide manufacturer's recommendations regarding appropriate pH range.

Use Precautions:

- Some plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised
 not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality. During
 periods of high temperature sulfur may burn foliage and fruit. Do not make applications of this product
 at excessively high temperatures (90 °F or higher).
- 2. Do not use First Choice Sanction Fungicide with oil or within 14 days of an oil application.
- 3. Do not include First Choice Sanction Fungicide with copper sprays.
- 4. Do not mix with any spray materials that warn against low pH (<5.5) applications.
- 5. First Choice Sanction Fungicide is a contact fungicide. The use of a spreading agent such as Widespread Max or Hi-Wett will aid in the distribution of product.
- 6. Do not use when plants are under severe stress conditions such as heat or water stress or pest pressure. Do not apply when temperatures exceed 90 °F.
- 7. Use caution when applying First Choice Sanction Fungicide in combinations with pesticides and adjuvants during the fruiting stage.
- 8. Use minimal effective rates of adjuvants during ripening. Organosilicone or high analysis non-ionic surfactants are not recommended during the ripening process. Hi-Wett at minimum recommended rates is recommended during the ripening period.
- 9. For maximum product effectiveness, apply early morning or after dusk.
- 10. Do not apply this product through any type of irrigation system.
- 11. Do not tank mix with any pesticide product which contains a prohibition on tank mixing.
- Follow the most restrictive precautions and Directions for Use of any tank mix partner.

In addition to the above 'Use Precautions', the following conditions must be followed:

- The operator of the property and the applicator must establish a buffer zone of enough distance to
 prevent drift onto non-target areas such as hospitals, clinics, residential areas, schools, and any other
 area designated by the county agricultural commissioner.
- 2. The operator of the property and the applicator must evaluate the method and equipment for each site to ensure proper and safe use. Evaluations shall include, but not be limited to the appropriateness of ground or air application.
- 3. This product must not be applied when the wind velocity exceeds 10 miles per hour. Applicators must be aware that in some areas, "dead calm" conditions are often associated with an inversion situation. In these areas, applying this product when there is a minimum air movement of 2 miles per hour will ensure that an inversion does not exist.
- 4. Before using this product, read and follow all applicable directions, restrictions and precautions on the EPA-registered label.

USE RATES (FOLIAR APPLICATION)¹

TREE	AND	VINE	CRO	DC
IIILL	MIND	ALIAL	UITU	1 3

CROP	RATE/ACRE	TIMING OF APPLICATIONS		
Apples	3 to 6 pts	Start spraying at tight cluster and continue spraying every 7 to 10 days until terminal shoots cease their vegetative growth. The rate of product per acre will vary depending upon the tree size (canopy development) and volume of water. Use on sulfur tolerant varieties only. Macintosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulfur under certain climatic conditions.		
Grapes	3 to 6 pts	Start spraying in the spring when shoots are 1 to 2 inches is (3 to 5 cm) in length and when disease pressure anticipated. Repeat every 10 to 14 days. For improved appearance on table grapes, use lower application rates.		
Stone Fruits (Apricots, Cherries, Nectarines, Peaches and Plums) 3 to 6 pts		Follow local recommendations for powdery mildew control timings, or apply when disease first appears and repeat every 7 to 14 days.		

CROP	RATE/ACRE	TIMING OF APPLICATIONS
Carrots	3 to 6 pts	Start spraying when disease is first observed. Repeat at 7 to 10 day intervals.
Cucurbits (including Cucumber, Summer and Winter Squash, Pumpkins and Watermelons	3 to 6 pts	Start spraying when plants begin to run or when disease pressure is anticipated. Repeat at 7 to 14 day intervals. Do not use on sulfur sensitive varieties.
Dry Beans (including Kidney, Pinto and Lima)	3 to 6 pts	Apply at first flower. Apply 2 additional times during the main filling stage of pod development, 7 to 10 days apart. Do not use on sulfur sensitive varieties.
Leafy Vegetables (including Lettuce)	3 to 6 pts	Start spraying when disease is first observed. Repeat at 10 to 14 day intervals.
Leafy Brassica Vegetables (including Broccoli, Brussels Sprouts, Cabbage and Cauliflower)	3 to 6 pts	Start spraying when disease is first observed. Repeat at 10 to 14 day intervals.
Tomatoes and Peppers	3 to 6 pts	Start spraying when disease is first observed. Repeat at 14 day intervals.

OTHER CROPS

CROP	RATE/ACRE	TIMING OF APPLICATIONS
Hops	3 to 6 pts	Start spraying when disease is first observed. Repeat at 7 to 10 day intervals.
Strawberries	3 to 6 pts	Apply 2 to 4 times during the harvest period on a 7 to 14 day schedule. Do not apply during hot weather. Sulfur may injure strawberries during certain climatic conditions. Do not use on sulfur sensitive varieties.

ORNAMENTALS

CROP	RATE/ACRE	TIMING OF APPLICATIONS
Ornamentals (including Woody and Herbaceous Ornamentals)	3 to 6 pts	Start spraying when conditions become favorable for disease development, or when disease is first observed and continue spraying on a 7 to 14 day schedule for the entire season.

- Use low rates per acre when disease pressure is low and high use rates when disease pressure increases.
- Rates suggested are for standard spray volumes of per acre. For higher volume sprays, rate should be adjusted.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in cool dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers and veterinary supplies. Avoid extreme variations in temperature. A reversible separation of ingredients may occur after prolonged storage. This separation has no effect on quality or effectiveness of product and agitation will resuspend mixture.

PESTICIDE DISPOSAL: Wastes resulting from this product may be disposed of on-site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for quidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org. If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For packages up to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows: Empty the

9

FIRST CHOICE® SANCTION® FUNGICIDE EPA REG. NO. 34704-1040

Storage & Disposal cont'd.:

remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. **For packages greater than 56 gallons:** To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVE-LAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS

PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Hi-Wett is a registered trademark of Elliott Chemicals, LTD. First Choice is a registered trademark of Crop Production Services, Inc. Sanction and Widespread are registered trademarks of Loveland Products, Inc.

FORMULATED FOR LOVELAND PRODUCTS, INC. P.O. BOX 1286, GREELEY, COLORADO 80632-1286