

34704-1054

1/2/2011

1/29



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

EPA Registration Number:

Date of Issuance:

34704-1054

FEB -2-2011

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Unconditional

Name of Pesticide Product:

LPI 6338-21

Name and Address of Registrant (include ZIP Code):

Loveland Products, Inc.  
P.O. Box 1286  
Greeley, Colorado 80632-1286

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5) provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data. If required, failure to submit acceptable data to fulfill these requirements may result in registration cancellation in accordance with FIFRA section 6(e).
- 2) Assure the type point is appropriate for the signal word and that the Keep out of Reach of Children statement meets the standards established by 40 CFR 156.60 (b) (1) and outlined in Chapter 3 of the Label Review Manual. For additional information on the appropriate type point, please consult "<http://www.epa.gov/oppfead1/labeling/lrm/chap-03.pdf>".
- 3) Under the PRECAUTIONARY STATEMENTS, replace the first two sentences with "Causes skin irritation. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid contact with eyes."
- 4) NOTE: The alternate CSF dated 7/7/10 is unacceptable because the total as stated does not add up to 100.00%. You may submit a revised alternate CSF through Front End Processing, if you so choose.

SEE NEXT PAGE FOR ADDITIONAL COMMENTS

Signature of Approving Official:

Kathryn V. Montague  
Product Manager 2B  
Herbicide Branch  
Registration Division (7505P)

Date:

5) The statements in the FIRST AID box must contain the following "If on skin" and "If in eyes" statements in this order:

"If on skin:

- Take off contaminated clothing
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

NOTE: The text that reads "Have the product container or label with you..." and "FOR A MEDICAL EMERGENCY..." must remain on the label. The "If swallowed" and "If inhaled" statements on the proposed label are not required. The optional "If swallowed" and "If inhaled" statements may remain on the label as additional safety language or they may be removed from the label.

6) Replace the last bullet on page 3 under the "Personal Protective Equipment" header that reads "When mixing and loading and when cleaning equipment, wear a chemical-resistant apron." with "Chemical-resistant apron when cleaning equipment, mixing, or loading."

7) Per the Metolachlor Reregistration Eligibility Document, replace the PPE bullet that reads "Coveralls" with "Coveralls over short-sleeve shirt and short pants" in the AGRICULTURAL USE REQUIREMENTS box.

8) Under the SOYBEAN APPLICATION DIRECTIONS on page 10, replace "cautions" with "precautions" in the last bullet under the "Restrictions" header for the statement to read "Observe all precautions and limitations on labeling of all products used in mixtures."

9) In the last sentence in the first bullet at the top of page 17, replace "cautions" with "precautions" for the sentence to read "Observe all precautions and limitations of all products used in tank mixtures."

10) Per the Metolachlor Reregistration Eligibility Document, assure the following endangered plants labeling is included in the Directions for Use under the "Restrictions" header on page 10.

"Do not apply under conditions which favor runoff or wind erosion of soil containing this product to non-target areas."

To prevent off-site movement due to run-off or wind erosion:

"Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation."

"Do not apply to impervious substrates such as pavement or highly compacted surfaces."

"Do not use tailwater from the first flood or furrow irrigation of treated fields to treat non-target crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation."

11) Per the Metolachlor Reregistration Eligibility Document, assure the following Rotational Crops Restriction is included in the Directions for Use under the "Restrictions" header on page 10.

"Do not rotate to food or feed crops other than those listed on this label."

12) Per the Metolachlor Reregistration Eligibility Document, assure the following skin sensitizer phrase is on the label under the "Hazards to Humans and Domestic Animals" header in the Precautionary Statements section:

"This product may cause skin sensitization reactions in some people."

13) The following changes must be made to the Container Disposal section of the Storage & Disposal box:

- a. In the first section (with the heading Nonrefillable Container), 7th line. The standard statement in the regulations is "...or contact [a pesticide container recycling organization] at [phone number] or [web site]. Therefore, you must add a phone number to the line that reads "To find the nearest site, contact your chemical dealer or manufacturer, or contact the Agricultural Container Recycling Council (ACRC) at *[you must include the appropriate phone number here and delete this italicized text and brackets]* or www.acrecycle.org."
- b. The term "packages" must be changed to "nonrefillable containers" in the three bold headers in this section to distinguish between nonrefillable and refillable (e.g., "For nonrefillable containers up to 5 gallons:").
- c. Replace the phrase "For refillable containers:" with "Refillable container." in the last section.
- d. In the last section, add a heading in the refillable container section that explains when the instructions are applicable. As appropriate, the heading could read "**For refillable containers from 56 to 330 gallons:** Refillable container. Refill this container with pesticide only. Do not reuse..." NOTE: The heading should identify the different size/type of containers as clearly as possible so the end user can easily find the right set of container handling instructions.
- e. Add the phrase "Offer for recycling or puncture and dispose of in a sanitary landfill or by incineration." as the last sentence to the refillable container section to explain what to do with the container.

14) NOTE: While no additional data is being requested at this time, any marketing claims made on the pesticide label must be substantiated by data maintained in your files. If data supporting marketing claims made on the product label is not available then those claims must be removed.

15) NOTE: Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

16) Submit one (1) copy of the revised final printed label before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

5/29

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

FEB - 2 2011  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

34704-1054



# LPI 6338-21

**Herbicide for preemergent control of certain grasses and broadleaf weeds in Soybeans**

ACTIVE INGREDIENTS:	By Wt.
Metolachlor* .....	43.72%
Metribuzin** .....	6.14%
Imazethapyr*** .....	1.38%
OTHER INGREDIENTS: .....	<u>48.76%</u>
TOTAL .....	100.00%

- \*contains 4.01 lbs of metolachlor per gallon.
- \*\*contains 0.56 lbs of metribuzin per gallon.
- \*\*\*contains 0.13 lbs of imazethapyr acid per gallon.

**KEEP OUT OF REACH OF CHILDREN  
WARNING-AVISO**


**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.**

**(If you do not understand the label, find someone to explain it to you in detail.)**

**EPA REG. NO. 34704-~~XXXX~~1054**

**EPA EST. NO. 34704-MS-1**

**NET CONTENTS 2.5 gal (9.46L)**



<b>FIRST AID</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911, or call an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</b>	
<b>FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565</b>	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING**

Causes skin irritation and moderate eye irritation. Do not get on skin or clothing and avoid contact with eyes. Wear protective eyewear. Wear coveralls, chemical-resistant gloves and chemical-resistant footwear. A chemical resistant apron plus chemical resistant gloves are required when mixing, loading, cleaning up spills or equipment or when otherwise exposed to this product's concentrate. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**Personal Protective Equipment**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category B on an EPA Chemical Resistance Category Selection Chart.

**Mixers, loaders, applicators, flaggers and other handlers must wear:**

- Protective eyewear,
- Coveralls worn over short-sleeved shirt and short pants,

- Chemical-resistant gloves made out of butyl rubber  $\geq$  14 mils or barrier laminate,
- Chemical-resistant footwear plus socks, and
- When mixing and loading and when cleaning equipment, wear a chemical-resistant apron.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

For terrestrial uses: do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

**Surface Water Advisory**

Metolachlor can contaminate surface water through ground spray drift. Under some conditions, metolachlor may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These conditions include: poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

**Ground Water Advisory and Mixing/Loading Instructions**

Metribuzin is a chemical which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Metribuzin has been found in ground water as a result of agricultural use. Users are advised not to apply metribuzin where the water table (ground water) is close to the surface, and where the soils are very permeable, i.e., well-drained soils such as loamy sands. Your local

agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Imazethapyr has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Metolachlor has the potential to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

DO NOT apply this product through any type of irrigation system.

Product must be used in a manner which will prevent back-siphoning in wells, spills or improper disposal of excess pesticide spray mixture.

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes or reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times.

The above specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.



9/29

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Observe all cautions and limitations on this label on the labels of products used in combination with this product.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protections of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protections Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls,
- Chemical-resistant gloves, such as butyl rubber  $\geq$  14 mils or barrier laminate,
- Chemical-resistant footwear plus socks.

#### PRODUCT INFORMATION

LPI 6338-21 kills weeds by root and/or foliage uptake and rapid translocation to the growing points. Adequate soil moisture is important for optimum activity of this product. When adequate soil moisture is present, this product will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

Occasionally, internode shortening and/or temporary yellowing of crop plants may occur following applications of this product. These effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.

When organophosphate (such as Lorsban®) or carbamate insecticides are tank mixed with

this product, temporary injury may result to the treated crops.

Use of this product in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product. Therefore, rotational crop injury is always possible.

Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall), this product may cause injury to subsequent planted crops. Vegetable crops (particularly sugar beets) are sensitive to residues of this product in the soil.

Naturally occurring biotypes\* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with either the ALS/AHAS enzyme inhibiting mode of action. Other herbicides with the ALS/AHAS enzyme inhibiting mode of action include the sulfonylureas (e.g., Accent®, Basis®, Classic®, Harmony® GT, Spirit®, Permit®, etc.), the sulfonamides (e.g., FirstRate®, etc.) and the pyrimidyl benzoates (e.g. Staple®, etc.).

\* A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants.

**AERIAL DRIFT REDUCTION ADVISORY INFORMATION**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
3. Where states have more stringent regulations, they must be observed.
4. The applicator must be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

**Information On Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity and Temperature Inversions).

**Controlling Droplet Size**

11/29

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

**Boom Length**

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

**Application Height**

**Do not make** applications at a height greater than 10 feet above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**Swath Adjustment**

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up-and-downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

**Wind**

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind directions and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions**

Because drift potential is high, do not apply during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in

a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Sensitive Areas**

This product may only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

**APPLICATION WITH HERBICIDE SPRAY EQUIPMENT**

Use a standard low-pressure (20 to 40 psi) herbicide boom sprayer equipped with suitable nozzles and screens no finer than 50-mesh in nozzle and in-line strainers. Agitate thoroughly before and during application with bypass agitation. Low pressure and high volume hand wand equipment is prohibited.

**Ground Application**

Apply the proper rate of this product in a minimum of 10 to 40 gallons of spray mixture per acre broadcast.

**Aerial Application**

Where permitted, apply specified rate in a minimum of 5 gallons of spray mixture per acre. Do not apply aerially when wind speed is greater than 10 mph.

**For All Applications**

Sprayer must be accurately calibrated before applying this product. Check sprayer during application to be sure it is working properly and delivering a uniform spray pattern. As the volume of spray mixture decreases per acre, the importance of accurate calibration and uniform application increases.

Avoid overapplication, misapplication, and boom and spray swath overlapping that will increase spray dosage. (Crop injury may occur as a result.) Avoid spray skips and gaps which allow weeds to grow in untreated soil. Do not apply when weather conditions favor spray drift and/or when sensitive or cool season crops (such as cole crops, onions, peas, or strawberries) are present in adjacent fields or in areas where wheat is growing in coarse textured soils.

**Sprayer Cleanup**

Spray equipment must be thoroughly cleaned to remove remaining traces of herbicide that might injure other crops to be sprayed. Drain any remaining spray solution of this product from the spray tank and dispose of according to label disposal instructions. Rinse

the spray tank and refill with water, adding a heavy-duty detergent at the rate of one cup per 20 gallons of water. Recycle this mixture through the equipment for 5 minutes and spray out. Repeat this procedure twice. Fill the spray tank with clean water, recycle for 5 minutes, and spray out. Clean pump and nozzle screens thoroughly. Wash away any spray mixture from the outside of spray tank, nozzles or spray rig. All rinse water must be disposed of in compliance with local, state, and Federal guidelines.

## MIXING INSTRUCTIONS

### Incorporation and Combination Uses

When this product is to be used in combination with another herbicide, follow the most restrictive directions on all product labels for combinations, rates, crops, incorporation, and special precautions.

When using this product, make sure the sprayer is completely clean, free of rust or corrosion which occurs from winter storage. Examine strainers and screens to be sure the sprayer is clean from previously used pesticides.

Any tank mix containing this product must be kept agitated and sprayed out immediately. Do not allow tank mixes to stand for prolonged periods of time.

The proper mixing procedure for this product alone or in tank-mix combinations with other herbicides is:

1. Fill the spray tank  $\frac{1}{4}$  to  $\frac{1}{3}$  full with clean water.
2. Add specified rate of this product while recirculating and with agitator running.
3. Mix thoroughly and add clean water to fill spray tank to desired level.
4. Add the other herbicide to tank last and agitate thoroughly.
5. Continue agitation during application and until sprayer tank is empty.

### Application of LPI 6338-21 In Fluid Fertilizers

This product may be applied in fluid fertilizer solutions by following the appropriate mixing procedures and compatibility check. When using tank mix combinations, be sure all components are compatible.

### Tank Mixing Guidelines for Fluid Fertilizer Mixtures

1. Add the required amount of water and compatibility agent (if required) to the tank. Start agitation system while adding this product and follow by adding the fluid fertilizer and agitate.
2. If a second herbicide is also to be used, follow as above in Step 1, but use twice the amount of water. Start agitation, add LPI 6338-21. Follow by adding the second herbicide, then continue filling the tank with fluid fertilizer.
3. Maintain continuous agitation to assure uniform spray mixture until the tank is emptied.

Make compatibility checks of this product plus fluid fertilizers and tank-mix combinations plus fluid fertilizers which include this product for each batch because of the variability of fluid fertilizers.

**THE FOLLOWING COMPATIBILITY CHECK SHOULD ONLY BE USED WHEN MIXING WITH FLUID FERTILIZERS.**

1. Pre-mix 8 teaspoons of water with 2 teaspoons of this product (4:1 ratio) in a quart jar by adding the water first and following with this product. Mix thoroughly. If a second herbicide is to be used, double the amount of water (8:1 ratio), mix in this product, and follow with the second herbicide.
2. Then pour 1 pint of fluid fertilizer into the quart jar and shake well.
3. Allow to stand for 5 minutes.

### **Interpretation of Results**

If the solution in the jar appears to be uniform, without signs of agglomeration, or without a separation of an oily film on top of the fertilizer, the mixture may be used. If not, repeat the compatibility check using twice the amount of water or add a compatibility agent to the water. If separation occurs, but the mixture can be resuspended by shaking, then application is possible with good agitation in the spray tank.

### **SOYBEAN APPLICATION DIRECTIONS**

**(Except California, and North Dakota and Minnesota North of Highway #210)**

This product may be applied preplant incorporated, preplant surface or preemergent surface, or as a sequential preemergent application. This product may also be used as an overlay application following a preplant incorporated application of a grass herbicide registered for this same use and in tank mix combinations for burndown weed control. All applications may be applied with ground equipment, and some may be applied with aerial spray equipment.

### **Restrictions**

- Do not apply more than 4.0 pints of LPI 6338-21 per acre per use season.
- Do not harvest within 85 days of the last application of LPI 6338-21.
- Do not graze or feed treated soybean forage, hay or straw to livestock.
- Do not rotate any crop not listed on this label for 40 months following application.
- Do not apply LPI 6338-21 preemergence to Soybeans in California, North Dakota and Minnesota north of highway #210.
- Do not incorporate into soil or apply more than once per season except where permitted as part of a sequential application.
- Do not allow sprays to drift onto adjacent desirable plants.
- Do not apply using low-pressure and high-volume hand-wand equipment.
- Observe all cautions and limitations on labeling of all products used in mixtures.

### **Soil Texture and Rate Ranges**

As used on this label, "Coarse soils" are loamy sand or sandy loam soils. "Medium soils" are loam, silt loam, silt, sandy clay, or sandy clay loam. "Fine soils" are silty clay, silty clay loam, clay, or clay loam. Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

Where a rate range is shown, use a lower rate on soils that are coarse-textured or low in organic matter. Use a higher rate on soils that are relatively fine-textured or high in organic matter.

### **Precautions**

LPI 6338-21 EPA REG. NO. 34704-XXXX

Injury to soybeans may occur when this product is used under the following conditions:

1. When soils have a calcareous surface area or a pH of 7.5 or higher.
2. When applied in conjunction with soil-applied organic phosphate pesticides.
3. With over-application or boom overlapping, which may result in stand loss and soil residues.
4. With uneven application or improper incorporation, which can decrease the level of weed control and/or increase the level of injury.
5. When applied to any soil with less than 0.5% organic matter.
6. When soil incorporation is deeper than recommended.
7. When sprayers are not calibrated accurately.
8. When heavy rains occur soon after application, especially in poorly drained areas where water may stand for several days.
9. When soybeans are planted less than 1-½ inches deep, particularly in preemergence application.
10. Where high soil levels of atrazine are present.
11. When using poor quality soybean seed.

Certain soybean varieties are sensitive to Metribuzin. Prior to use of this product, consult your soybean seed supplier for more information on the tolerance of soybean varieties to LPI 6338-21.

**Activation**

A minimum amount of soil moisture is required to activate this product. In areas of low rainfall, preemergence applications to dry soil should be followed with light irrigation of ¼ acre-inch of water. Do not apply heavy irrigation immediately after application. As with many surface-applied herbicides, weed control and crop tolerance may vary with rainfall and/or soil texture.

**Replanting**

If replanting is necessary in fields treated with this product as directed on this label, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. Do not apply more than once per season except where permitted as part of a sequential application as injury to soybeans may occur. Maximum application rate is 4.0 pints of LPI 6338-21 per acre per use season. Do not exceed this amount in any use pattern: single application, replant or sequential application.

**TABLE 1: ANNUAL BROADLEAF WEEDS CONTROLLED BY LPI 6338-21**

C = Control S = Suppression or Erratic Control P = Poor or No Control U= Unknown	
Weed Controlled	Level of Control
Bristly Starbur ( <i>Acanthospermum hispidum</i> )	C
Buffalobur ( <i>Solanum rostratum</i> )	C
Carpetweed ( <i>Mollugo verticillata</i> )	C
Cocklebur ( <i>Xanthium pensylvanicum</i> )	S
Common Chickweed ( <i>Stellaria media</i> )	C
Copperleaf, Hophornbeam ( <i>Acalypha ostryifolia</i> )	C
Field Pennycress ( <i>Thlaspi arvense</i> )	C
Florida Beggarweed ( <i>Desmodium tortuosum</i> )	C

LPI 6338-21 EPA REG. NO. 34704-XXXX

Florida Pusley ( <i>Richardia scabra</i> )	C
Galinsoga ( <i>Galinsoga</i> spp.)	C
Henbit ( <i>Lamium amplexicaule</i> )	C
Horseweed (Marestail) ( <i>Conyza canadensis</i> )	U
Jimsonweed ( <i>Datura stramonium</i> )	C
Knotweed ( <i>Polygonum</i> spp.)	C
Kochia ( <i>Kochia scoparia</i> )	C
Lambsquarters ( <i>Chenopodium</i> spp.)	C
Marshelder ( <i>Iva Annua</i> )	C
Morningglory	
Entireleaf ( <i>Ipomoea hederacea</i> var. <i>integriuscula</i> )	S
Ivyleaf ( <i>Ipomoea hederacea</i> )	S
Pitted ( <i>Ipomoea lacunosa</i> )	S
Smallflower ( <i>Jacquemontia tamnifolia</i> )	C
Tall ( <i>Ipomoea purpurea</i> )	S
Mustard spp.	C
Nightshade	
Black ( <i>Solanum nigrum</i> )	C
Eastern Black ( <i>Solanum ptycanthum</i> )	C
Hairy ( <i>Solanum villosum</i> )	C
Pigweed	
Redroot ( <i>Amaranthus retroflexus</i> )	C
Smooth ( <i>Amaranthus hybridus</i> )	C
Spiny ( <i>Amaranthus spinosus</i> )	C
Poinsettia, Wild ( <i>Euphorbia cyathophora</i> )	C
Prickly Lettuce ( <i>Lactuca scariola</i> )	C
Prickly Sida/Teaweed ( <i>Sida spinosa</i> )	C
Puncturevine ( <i>Tribulus terrestris</i> )	C
Purslane ( <i>Portulaca oleracea</i> )	C
Ragweed	
Common ( <i>Ambrosia artemisiifolia</i> )	C
Giant ( <i>Ambrosia trifida</i> )	S
Redweed	C
Russian Thistle ( <i>Salsola kali</i> )	C
Sage, Barnyard	S
Sesbania ( <i>Sesbania</i> spp.)	C
Shepherd's Purse ( <i>Capsella bursa-pastoris</i> )	C
Sicklepod ( <i>Cassia obtusifolia</i> ) <sup>1</sup>	C
Smartweeds ( <i>Polygonum</i> spp.)	
Ladysthumb ( <i>Polygonum persicaria</i> )	C
Pennsylvania ( <i>Polygonum pennsylvanicum</i> )	C
Spurge	
Prostrate ( <i>Euphorbia humistrata</i> )	C
Spotted ( <i>Euphorbia maculata</i> )	C
Spurred Anoda ( <i>Anoda cristata</i> )	C



LPI 6338-21 EPA REG. NO. 34704-XXXX

Sunflower ( <i>Helianthus</i> spp.)	C
Velvetleaf ( <i>Abutilon theophrasti</i> )	C
Venice Mallow ( <i>Hibiscus trionum</i> )	C
Virginia Pepperweed ( <i>Lepidium virginicum</i> )	C
Waterhemp ( <i>Amaranthus rudis</i> )	C
Wild Mustards ( <i>Brassica</i> spp.)	C

<sup>1</sup>For maximum control of sicklepod, use a preemergence application.

**TABLE 2: ANNUAL GRASSES AND SEDGES CONTROLLED BY LPI 6338-21**

C = Control S = Suppression or Erratic Control P = Poor or No Control	
Weed Controlled	Level of Control
Barnyardgrass ( <i>Echinochloa crus-galli</i> )	C
Bluegrass ( <i>Poa annua</i> )	C
Broadleaf Signalgrass ( <i>Brachiaria platyphylla</i> )	C
Browntop Millet ( <i>Panicum ramosa</i> )	C
Crabgrass ( <i>Digitaria</i> spp.)	C
Crowfootgrass ( <i>Dactyloctenium aegyptium</i> )	C
Cupgrass ( <i>Eriochloa gracilis</i> )	C
Foxtails ( <i>Setaria</i> spp.)	C
Goosegrass ( <i>Eleusine indica</i> )	C
Johnsongrass, Seedling ( <i>Sorghum halepense</i> )	C
Junglerice ( <i>Echinochloa colona</i> )	C
Millet, Wild-proso ( <i>Panicum miliaceum</i> )	S
Nutsedge	
Yellow ( <i>Cyperus esculentus</i> )	S
Purple ( <i>Cyperus rotundus</i> )	S
Panicum, Fall ( <i>Panicum dichotomiflorum</i> )	C
Panicum, Texas ( <i>Panicum, texanum</i> )	S
Red Rice ( <i>Oryza sativa</i> )	C
Sandbur ( <i>Cenchrus</i> spp.)	S
Shattercane ( <i>Sorghum bicolor</i> )	S
Sorghum, Volunteer ( <i>Sorghum</i> spp.)	S
Sprangletop, ( <i>Leptochloa</i> spp.)	P
Stinkgrass ( <i>Eragrostis</i> spp.)	P
Wheat, Volunteer ( <i>Triticum</i> spp.)	S
Witchgrass ( <i>Panicum capillare</i> )	C

**LPI 6338-21 USE RATES FOR COVENTIONAL TILLAGE SYSTEMS**

**LPI 6338-21 used alone in Preplant Incorporated Application**

Incorporate LPI 6338-21 uniformly into the top 2 inches of soil within 14 days before planting using a disk, field cultivator, rolling cultivator or similar equipment. Use incorporated application if furrow irrigation is used or when a period of dry weather after application is expected.

**LPI 6338-21 used alone in Preemergence Application**

When used alone, LPI 6338-21 can be applied broadcast by ground or by aerially. This application may be made during planting or as a separate operation after planting but must be made before crop emergence. If dry weather follows preemergence application, cultivate uniformly with shallow tilling equipment that will not damage soybeans.

**Restrictions**

Do not apply to sand soils, or to sandy loam or loamy sand soils containing less than 2% organic matter.

**TABLE 3: LPI 6338-21 RATES WHEN USED ALONE IN PREPLANT OR PREEMERGENCE APPLICATION**

Soil Texture		Organic Matter		
		0.5 to 2%	2 to 3%	Over 3% <sup>3</sup>
Pints of LPI 6338-21 Per Acre				
Coarse Soils <sup>1</sup>	Sandy Loam	1.6 to 2.0 <sup>4</sup>	1.6 to 2.0 <sup>4</sup>	2.0 to 2.4
	Loamy Sand	n/a <sup>1</sup>	1.6 to 2.0 <sup>4</sup>	2.0 to 2.4
Medium Soils (Loam, silt loam, silt, sandy clay, sandy clay loam)		2.4 to 2.7		2.7 to 3.1
Fine Soils (Silty clay, silty clay loam <sup>2</sup> , clay, clay loam)		3.1 to 3.5		3.5 to 3.9

<sup>1</sup> Do not use on sand soils. On coarse-textured soils, do not use on loamy sand with less than 2% organic matter.

<sup>2</sup> Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

<sup>3</sup>For preplant incorporated application, use the lower rate.

<sup>4</sup>For AL, AR, FL, GA, LA, MS, MO, NC, OK, SC, TN, TX, VA, see section below **In Coarse (Light) Soils.**

On soils with pH above 7.0, soybean injury caused by the metribuzin in this product may occur at rates higher than 3.3 pts/A. To avoid injury, do not use this product at rates greater than 3.3 pts/A on soils above pH 7.0.

**In Coarse (Light) Soils**

(Only in AL, AR, FL, GA, LA, MS, MO, NC, OK, SC, TN, TX, VA)

This product is may be used at the rates specified in Table 4 as a preplant incorporated or preemergence application in coarse-textured, low organic matter soils in the states listed above. Refer to Table 4 and to the appropriate sections of this label for specific directions on use and restrictions.

29/29

**TABLE 4: LPI 6338-21 RATES WHEN USED ALONE IN PREPLANT OR PREEMERGENCE APPLICATIONS ON COARSE SOILS (Only in AL, AR, FL, GA, LA, MS, MO, NC, OK, SC, TN, TX, VA)**

		Organic Matter	
		0.5 to 1%	1% or above
Soil Texture		Pints of LPI 6338-21 Per Acre <sup>2</sup>	
Coarse Soils	Sand	n/a <sup>1</sup>	1.6 to 2.7
	Sandy loam, loamy sand	1.6 to 2.7	1.6 to 2.7

<sup>1</sup>Do not use on sand with less than 1% organic matter.

<sup>2</sup>Use the higher rate under heavy weed pressure and/or soils higher in organic matter.

On soils with pH above 7.0, soybean injury caused by the metribuzin in this product may occur at rates higher than 3.3 pts/A. To avoid injury, do not use this product at rates greater than 3.3 pts/A on soils above pH 7.0.

**HERBICIDES THAT MAY BE APPLIED POSTEMERGENCE FOLLOWING LPI 6338-21**

If required, application of this product alone or in tank mixture may be followed by an application of a postemergence herbicide to provide additional control of certain weeds. The following postemergence herbicides may be applied:

Aim <sup>®</sup>	glyphosate herbicides <sup>1</sup> (such as Makaze or Mad Dog)
Arrow <sup>™</sup>	Harmony <sup>®</sup> GT XP
Assure <sup>®</sup> II	Intensity <sup>®</sup>
Basagran <sup>®</sup>	Poast <sup>®</sup>
Classic <sup>®</sup>	Poast Plus <sup>®</sup>
Cobra <sup>®</sup>	Reflex <sup>®</sup>
FirstRate <sup>®</sup>	Resource <sup>®</sup>
Flexstar <sup>®</sup>	Rezult <sup>®</sup> A&B
Fusilade <sup>®</sup> DX	Storm <sup>®</sup>
Frontrow <sup>®</sup>	Synchrony <sup>®</sup> XP <sup>2</sup>
Fusion <sup>®</sup>	Ultra Blazer <sup>®</sup>

<sup>1</sup>Use on Roundup-Ready<sup>®</sup> or glyphosate tolerant soybean varieties only.

<sup>2</sup>Use on STSTM soybean varieties only.

Refer to the **Directions for Use** on this label and the individual product labels for use directions, use rates, and special precautions and/or restrictions.

**BURNDOWN WEED CONTROL**

This product can be used as part of a burndown herbicide program for control of existing vegetation prior to soybean emergence in conservation tillage (reduced-tillage/no-till)

systems. This product may be tank mixed with a 2,4-D low volatile ester (LVE) (such as Whiteout™) and/or glyphosate herbicides (such as Mad Dog® and Makaze® brands) for control of emerged weeds prior to crop emergence. Burndown tank mixes with LPI 6338-21 can be applied before planting or prior to crop emergence.

**DO NOT** tank mix LPI 6338-21 with clomazone containing herbicides (Command®).

**Application**

This product may be applied up to 30 days before planting or preemergence. Apply only by ground equipment when this product is used for burndown of existing vegetation in conservation tillage systems. Use the high end of the rate range for applications of this product made 14 to 30 days before planting. Refer to Tables 3 and 4 for rates of LPI 6338-21 alone and to Table 5 for rates of tank mix partners.

**TABLE 5: RATES OF TANK MIX PARTNERS TO BE USED IN COMBINATION WITH LPI 6338-21 FOR BURNDOWN APPLICATIONS**

Product	Rate of Tank Mix Partner	Directions and Remarks
2,4-D LVE (Whiteout 2,4-D)	0.25 to 1 lb. a.e. <sup>1</sup> /A.	Apply at least 7 days preplant when using Whiteout at 0.25 to 0.5 lb. a.e. <sup>1</sup> /A and at least 30 days preplant with rates greater than 0.5 lb. a.e. <sup>1</sup> /A. Include crop oil concentrate (COC) at the rate of 1 gal./100 gal. of spray solution (1% v/v).
Glyphosate (Mad Dog, or Makaze brands)	Refer to product label for use rates.	Must be applied prior to crop emergence. Use the higher rates within the specified range as weeds approach the maximum weed heights listed in Table 6. Apply in 10 to 20 gal. of water per acre. Refer to the Mad Dog or Makaze label for spray adjuvant instructions. Any glyphosate formulation registered and labeled for use in soybeans may be tank mixed with this product.
Glyphosate (Mad Dog or, Makaze brands) + 2,4-D LVE (Whiteout 2,4-D)	Refer to the product label for use rates + 0.25 lb. a.e. <sup>1</sup> /A.	Follow the Directions and Remarks section above for Whiteout 2,4-D and Mad Dog/Makaze, paying special attention to planting restrictions with Whiteout. Refer to the Mad Dog or Makaze label for spray adjuvant instructions. Do not use crop oil concentrate (COC).

<sup>1</sup>a.e. = 2,4-D acid equivalent

**Precautions**

Do not apply these treatments after crop emergence. Observe all precautions and limitations on the labeling of all products used in tank mixtures.

- Apply only 2,4-D LVE formulations (such as Whiteout 2,4-D ) that are registered

and recommended for preplant or burndown use.

- Do not apply tank mixtures containing 2,4-D LVE (Whiteout 2,4-D ) if wind is blowing toward desired susceptible plants (i.e., cotton, tobacco, tomato, etc.) or when wind speeds exceed 6 miles per hour. Observe all cautions and limitations of all products used in tank mixtures.

Follow the most restrictive preharvest interval of all products used in a tank mixture.

**Weeds Controlled**

LPI 6338-21 in tank mixtures with the herbicides listed in Table 6 will provide burndown control of the weeds listed below.

**TABLE 6: WEEDS CONTROLLED WITH TANK MIXES OF LPI 6338-21 IN BURNDOWN APPLICATION**

Weeds Controlled	Whiteout 2,4-D	Mad Dog/Makaze	Mad Dog/Makaze + Whiteout 2,4-D
<b>Annual Grasses</b>	<b>Maximum Burndown Height (Inches)</b>		
Barley	Does not improve control of these species	8	
Barnyardgrass		6	
Crabgrass spp.		6	
Foxtail spp.		8	
Johnsongrass, Seedling		8	
Panicum, Fall		6	
Sandbur, Field		8	
Wheat, Volunteer		6	
Witchgrass		6	
<b>Broadleaves</b>		<b>Maximum Burndown Height (Inches)</b>	
Buffalobur	-	6	6
Chickweed, Common	6	6	6
Cocklebur, Common	6	6	8
Dandelion, Common	6 dia. <sup>1</sup>	2 dia. <sup>2</sup>	6 dia. <sup>1</sup>
Henbit	4	4	4
Horseweed (Marestail)	6 <sup>1</sup>	4 <sup>2</sup>	6
Jimsonweed	6	6	6
Kochia	4 <sup>1</sup>	4	4
Ladysthumb	6	6	8
Lambsquarters, Common	6	6	8
Lettuce, Prickly	6	4	6
Mallow, Venice	6	6	6
Morningglory spp.	6	2	4
Mustard spp.	6	6	8
Pennycress, Field	6	6	6
Pigweed spp. (annual)	6	6	8
Ragweed, Common	6	6 <sup>2</sup>	8
Ragweed, Giant	6 <sup>1</sup>	4 <sup>2</sup>	6
Shepherd's Purse	6	6	6
Sida, Prickly	6	4	4
Smartweed, Pennsylvania	6	6	8
Sunflower, Common	6	6	6

LPI 6338-21 EPA REG. NO. 34704-XXXX

Thistle, Russian	4 <sup>1</sup>	2-4 <sup>2</sup>	4
Velvetleaf	6	6	8
Waterhemp spp.	6	6	8
<sup>1</sup> Use Whiteout 2,4-D at 0.5 lb. ai/A.			
<sup>2</sup> Use a minimum of 0.75 lb. ai/A of Mad Dog or Makaze.			

**LPI 6338-21 USE RATES FOR REDUCED- AND NO-TILL SYSTEMS**

**Preplant Surface Application**

LPI 6338-21 may also be used in reduced-till and no-till systems. Applications may be made up to 30 days before planting or after planting, but before soybean emergence. Residual herbicides such as Canopy, FirstRate, Command, Python®, and Stealth® may be tank mixed for additional weed control. If weeds are present at time of application, burndown herbicides may be added to the tank mixes (see **Burndown Weed Control** section). Refer to the tank mix partner product labels for specific rates and use directions.

**TABLE 7: LPI 6338-21 RATES FOR REDUCED AND NO-TILL SYSTEMS**

Soil Texture <sup>1</sup>	LPI 6338-21 (Pt./A <sup>1</sup> )
COARSE <sup>2</sup> (Loamy sand, sandy loam)	1.6 to 2.7
MEDIUM (Loam, silt loam, silt, sandy clay, sandy clay loam)	2.7 to 4.0
FINE (Silty clay, silty clay loam <sup>3</sup> , clay, clay loam)	4.0
<sup>1</sup> Use low rate in specified range for low residue level or soils with less than 3% organic matter. Use the higher rate in specified range for high residue level or soils with greater than 3% organic matter.	
<sup>2</sup> Do not use on sand soils. On coarse-textured soils, do not use on loamy sand soils with less than 2% organic matter.	
<sup>3</sup> Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S. When using LPI 6338-21, treat this soil as fine-textured.	

**LPI 6338-21 SEQUENTIAL APPLICATION**

More consistent control of broadleaf and grass weeds may be obtained by an early preplant (surface-applied or shallow incorporated) application of LPI 6338-21, followed by a second preemergence application after planting but before soybean emergence. A sequential application will decrease the need for tillage and/or burndown herbicides for the control of existing vegetation before planting, while providing residual control of weeds after planting.

**Application**

An early preplant application may be made 15 to 30 days before planting soybeans. Follow this application with a preemergence overlay application of LPI 6338-21 after planting but before crop emergence. Follow directions on this label for sequential applications from 0 to 14 days before planting.

Where a rate range is recommended, use the higher rates:

- In fields with a history of severe weed pressure.
- When the time between early preplant and preemergence overlay applications

**LPI 6338-21 EPA REG. NO. 34704-XXXX**

approaches the maximum 30 days.

- When the organic matter content of the soil is over 3%.
- When heavy crop residues are present on the soil surface.

When weeds exceed 1 to 1-1/2 inches in height or diameter at application, use a burndown herbicide, such as Mad Dog, Makaze, Gramoxone Inteon or Whiteout.

**Weeds Controlled**

In addition to weeds controlled by LPI 6338-21 alone, the sequential application improves control of the following annual broadleaf weeds: buffalobur, cocklebur, common ragweed, velvetleaf, and sunflower.

**TABLE 8: LPI 6338-21 SEQUENTIAL USE RATES FOR REDUCED-TILL AND NO-TILL SYSTEMS (BROADCAST RATES)**

Soil Texture <sup>1</sup>	Early Preplant Application LPI 6338-21(Pt./A)	-Followed By -	Preemergence Overlay Application LPI 6338-21(Pt./A) <sup>2</sup>
COARSE <sup>1</sup> (Sand, loamy sand, sandy loam)	1.6 to 2.4	-followed by -	0.4 to 1.2
MEDIUM (Loam, silt loam, sandy clay loam, silt, sandy clay)	2.0 to 2.7	-followed by -	0.8 to 1.6
FINE (Silty clay loam <sup>3</sup> , clay loam, silty clay, clay)	2.4 to 3.5	-followed by -	1.2 to 2.0

<sup>1</sup>On coarse-textured soils, do not use on sandy soils with less than 1% organic matter. However, on coarse-textured soils with a calcareous surface area or a pH of 7.5 or higher, do not use on sand soils with less than 2% organic matter, or on loamy sand or sandy loam soils with less than 1% organic matter.  
<sup>2</sup>Total not to exceed 4.0 pints of LPI 6338-21 per acre per use season.  
<sup>3</sup>Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S. When using LPI 6338-21, treat this soil as "fine-textured."

**CROP ROTATION INTERVALS**

Full-rate application of products containing chlorimuron-ethyl (**Classic herbicide®**, etc.), chloransulam-methyl (**FirstRate®**), flumetsulam (**Hornet®**), imazaquin (**Scepter® 70 DG herbicide**) the same year as **LPI 6338-21** may increase the risk of injury to sensitive followcrops. Consult the product labels for recommended uses of these products in combinations.

Only rotational crops harvested at maturity may be used for feed or food.

**TABLE 9: CROP ROTATION INTERVALS**

CROP	Crop Rotation Intervals (Months)	CROP	Crop Rotation Intervals (Months)
Alfalfa	4.5	Peanuts	12
Asparagus	40	Peas	8
Bahiagrass <sup>6</sup>	40	Popcorn <sup>5</sup>	18
Barley, Spring (except North Dakota) <sup>2</sup>	9.5	Potatoes <sup>6</sup>	26
Barley, Winter (except North Dakota) <sup>2</sup>	9.5	Rice	40
Cabbage <sup>6</sup>	40	Root crops	40
Canola <sup>7</sup>	40	Rye	12
Cantaloupe <sup>6</sup>	40	Sainfoin	40
CLEARFIELD <sup>®</sup> Corn	8	Southern Peas	12
Clover	12	Sweet Pepper Transplants <sup>6</sup>	40
Cotton	18	Sweet Potato Transplants <sup>6</sup>	40
Cucumber <sup>6</sup>	40	Soybeans	0
Edible Beans	12	Sugarcane	40
Field Corn <sup>3,4</sup>	8.5	Sweet corn <sup>5</sup>	18
Field Corn (Seed) <sup>3,4</sup>	8.5	Tobacco	12
Flax	26	Tomatoes	40
Forage Grasses	40	Tomato Transplants <sup>6</sup>	40
Lentils	40	Watermelon <sup>6</sup>	40
Lettuce	18	Wheat, Spring	8
Lima	12	Wheat, Winter <sup>1</sup>	4.5
Oats	18	Other Crops not listed	40
Onion <sup>6</sup>	40	---	--

<sup>1</sup>If soybeans are furrow irrigated, till the soil prior to planting winter wheat. The beds should be broken up and the soil mixed with tillage equipment set to cut 4-6 inches deep.

<sup>2</sup>**Delaware, Indiana, Kentucky, Maryland, New Jersey, Ohio, Pennsylvania, and Virginia only:** Barley may be planted 4 months following a LPI 6338-21 application in these states.

**North Dakota only:** Barley may be planted 18 months following a LPI 6338-21 application.

<sup>3</sup>**Corn inbred lines:** Corn inbred seed lines may be planted the year following an application of LPI 6338-21. Several seed companies have tested a wide range of inbreds for sensitivity to LPI 6338-21 soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, Loveland Products, Inc. has not been given access to



the inbred data. Growers are directed to contact the seed company for information and directions regarding the planting of corn grown for seed in fields treated with LPI 6338-21 the previous year. **Since growing conditions, environmental conditions and grower practices are beyond the control of Loveland Products, Inc. all risks and consequences associated with planting seed corn inbreds into fields treated previously with LPI 6338-21 shall be assumed by the user.**

<sup>4</sup>**Arizona, Hawaii, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming only:** Field corn and field corn grown for seed may be planted 9½ months after LPI 6338-21 application.

<sup>5</sup>**Illinois, Indiana, Iowa, Minnesota, Ohio, Tennessee, and Wisconsin only:** Sweet corn and popcorn varieties may be planted the year following an application of LPI 6338-21. Some sweet corn and popcorn varieties may be injured when planted at less than 18 months following an application of LPI 6338-21. Before planting sweet corn for processing, contact the processor company for information and directions regarding the tolerance of sweet corn varieties planned for fields treated with LPI 6338-21 the previous year. **DO NOT** plant fresh market sweet corn varieties prior to 18 months after LPI 6338-21 use. Before planting popcorn, contact the popcorn company for information and directions regarding the tolerance of popcorn varieties planned for fields treated with LPI 6338-21 the previous year. **Since growing conditions, environmental conditions and grower practices are beyond the control of Loveland Products, Inc., to the extent consistent with applicable law, all risks and consequences associated with planting sweet corn or popcorn varieties into fields treated previously with LPI 6338-21 shall be assumed by the user.** Stunting and maturity delay or other adverse effects may result when sweet corn or popcorn are planted following LPI 6338-21 use.

<sup>6</sup>**Alabama, Delaware, Florida, Georgia, Indiana, Kentucky, Maryland, New Jersey, North Carolina, Pennsylvania, South Carolina, and Virginia only:** This crop may be planted 18 months following the last application of LPI 6338-21.

<sup>7</sup>**CLEARFIELD® Canola:** CLEARFIELD varieties of canola may be planted as a rotational crop the twelve months after an application of LPI 6338-21 at specified rates on soybeans.

26/29

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully dam up material to prevent runoff. Refer to **Precautionary Statements** on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed below. In spill or leak incidents, keep unauthorized people away. Maintaining a spill kit and fire extinguisher on hand and having emergency phone numbers posted will allow you to be prepared for emergencies

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:**

**Nonrefillable container.** Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at [www.acrecycle.org](http://www.acrecycle.org). If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**For packages up to 5 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:**

Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure**

**rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 56 gallons:** To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**For refillable containers:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY BEFORE BUYING OR USING THIS PRODUCT,** read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance

28/29

with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Accent, Assure, Basis, Classic, Harmony, Staple, and Synchrony are registered trademarks of E.I. DuPont de Nemours and Company.

Aim and Command are registered trademarks of FMC Corporation.

Arrow is a registered trademark of Makhteshim Agan of North America, Inc.

Basagran, Clearfield, Poast, Poast Plus, Pursuit, Rezult, and Scepter are registered trademarks of BASF.

Cobra and Resource are registered trademarks of Valent U.S.A. Corporation.

FirstRate, Frontrow, Hornet, Lorsban, and Python are registered trademarks of Dow AgroSciences LLC.

Flexstar, Fusilade, Fusion, Gramoxone Inteon, Reflex, Spirit, and Touchdown are registered trademarks of a Syngenta Group Company.

Intensity Post-emergence Grass Herbicide, Mad Dog, Makaze, and Stealth are registered trademarks and Whiteout is a trademark of Loveland Products, Inc.

Micro-Cel is a registered trademark of Celite Corporation.

Permit is a registered trademark of Nissan Chemical Industries, Ltd.

29/29

LPI 6338-21 EPA REG. NO. 34704-XXXX

Pioneer is a registered trademark of Pioneer Hi-Bred International.  
Roundup and Roundup Ready are registered trademarks of Monsanto Technology LLC.  
Storm and Ultra Blazer are registered trademarks of United Phosphorus, Inc.  
**Gramoxone Inteon and Lorsban 4E are restricted use pesticides.**

FORMULATED FOR



P.O. BOX 1286,  
GREELEY, COLORADO 80634-1286