



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 21, 2015

Chris Mason
Manager of Registrations
Loveland Products Inc.
P.O. Box 1286
Greeley, CO 80632-1286

Subject: Label Amendment – Adding language for North Dakota and Minnesota
Product Name: Matador-S
EPA Registration Number: 34704-1067
Application Date: September 23, 2014
Decision Number: 495983

Dear Dr. Mason:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 34704-1067
Decision No. 495983

with FIFRA section 6. If you have any questions, please contact Shanta Adeeb by phone at 703-347-0502, or via email at adeeb.shanta@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long horizontal flourish extending to the right.

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure




MATADOR®-S

GROUP 2 | 5 | 15 HERBICIDE

Herbicide for pre-emergent control of certain grasses and broadleaf weeds in soybeans.

ACTIVE INGREDIENTS:	By Wt
S-Metolachlor*	37.08%
Metribuzin**	8.23%
Imazethapyr***	1.83%
OTHER INGREDIENTS:	52.86%
	TOTAL 100.00%

*contains 3.38 pounds of S-metolachlor per gallon.
 **contains 0.75 pound of metribuzin per gallon.
 ***contains 0.17 pound of imazethapyr acid per gallon.



**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911, or call an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.
FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.

EPA REG. NO. 34704-1067

EPA EST. NO. 34704-MS-001

NET CONTENTS 2.5 GAL (9.46L)

EXP 09/14 MN ND

MATADOR®-S
EPA REG. NO. 34704-1067

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING

Harmful if absorbed through skin. Harmful if swallowed. Causes moderate eye irritation. Wear protective eyewear. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some people. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below.

Mixers, loaders, applicators, flaggers and other handlers must wear:

- Protective eyewear,
- Coveralls worn over short-sleeved shirt and short pants,
- Chemical-resistant gloves made out of butyl rubber \geq 14 mils or barrier laminate,
- Chemical-resistant footwear plus socks, and
- Chemical-resistant apron when cleaning equipment, spills, mixing and loading and when otherwise exposed to this product's concentrate.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove and wash contaminated clothing before reuse.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Surface Water Advisory

Metolachlor can contaminate surface water through ground spray drift. Under some conditions, metolachlor may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These conditions include: poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

Ground Water Advisory

Metribuzin is a chemical which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Metribuzin has been found in ground water as a result of agricultural use. Users are advised not to apply metribuzin where the water table (ground water) is close to the surface, and where the soils are very permeable, i.e., well-drained soils such as loamy sands. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Imazethapyr has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Metolachlor has the potential to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

MATADOR®-S
EPA REG. NO. 34704-1067

Mixing/Loading Instructions

Product must be used in a manner which will prevent back-siphoning in wells, spills or improper disposal of excess pesticide spray mixture.

This product may not be mixed or loaded within 50 feet of any wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes or reservoirs. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times.

The above specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

In New York State - not for sale or use in Nassau and Suffolk Counties.

Observe all restrictions, precautions and limitations on this label and on the labels of products used in combination with this product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protections of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protections Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Protective Eyewear.
- Coveralls over short-sleeve shirt and short pants,
- Chemical-resistant gloves, such as butyl rubber \geq 14 mils or barrier laminate,
- Chemical-resistant footwear plus socks.

PRODUCT INFORMATION

Matador®-S is an emulsifiable concentrate that kills weeds by root and/or foliage uptake and rapid translocation to the growing points. Adequate soil moisture is important for optimum activity of this product. When adequate soil moisture is present, this product will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

Occasionally, internode shortening and/or temporary yellowing of crop plants may occur following applications of this product. These effects occur infrequently and are temporary. Normal growth and appearance should resume within 1 to 2 weeks.

When organophosphate (such as Lorsban®) or carbamate insecticides are tank mixed with this product, temporary injury may result to the treated crops.

Use of this product in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product.

MATADOR®-S
EPA REG. NO. 34704-1067

Therefore, rotational crop injury is always possible.

Under some conditions (such as heavy texture soil, high organic matter, low pH or low rainfall), this product may cause injury to subsequent planted crops. Vegetable crops (particularly sugar beets) are sensitive to residues of this product in the soil.

Naturally occurring biotypes* of some of the weeds listed on this label may not be effectively controlled by this and/or other products with either the ALS/AHAS enzyme inhibiting mode of action. Other herbicides with the ALS/AHAS enzyme inhibiting mode of action include the sulfonyleureas (e.g., Accent®, Basis®, Classic®, Harmony® GT, Spirit®, Permit®, etc.), the sulfonamides (e.g., FirstRate®, etc.) and the pyrimidyl benzoates (e.g. Staple®, etc.).

* A weed biotype is a naturally occurring plant within a given species that has a slightly different, but distinct, genetic makeup from other plants.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to <http://www.epa.gov/espp/litstatus/wtc>.

SPRAY DRIFT MANAGEMENT

Preventing spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to prevent off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they must be observed.

The applicator must be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity and Temperature Inversions).

Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Do not make applications at a height greater than 10 feet above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up-and-downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

MATADOR®-S
EPA REG. NO. 34704-1067

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind directions and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Because drift potential is high, do not apply during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

This product may only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

APPLICATION WITH HERBICIDE SPRAY EQUIPMENT

Use a standard low-pressure (20 to 40 psi) herbicide boom sprayer equipped with suitable nozzles and screens no finer than 50-mesh in nozzle and in-line strainers. Agitate thoroughly before and during application with bypass agitation. Low pressure and high volume hand wand equipment is prohibited.

Ground Application

Apply the proper rate of this product in a minimum of 10.0 to 40.0 gallons of spray mixture per acre broadcast.

Aerial Application

Where permitted, apply specified rate in a minimum of 5.0 gallons of spray mixture per acre. Do not apply aerially when wind speed is greater than 10 mph.

For All Applications

Sprayer must be accurately calibrated before applying this product. Check sprayer during application to be sure it is working properly and delivering a uniform spray pattern. As the volume of spray mixture decreases per acre, the importance of accurate calibration and uniform application increases.

Avoid overapplication, misapplication, and boom and spray swath overlapping that will increase spray dosage. (Crop injury may occur as a result.) Avoid spray skips and gaps which allow weeds to grow in untreated soil. Do not apply when weather conditions favor spray drift and/or when sensitive or cool season crops (such as cole crops, onions, peas, or strawberries) are present in adjacent fields or in areas where wheat is growing in coarse textured soils.

Sprayer Cleanup

Spray equipment must be thoroughly cleaned to remove remaining traces of herbicide that might injure other crops to be sprayed. Drain any remaining spray solution of this product from the spray tank and dispose of according to label disposal instructions. Rinse the spray tank and refill with water, adding a heavy-duty detergent at the rate of 1 cup per 20.0 gallons of water. Recycle this mixture through the equipment for 5 minutes and spray out. Repeat this procedure twice. Fill the spray tank with clean water, recycle for 5 minutes, and spray out. Clean pump and nozzle screens thoroughly. Wash away any spray mixture from the outside of spray tank, nozzles or spray rig. All rinse water must be disposed of in compliance with local, state, and Federal guidelines.

MIXING INSTRUCTIONS

Incorporation and Combination Uses

When this product is to be used in combination with another herbicide, follow the most restrictive directions on all product labels for combinations, rates, crops, incorporation, and special precautions.

When using this product, make sure the sprayer is completely clean, free of rust or corrosion which occurs from winter storage. Examine strainers and screens to be sure the sprayer is clean from previously used pesticides.

Any tank mix containing this product must be kept agitated and sprayed out immediately. Do not allow tank mixes to stand for prolonged periods of time.

MATADOR®-S
EPA REG. NO. 34704-1067

The proper mixing procedure for this product alone or in tank-mix combinations with other herbicides is:

1. Fill the spray tank 1/4 to 1/3 full with clean water.
2. Add specified rate of this product while recirculating and with agitator running.
3. Mix thoroughly and add clean water to fill spray tank to desired level.
4. Add the other herbicide to tank last and agitate thoroughly.
5. Continue agitation during application and until sprayer tank is empty.

Application of Matador-S In Fluid Fertilizers

This product may be applied in fluid fertilizer solutions by following the appropriate mixing procedures and compatibility check. When using tank mix combinations, be sure all components are compatible.

Tank Mixing Guidelines for Fluid Fertilizer Mixtures

1. Add the required amount of water and compatibility agent (if required) to the tank. Start agitation system while adding this product and follow by adding the fluid fertilizer and agitate.
 2. If a second herbicide is also to be used, follow as above in Step 1, but use twice the amount of water. Start agitation, add Matador-S. Follow by adding the second herbicide, then continue filling the tank with fluid fertilizer.
 3. Maintain continuous agitation to assure uniform spray mixture until the tank is emptied.
- Make compatibility checks of this product plus fluid fertilizers and tank-mix combinations plus fluid fertilizers which include this product for each batch because of the variability of fluid fertilizers.

The following compatibility check should only be used when mixing with fluid fertilizers.

1. Pre-mix 8.0 teaspoons of water with 2.0 teaspoons of this product (4:1 ratio) in a quart jar by adding the water first and following with this product. Mix thoroughly. If a second herbicide is to be used, double the amount of water (8:1 ratio), mix in this product, and follow with the second herbicide.
2. Then pour 1.0 pint of fluid fertilizer into the quart jar and shake well.
3. Allow to stand for 5 minutes.

Interpretation of Results

If the solution in the jar appears to be uniform, without signs of agglomeration, or without a separation of an oily film on top of the fertilizer, the mixture may be used. If not, repeat the compatibility check using twice the amount of water or add a compatibility agent to the water. If separation occurs, but the mixture can be resuspended by shaking, then application is possible with good agitation in the spray tank.

SOYBEAN APPLICATION DIRECTIONS
(Except California)

This product may be applied preplant incorporated, preplant surface or pre-emergent surface, or as a sequential pre-emergent application. This product may also be used as an overlay application following a preplant incorporated application of a grass herbicide registered for this same use and in tank mix combinations for burndown weed control. All applications may be applied with ground equipment, and some may be applied with aerial spray equipment.

Restrictions

- Do not apply more than 3.0 pints of Matador-S per acre per use season.
- In North Dakota, and in Minnesota north of Highway #210, do not apply more than 2.2 pints of Matador-S per acre per use season.
- Do not harvest within 85 days of the last application of Matador-S.
- Do not graze or feed treated soybean forage, hay or straw to livestock.
- Do not rotate any crop not listed on this label for 40 months following application.
- Do not rotate to food or feed crops other than those listed on this label.
- Do not apply Matador-S preemergence to soybeans in California.
- Do not incorporate into soil or apply more than once per season except where permitted as part of a sequential application.
- Do not allow sprays to drift onto adjacent desirable plants.
- Do not apply this product through any type of irrigation system
- Do not apply under conditions which favor runoff or wind erosion of soil containing this product to non-target areas.
- To prevent off-site movement due to run-off or wind erosion:
The soil surface must first be settled by rainfall or irrigation if treating powdery dry or light sand soils where conditions are favorable for wind erosion.
Do not apply to impervious substrates such as paved or highly compacted surfaces.
Do not use tailwater from the first flood or furrow irrigation of treated fields to treat non-target crops unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.
- Do not apply using low-pressure and high-volume hand-wand equipment.
- Observe all restrictions, precautions and limitations on labeling of all products used in mixtures.

MATADOR®-S
EPA REG. NO. 34704-1067

Soil Texture and Rate Ranges

As used on this label, "Coarse soils" are loamy sand or sandy loam soils. "Medium soils" are loam, silt loam, silt, sandy clay, or sandy clay loam. "Fine soils" are silty clay, silty clay loam, clay, or clay loam. Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

Where a rate range is shown, use a lower rate on soils that are coarse-textured or low in organic matter. Use a higher rate on soils that are relatively fine-textured or high in organic matter.

Precautions

Injury to soybeans may occur when this product is used under the following conditions:

1. When soils have a calcareous surface area or a pH of 7.5 or higher.
2. When applied in conjunction with soil-applied organic phosphate pesticides.
3. With over-application or boom overlapping, which may result in stand loss and soil residues.
4. With uneven application or improper incorporation, which can decrease the level of weed control and/or increase the level of injury.
5. When applied to any soil with less than 0.5% organic matter.
6. When soil incorporation is deeper than is listed.
7. When sprayers are not calibrated accurately.
8. When heavy rains occur soon after application, especially in poorly drained areas where water may stand for several days.
9. When soybeans are planted less than 1-1/2 inches deep, particularly in preemergence application.
10. Where high soil levels of atrazine are present.
11. When using poor quality soybean seed.

Certain soybean varieties are sensitive to Metribuzin. Prior to use of this product, consult your soybean seed supplier for more information on the tolerance of soybean varieties to Matador-S.

Activation

A minimum amount of soil moisture is required to activate this product. In areas of low rainfall, follow preemergence applications to dry soil with light irrigation of 1/4 acre-inch of water. Do not apply heavy irrigation immediately after application. As with many surface-applied herbicides, weed control and crop tolerance may vary with rainfall and/or soil texture.

Replanting

If replanting is necessary in fields treated with this product as directed on this label, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. Do not apply more than once per season except where permitted as part of a sequential application as injury to soybeans may occur. Maximum application rate is 3.0 pints of Matador-S per acre per use season. Do not exceed this amount in any use pattern: single application, replant or sequential application.

TABLE 1: ANNUAL BROADLEAF WEEDS CONTROLLED BY MATADOR-S

C = Control S = Suppression or Erratic Control P = Poor or No Control U= Unknown

Weed Controlled	Level of Control
Bristly starbur (<i>Acanthospermum hispidum</i>)	C
Buffalobur (<i>Solanum rostratum</i>)	C
Carpetweed (<i>Mollugo verticillata</i>)	C
Cocklebur (<i>Xanthium pensylvanicum</i>)	S
Common chickweed (<i>Stellaria media</i>)	C
Copperleaf, Hophornbeam (<i>Acalypha ostryifolia</i>)	C
Field pennycress (<i>Thlaspi arvense</i>)	C
Florida beggarweed (<i>Desmodium tortuosum</i>)	C
Florida pusley (<i>Richardia scabra</i>)	C
Galinsoga (<i>Galinsoga</i> spp.)	C
Henbit (<i>Lamium amplexicaule</i>)	C
Horseweed (Marestail) (<i>Conyza canadensis</i>)	U
Jimsonweed (<i>Datura stramonium</i>)	C
Knotweed (<i>Polygonum</i> spp.)	C
Kochia (<i>Kochia scoparia</i>)	C
Lambsquarters (<i>Chenopodium</i> spp.)	C
Marshelder (<i>Iva Annua</i>)	C
Morningglory	
Entireleaf (<i>Ipomoea hederacea</i> var. <i>integriuscula</i>)	S
Ivyleaf (<i>Ipomoea hederacea</i>)	S
Pitted (<i>Ipomoea lacunosa</i>)	S
Smallflower (<i>Jacquemontia tamnifolia</i>)	C
Tall (<i>Ipomoea purpurea</i>)	S
Mustard spp.	C

MATADOR®-S
EPA REG. NO. 34704-1067

Table 1: Annual Broadleaf Weeds Controlled by Matador-S cont'd.:

C = Control S = Suppression or Erratic Control P = Poor or No Control U= Unknown

Weed Controlled	Level of Control
Nightshade	
Black (<i>Solanum nigrum</i>)	C
Eastern black (<i>Solanum ptycanthum</i>)	C
Hairy (<i>Solanum villosum</i>)	C
Pigweed	
Redroot (<i>Amaranthus retroflexus</i>)	C
Smooth (<i>Amaranthus hybridus</i>)	C
Spiny (<i>Amaranthus spinosus</i>)	C
Poinsettia, wild (<i>Euphorbia cyathophora</i>)	C
Prickly lettuce (<i>Lactuca serriola</i>)	C
Prickly sida/Teaweed (<i>Sida spinosa</i>)	C
Puncturevine (<i>Tribulus terrestris</i>)	C
Purslane (<i>Portulaca oleracea</i>)	C
Ragweed	
Common (<i>Ambrosia artemisiifolia</i>)	C
Giant (<i>Ambrosia trifida</i>)	S
Redweed	C
Russian thistle (<i>Salsola kali</i>)	C
Sage, barnyard	S
Sesbania (<i>Sesbania</i> spp.)	C
Shepherd's-purse (<i>Capsella bursa-pastoris</i>)	C
Sicklepod (<i>Cassia obtusifolia</i>) ¹	C
Smartweeds (<i>Polygonum</i> spp.)	
Ladysthumb (<i>Polygonum persicaria</i>)	C
Pennsylvania (<i>Polygonum pennsylvanicum</i>)	C
Spurge	
Prostrate (<i>Euphorbia humistrata</i>)	C
Spotted (<i>Euphorbia maculata</i>)	C
Spurred anoda (<i>Anoda cristata</i>)	C
Sunflower (<i>Helianthus</i> spp.)	C
Velvetleaf (<i>Abutilon theophrasti</i>)	C
Venice mallow (<i>Hibiscus trionum</i>)	C
Virginia pepperweed (<i>Lepidium virginicum</i>)	C
Waterhemp (<i>Amaranthus rudis</i>)	C
Wild mustards (<i>Brassica</i> spp.)	C

¹For maximum control of sicklepod, use a preemergence application.

TABLE 2: ANNUAL GRASSES AND SEDGES CONTROLLED BY MATADOR-S

C = Control S = Suppression or Erratic Control P = Poor or No Control

Weed Controlled	Level of Control
Barnyardgrass (<i>Echinochloa crus-galli</i>)	C
Bluegrass (<i>Poa annua</i>)	C
Broadleaf signalgrass (<i>Brachiaria platyphylla</i>)	C
Browntop millet (<i>Panicum ramosa</i>)	C
Crabgrass (<i>Digitaria</i> spp.)	C
Crowfootgrass (<i>Dactyloctenium aegyptium</i>)	C
Cupgrass (<i>Eriochloa gracilis</i>)	C
Foxtails (<i>Setaria</i> spp.)	C
Goosegrass (<i>Eleusine indica</i>)	C
Johnsongrass, Seedling (<i>Sorghum halepense</i>)	C
Junglerice (<i>Echinochloa colona</i>)	C
Millet, wild-proso (<i>Panicum miliaceum</i>)	S
Nutsedge	
Yellow (<i>Cyperus esculentus</i>)	S
Purple (<i>Cyperus rotundus</i>)	S
Panicum, fall (<i>Panicum dichotomiflorum</i>)	C
Panicum, Texas (<i>Panicum, texanum</i>)	S
Red rice (<i>Oryza sativa</i>)	C
Sandbur (<i>Cenchrus</i> spp.)	S

MATADOR®-S
EPA REG. NO. 34704-1067

Table 2: Annual Grasses and Sedges Controlled by Matador-S cont'd.:

Weed Controlled	Level of Control
Shattercane (<i>Sorghum bicolor</i>)	S
Sorghum, volunteer (<i>Sorghum</i> spp.)	S
Sprangletop (<i>Leptochloa</i> spp.)	P
Stinkgrass (<i>Eragrostis</i> spp.)	P
Wheat, volunteer (<i>Triticum</i> spp.)	S
Witchgrass (<i>Panicum capillare</i>)	C

MATADOR-S USE RATES FOR COVENTIONAL TILLAGE SYSTEMS

Matador-S used alone in Preplant Incorporated Application

Incorporate Matador-S uniformly into the top 2 inches of soil within 14 days before planting using a disk, field cultivator, rolling cultivator or similar equipment. Use incorporated application if furrow irrigation is used or when a period of dry weather after application is expected.

Matador-S used alone in Preemergence Application

When used alone, Matador-S can be applied broadcast by ground or aerially. This application may be made during planting or as a separate operation after planting but must be made before crop emergence. If dry weather follows preemergence application, cultivate uniformly with shallow tilling equipment that will not damage soybeans.

Restrictions

Do not apply to sand soils, or to sandy loam or loamy sand soils containing less than 2% organic matter.

TABLE 3: MATADOR-S RATES WHEN USED ALONE IN PREPLANT OR PREEMERGENCE APPLICATION

***(In North Dakota, and in Minnesota north of Highway #210, do not apply more than 2.2 pints of Matador-S per acre per use season.)**

Soil Texture		Organic Matter		
		0.5 to 2.0%	2.0 to 3.0%	Over 3.0% ³
		Pints of Matador-S Per Acre		
Coarse Soils¹	Sandy loam	1.9 to 2.4*	1.9 to 2.4*	2.4 to 2.9*
	Loamy sand	n/a ¹	1.9 to 2.4*	2.4 to 2.9*
Medium Soils (Loam, silt loam, silt, sandy clay, sandy clay loam)		3.0*		
Fine Soils (Silty clay, silty clay loam ² , clay, clay loam)				

¹ Do not use on sand soils. On coarse-textured soils, do not use on loamy sand with less than 2% organic matter.

² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S.

³ For preplant incorporated application, use the lower rate.

⁴ For Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, see section below **In Coarse (Light) Soils**.

In Coarse (Light) Soils

(Only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia)

This product can be used at the rates specified in Table 4 as a preplant incorporated or preemergence application in coarse-textured, low organic matter soils in the states listed above. Refer to Table 4 and to the appropriate sections of this label for specific directions on use and restrictions.

TABLE 4: MATADOR-S RATES WHEN USED ALONE IN PREPLANT OR PREEMERGENCE APPLICATIONS ON COARSE SOILS (Only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Missouri, North Carolina, Oklahoma, South Carolina, Tennessee, exas, Virginia)

Soil Texture		Organic Matter	
		0.5 to 1.0%	1.0% or above
		Pints of Matador-S Per Acre ²	
Coarse Soils	Sand	n/a ¹	1.9 to 3.0
	Sandy loam, loamy sand	1.9 to 3.0	1.9 to 3.0

¹ Do not use on sand with less than 1% organic matter.

² Use the higher rate under heavy weed pressure and/or soils higher in organic matter.

MATADOR®-S
EPA REG. NO. 34704-1067

HERBICIDES THAT MAY BE APPLIED POSTEMERGENCE FOLLOWING MATADOR-S

If required, application of this product alone or in tank mixture may be followed by an application of a postemergence herbicide to provide additional control of certain weeds. The following postemergence herbicides may be applied:

Aim®	FirstRate®	Harmony® GT XP	Storm®
Arrow®	Flexstar®	Intensity®	Synchrony® XP ²
Assure® II	Fusilade® DX	Poast®	Ultra Blazer®
Basagran®	Frontrow®	Poast Plus®	
Classic®	Fusion®	Reflex®	
Cobra®	glyphosate herbicides ¹ (such as Makaze or Mad Dog)	Resource®	
		Rezult® A&B	

¹ Use on Roundup Ready® or glyphosate tolerant soybean varieties only.

² Use on STSTM soybean varieties only.

Refer to the **Directions for Use** on this label and the individual product labels for use directions, use rates, and special precautions and/or restrictions.

BURNDOWN WEED CONTROL

This product can be used as part of a burndown herbicide program for control of existing vegetation prior to soybean emergence in conservation tillage (reduced-tillage/no-till) systems. This product may be tank mixed with a 2,4-D low volatile ester (LVE) (such as Whiteout®) and/or glyphosate herbicides (such as Mad Dog® and Makaze® brands) for control of emerged weeds prior to crop emergence. Burndown tank mixes with Matador-S can be applied before planting or prior to crop emergence.

DO NOT tank mix Matador-S with clomazone containing herbicides (Command®).

Application

This product may be applied up to 30 days before planting or preemergence. Apply only by ground equipment when this product is used for burndown of existing vegetation in conservation tillage systems. Use the high end of the rate range for applications of this product made 14 to 30 days before planting. Refer to Tables 3 and 4 for rates of Matador-S alone and to Table 5 for rates of tank mix partners.

TABLE 5: RATES OF TANK MIX PARTNERS TO BE USED IN COMBINATION WITH MATADOR-S FOR BURNDOWN APPLICATIONS

Product	Rate of Tank Mix Partner	Directions and Remarks
2,4-D LVE (Whiteout 2,4-D)	0.25 to 1.0 lb AE ¹ /A	Apply at least 7 days preplant when using Whiteout at 0.25 to 0.5 lb AE ¹ /A and at least 30 days preplant with rates greater than 0.5 lb AE ¹ /A. Include crop oil concentrate (COC) at the rate of 1.0 gal/100 gal of spray solution (1% v/v).
Glyphosate (Mad Dog, or Makaze brands)	Refer to product label for use rates.	Must be applied prior to crop emergence. Use the higher rates within the specified range as weeds approach the maximum weed heights listed in Table 6. Apply in 10.0 to 20.0 gal of water/A. Refer to the Mad Dog or Makaze label for spray adjuvant instructions. Any glyphosate formulation registered and labeled for use in soybeans may be tank mixed with this product.
Glyphosate (Mad Dog or Makaze brands) + 2,4-D LVE (Whiteout 2,4-D)	Refer to the product label for use rates + 0.25 lb AE ¹ /A.	Follow the Directions and Remarks section above for Whiteout 2,4-D and Mad Dog/Makaze, paying special attention to planting restrictions with Whiteout. Refer to the Mad Dog or Makaze label for spray adjuvant instructions. Do not use crop oil concentrate (COC).

¹AE = 2,4-D acid equivalent

Restrictions

Do not apply these treatments after crop emergence. Observe all restrictions, precautions and limitations on the labeling of all products used in tank mixtures.

- Apply only 2,4-D LVE formulations (such as Whiteout 2,4-D) that are registered and specified for preplant or burndown use.
- Do not apply tank mixtures containing 2,4-D LVE (Whiteout 2,4-D) if wind is blowing toward desired susceptible plants (i.e., cotton, tobacco, tomato, etc.) or when wind speeds exceed 6 miles per hour. Observe all precautions and limitations of all products used in tank mixtures.

Follow the most restrictive preharvest interval of all products used in a tank mixture.

MATADOR®-S
EPA REG. NO. 34704-1067

Weeds Controlled

Matador-S in tank mixtures with the herbicides listed in Table 6 will provide burndown control of the weeds listed below.

TABLE 6: WEEDS CONTROLLED WITH TANK MIXES OF MATADOR-S IN BURNDOWN APPLICATION

Weeds Controlled	Whiteout 2,4-D	Mad Dog/Makaze	Mad Dog/Makaze + Whiteout 2,4-D
Annual Grasses	Maximum Burndown Height (Inches)		
Barley	Does not improve control of these species		8
Barnyardgrass			6
Crabgrass spp.			6
Foxtail spp.			8
Johnsongrass, seedling			8
Panicum, fall			6
Sandbur, field			8
Wheat, volunteer			6
Witchgrass			6
Broadleaves		Maximum Burndown Height (Inches)	
Buffalobur	-	6	6
Chickweed, common	6	6	6
Cocklebur, common	6	6	8
Dandelion, common	6 dia ¹	2 dia ²	6 dia ¹
Henbit	4	4	4
Horseweed (Marestail)	6 ¹	4 ²	6
Jimsonweed	6	6	6
Kochia	4 ¹	4	4
Ladysthumb	6	6	8
Lambsquarters, common	6	6	8
Lettuce, prickly	6	4	6
Mallow, Venice	6	6	6
Morningglory spp.	6	2	4
Mustard spp.	6	6	8
Pennycress, field	6	6	6
Pigweed spp. (annual)	6	6	8
Ragweed, common	6	6 ²	8
Ragweed, giant	6 ¹	4 ²	6
Shepherd's-purse	6	6	6
Sida, prickly	6	4	4
Smartweed, Pennsylvania	6	6	8
Sunflower, common	6	6	6
Thistle, Russian	4 ¹	2 to 4 ²	4
Velvetleaf	6	6	8
Waterhemp spp.	6	6	8

¹Use Whiteout 2,4-D at 0.5 pound active ingredient per acre.

²Use a minimum of 0.75 pound active ingredient per acre of Mad Dog or Makaze.

MATADOR-S USE RATES FOR REDUCED- AND NO-TILL SYSTEMS

Preplant Surface Application

Matador-S may also be used in reduced-till and no-till systems. Applications may be made up to 30 days before planting or after planting, but before soybean emergence. Residual herbicides such as Canopy®, FirstRate, Command, Python®, and Stealth® may be tank mixed for additional weed control. If weeds are present at time of application, burndown herbicides may be added to the tank mixes (see **Burndown Weed Control** section). Refer to the tank mix partner product labels for specific rates and use directions.

TABLE 7: MATADOR-S RATES FOR REDUCED AND NO-TILL SYSTEMS

**(In North Dakota, and in Minnesota north of Highway #210, do not apply more than 2.2 pints of Matador-S per acre per use season.)*

Soil Texture		Organic Matter		
		Up to 2.0%	2.0 to 3.0%	3.0% or above
Coarse ¹	Sand, loamy sand	n/a ¹	1.9	3.0*
	Sandy loam	1.9	1.9	3.0*
Medium (Loam, silt loam, silt, sandy clay, sandy clay loam)			3.0*	
Fine (Silty clay, silty clay loam ² , clay, clay loam)			3.0*	

Cont'd. next page

MATADOR®-S

EPA REG. NO. 34704-1067

- ¹ Do not use on sand soils. On coarse-textured soils, do not use on loamy sand soils with less than 2% organic matter.
- ² Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S. When using Matador-S, treat this soil as fine-textured.

MATADOR-S SEQUENTIAL APPLICATION

More consistent control of broadleaf and grass weeds may be obtained by an early preplant (surface-applied or shallow incorporated) application of Matador-S, followed by a second preemergence application after planting but before soybean emergence. A sequential application will decrease the need for tillage and/or burndown herbicides for the control of existing vegetation before planting, while providing residual control of weeds after planting.

Application

An early preplant application may be made 15 to 30 days before planting soybeans. Follow this application with a preemergence overlay application of Matador-S after planting but before crop emergence. Follow directions on this label for sequential applications from 0 to 14 days before planting.

Where a rate range is listed, use the higher rates:

- In fields with a history of severe weed pressure.
- When the time between early preplant and preemergence overlay applications approaches the maximum 30 days.
- When the organic matter content of the soil is over 3%.
- When heavy crop residues are present on the soil surface.

When weeds exceed 1.0 to 1.5 inches in height or diameter at application, use a burndown herbicide, such as Mad Dog, Makaze, Gramoxone Inteon or Whiteout.

Weeds Controlled

In addition to weeds controlled by Matador-S alone, the sequential application improves control of the following annual broadleaf weeds: buffalobur, cocklebur, common ragweed, velvetleaf, and sunflower.

TABLE 8: MATADOR-S SEQUENTIAL USE RATES FOR REDUCED-TILL AND NO-TILL SYSTEMS (BROADCAST RATES)

*** (In North Dakota, and in Minnesota north of Highway #210, do not apply more than 2.2 pints of Matador-S per acre per use season.)**

Soil Texture ¹		Early Preplant Application Matador-S (Pt/A)	- Followed By -	Preemergence Overlay Application Matador-S (Pt/A) ²
Coarse ¹ (Sand, loamy sand, sandy loam)	Organic Matter	--	--	--
	0.5 to 3.0%	1.2 to 1.8	- followed by -	1.2 to 0.6*
	Over 3.0%	1.5 to 2.0	- followed by -	1.5 to 0.9*
Medium (Loam, silt loam, sandy clay loam, silt, sandy clay)		1.5 to 2.0	- followed by -	1.5 to 1.0*
Fine (Silty clay loam ³ , clay loam, silty clay, clay)		1.5 to 2.0	- followed by -	1.5 to 1.0*

¹ On coarse-textured soils, do not use on sandy soils with less than 1% organic matter. On coarse-textured soils with a calcareous surface area or a pH of 7.5 or higher, do not use on sand soils with less than 2% organic matter, or on loamy sand or sandy loam soils with less than 1% organic matter.

² Total not to exceed 3.0 pints of Matador-S per acre per use season. For coarse soils do not exceed total rates in Table 3.

³ Silty clay loam soils are transitional soils and may be classified as medium-textured soils in some regions of the U.S. When using Matador-S, treat this soil as "fine-textured."

CROP ROTATION INTERVALS

Full-rate application of products containing chlorimuron-ethyl (Classic Herbicide, etc.), chloransulam-methyl (FirstRate), flumetsulam (Hornet®), imazaquin (Scepter® 70 DG Herbicide) the same year as Matador-S may increase the risk of injury to sensitive followcrops. Consult the product labels for listed uses of these products in combinations.

Only rotational crops harvested at maturity may be used for feed or food.

MATADOR®-S
EPA REG. NO. 34704-1067

TABLE 9: CROP ROTATION INTERVALS

Crop	Crop Rotation Intervals (Months)	Crop	Crop Rotation Intervals (Months)
Alfalfa	4.5	Popcorn ⁵	18
Asparagus	40	Potatoes ⁶	26
Bahiagrass ⁶	40	Rice	40
Barley, spring (except North Dakota) ²	9.5	Root crops	40
Barley, winter (except North Dakota) ²	9.5	Rye (except in North Dakota, and in Minnesota north of Highway #210)	12
Cabbage ⁶	40	Rye in North Dakota, and in Minnesota north of Highway #210	18
Canola ⁷	40	Sainfoin	40
Cantaloupe ⁶	40	Safflower	18
Clearfield® Corn	8	Southern peas	12
Clover	12	Sorghum	18
Cotton	18	Soybeans	0
Cucumber ⁶	40	Sugarcane	40
Edible beans	12	Sunflower	18
Field corn ^{3,4}	8.5	Sweet corn ⁵	18
Field corn (seed) ^{3,4}	8.5	Sweet pepper transplants ⁶	40
Flax	26	Sweet potato transplants ⁶	40
Forage grasses	40	Tobacco	12
Lentils	40	Tomatoes	40
Lettuce	18	Tomato transplants ⁶	40
Lima	12	Watermelon ⁶	40
Oats	18	Wheat, spring	8
Onion ⁶	40	Wheat, winter ¹	4.5
Peanuts	12	Other crops not listed	40
Peas	8		

¹ If soybeans are furrow irrigated, till the soil prior to planting winter wheat. The beds should be broken up and the soil mixed with tillage equipment set to cut 4 to 6 inches deep.

² **Delaware, Indiana, Kentucky, Maryland, New Jersey, Ohio, Pennsylvania, and Virginia only:** Barley may be planted 4 months following a Matador-S application in these states.

North Dakota only: Barley may be planted 18 months following a Matador-S application.

³ **Corn inbred lines:** Corn inbred seed lines may be planted the year following an application of Matador-S. Several seed companies have tested a wide range of inbreds for sensitivity to Matador-S soil residues and have reported good crop safety. However, due to the proprietary nature of seed production, Loveland Products, Inc. has not been given access to the inbred data. Growers are directed to contact the seed company for information and directions regarding the planting of corn grown for seed in fields treated with Matador-S the previous year. **Since growing conditions, environmental conditions and grower practices are beyond the control of Loveland Products, Inc. all risks and consequences associated with planting seed corn inbreds into fields treated previously with Matador-S shall be assumed by the user.**

⁴ **Arizona, Hawaii, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming only:** Field corn and field corn grown for seed may be planted 9-1/2 months after Matador-S application.

⁵ **Illinois, Indiana, Iowa, Minnesota, Ohio, Tennessee, and Wisconsin only:** Sweet corn and popcorn varieties may be planted the year following an application of Matador-S. Some sweet corn and popcorn varieties may be injured when planted at less than 18 months following an application of Matador-S. Before planting sweet corn for processing, contact the processor company for information and directions regarding the tolerance of sweet corn varieties planned for fields treated with Matador-S the previous year. **DO NOT** plant fresh market sweet corn varieties prior to 18 months after Matador-S use. Before planting popcorn, contact the popcorn company for information and directions regarding the tolerance of popcorn varieties planned for fields treated with Matador-S the previous year. **Since growing conditions, environmental conditions and grower practices are beyond the control of Loveland Products, Inc., to the extent consistent with applicable law, all risks and consequences associated with planting sweet corn or popcorn varieties into fields treated previously with Matador-S shall be assumed by the user. Stunting and maturity delay or other adverse effects may result when sweet corn or popcorn are planted following Matador-S use.**

⁶ **Alabama, Delaware, Florida, Georgia, Indiana, Kentucky, Maryland, New Jersey, North Carolina, Pennsylvania, South Carolina, and Virginia only:** This crop may be planted 18 months following the last application of Matador-S.

⁷ **Clearfield Canola:** Clearfield varieties of canola may be planted as a rotational crop the 12 months after an application of Matador-S at specified rates on soybeans.

MATADOR®-S
EPA REG. NO. 34704-1067

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully dam up material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed below. In spill or leak incidents, keep unauthorized people away. Maintaining a spill kit and fire extinguisher on hand and having emergency phone numbers posted will allow you to be prepared for emergencies

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable container. Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at 1-877-952-2272 or www.acrecycle.org. If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For nonrefillable containers up to 5 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For nonrefillable containers greater than 5 gallons and less than 56 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For nonrefillable containers greater than 56 gallons: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For refillable containers from 55 to 330 gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Offer for recycling or puncture and dispose of in a sanitary landfill or by incineration.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such

MATADOR®-S
EPA REG. NO. 34704-1067

Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Accent, Assure, Basis, Canopy, Classic, Harmony, Staple, and Synchrony are registered trademarks of E.I. DuPont de Nemours and Company.

Aim and Command are registered trademarks of FMC Corporation.

Arrow is a registered trademark of Makhteshim Agan of North America, Inc.

Basagran, Clearfield, Poast, Poast Plus, Pursuit, Rezult, and Scepter are registered trademarks of BASF.

Cobra and Resource are registered trademarks of Valent U.S.A. Corporation.

FirstRate, Frontrow, Hornet, Lorsban, and Python are registered trademarks of Dow AgroSciences LLC.

Flexstar, Fusilade, Fusion, Gramoxone Inteon, Reflex, Spirit, and Touchdown are registered trademarks of a Syngenta Group Company.

Intensity Post-emergence Grass Herbicide, Mad Dog, Makaze, Matador, Stealth and Whiteout are registered trademarks Loveland Products, Inc.

Permit is a registered trademark of Nissan Chemical Industries, Ltd.

Roundup and Roundup Ready are registered trademarks of Monsanto Technology LLC.

Storm and Ultra Blazer are registered trademarks of United Phosphorus, Inc.

FORMULATED FOR
LOVELAND PRODUCTS, INC.
P.O. BOX 1286, GREELEY, COLORADO 80632-1286