

34704-1075

04/10/2013

1/18



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7504P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

EPA Reg. Number:  
34704-1075

Date of Issuance:  
APR 10 2013

NOTICE OF PESTICIDE:  
 Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance: Unconditional

Name of Pesticide Product:  
Loveland Captan 80 DF

Name and Address of Registrant (include ZIP Code):

Robert Avalos  
Loveland Products, Inc.  
P.O. Box 1286  
Greeley, CO  
80632-1286

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5), provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA Section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for re-registration of your product under FIFRA Section 4.
2. You must submit two copies of a final printed label within 30 days from the date of this notice (along with a copy of this notice) which makes the following changes:
  - A. Revise the EPA registration number to be: 34704-1075.

(continued)

Signature of Approving Official:

Tony Kish  
Product Manager, Team 22  
Fungicide Branch  
Registration Division (7504P)

Date:


APR 10 2013

page 2  
EPA Reg. No. 34704-1075

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted" is enclosed for your records.

Sincerely,



Tony Kish  
Product Manager 22  
Fungicides Branch  
Registration Division (7504P)

# LOVELAND CAPTAN 80 DF

## A FUNGICIDE FOR PLANT DISEASE CONTROL

LOVELAND CAPTAN 80 DF is a water dispersible granule for use in water as a spray for the control of listed fungal diseases of fruit and as a soil treatment for the control of listed seed rots and damping-off diseases.

ACTIVE INGREDIENTS:	% BY WT.
Captan: N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide . . . . .	80.0%
OTHER INGREDIENTS: . . . . .	<u>20.0%</u>
TOTAL: . . . . .	100.0%

KEEP OUT OF REACH OF CHILDREN


# DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

CAUSES IRREVERSIBLE EYE DAMAGE

<b>FIRST AID</b>	
IF IN EYES:	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF SWALLOWED:	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
IF INHALED	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.	

EPA Reg. No. 34704-XXXX  
 EPA Est. No. 34704-MT-001  
 NET CONTENTS \_\_\_\_ GALS. ( \_\_\_\_ L)



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Do not get in eyes. Avoid contact with skin and clothing. Avoid inhalation of dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an ERA chemical resistance category selection chart. All mixers, loaders, applicators, and other handlers (including handlers participating in seeding and transplanting as part of root-dip or greenhouse-soil treatments must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear
- Chemical-resistant gloves made of any waterproof material (except pilots, flaggers or applicators driving motorized equipment) such as barrier laminate, butyl rubber, natural rubber, neoprene rubber, or nitrile rubber > 14 mils
- Chemical-resistant apron when mixing, loading, participating in dip treatments, cleaning up spills, cleaning equipment, or otherwise exposed to the concentrate.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of: 24 hours for strawberries, almonds, apples, apricots, cherries, nectarines, plums/fresh prunes, and peaches.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of: 48 hours for blueberries, grapes, raspberries, blackberries, and dewberries.

48 hours for soil treatments and root dips: For soil and greenhouse bench treatments and root dips, once the treatment and any seeding or transplanting tasks done as part of the treatment are complete, the 48-hour REI begins. Exception: Once the seeds or transplants are planted in the soil, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no contact with the soil subsurface.

**EARLY ENTRY PPE:**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Protective eyewear
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

Eye Protection: To mitigate eye irritation concerns from post-application exposure, it is required that, for at least seven days following the application of captan: 1. at least one container designed specifically for flushing eyes is available in operating condition at the WPS-required decontamination site for workers entering the area treated with captan, and 2. workers are informed orally, in a manner they can understand: -that residues in the treated area may be highly irritating to their eyes,-that they should take precautions such as refraining from rubbing their eyes to keep the residues out of their eyes, -that if they do get residues in their eyes, they should immediately flush their eyes with the eyeflush container that is located at the decontamination site, and -how to operate the eyeflush container.

**DOUBLE NOTIFICATION:**

Notify workers of the application by warning them orally or by posting warning signs at entrances to treated areas.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restriction: Do not allow people or pets to enter treated areas until sprays have dried.

For post-application fruit dips: Do not contact or allow others to contact the treated fruit until the treatment solution on the fruit has dried.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label.

In order that pesticide residues on food and forage crops will not exceed federal tolerances, use only at labeled rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to adjoining food, fiber, or pasture crops. Drift of LOVELAND CAPTAN 80 DF onto sensitive crops (e.g., D'Anjou Pears) can cause severe phytotoxicity and crop loss.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2-10 mph at the application site.

Additional requirements for aerial applications:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications into temperature inversions.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

COMPATIBILITY AND PLANT SAFETY

LOVELAND CAPTAN 80 DF can be combined safely and effectively at recommended dosage rates with most commonly used fungicides and insecticides, with the exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur, and Bordeaux mixture will reduce the fungicidal activity of LOVELAND CAPTAN 80 DF. Do not apply LOVELAND CAPTAN 80 DF in combination with or immediately

before or closely following oil sprays. Do not allow oil sprays on adjacent crops to drift onto crops which have been or will shortly be treated with LOVELAND CAPTAN 80 DF. The time factor governing the safe interval between LOVELAND CAPTAN 80 DF and oil sprays varies due to general climatic conditions, therefore, consult local agricultural spray programs and authorities to determine the proper timing. The use of spreaders which cause excessive wetting is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of LOVELAND CAPTAN 80 DF and sulfur should not be used on crops sensitive to sulfur. Used at high rates or in drenching sprays, LOVELAND CAPTAN 80 DF may cause a necrotic spotting of tender, immature leaves of certain varieties of apples, peaches, plums, and cherries. This type of injury is most likely to occur in the early cover sprays during long periods of warm, cloudy, humid weather. To avoid the hazard of leaf spotting under such conditions, use LOVELAND CAPTAN 80 DF and other spray materials at lowest rates and avoid drenching trees.

Observe the most restrictive of the labeling limitations and precautions of all products used in mixtures.

Applications can be made by aircraft or ground power equipment (including concentrate and semi-concentrate equipment). Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wetttable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read compatibility and plant safety information).

For aerial or concentrate spray applications, apply the same amount of LOVELAND CAPTAN 80 DF per acre as would normally be applied for dilute spray applications. Apply aerial or concentrate sprays in sufficient water for coverage.

#### CHEMIGATION STATEMENT

Do not apply this product through any type of irrigation system.

#### USE PRECAUTIONS

Except as specified, begin applications before or at first sign of disease and repeat as needed to maintain control but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. Maximum application is for a crop cycle. Crop cycle is defined as prebloom through postharvest. Apply the high rate and/or spray at shorter intervals when climatic conditions most favor disease(s). Apply the low rate and/or spray at larger intervals when climatic conditions least favor disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your State Agricultural Extension Service to learn of these conditions.

**IMPORTANT:** Read label carefully. Although most of the directions on this label may be followed nationwide, a few are limited to either the eastern or western U.S. Follow those directions for your growing area where specified.

FRUIT AND NUT CROPS

ALMONDS		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Brown rot twig and blossom blight, Shothole, Scab, Leaf blight, Anthracnose (for control of anthracnose, use in a disease and resistance management program of rotational sprays with other approved materials)	2.5 to 5.6	Apply at popcorn, bloom, petal fall, post petal fall, and full cover sprays.  Apply in 20 to 300 gallons of water using ground equipment or in 5 to 20 gallons of water by air.  Use 3.75 to 5.6 pounds of product per acre when captan is used alone. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, LOVELAND CAPTAN 80 DF may be used in a tank mix at a rate of 2.5 to 3.75 lbs. product per acre.  Almond hulls may be fed to livestock.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply within 30 days of harvest.</li> <li>Do not apply more than 25 pounds LOVELAND CAPTAN 80 DF per acre per crop cycle.</li> <li>REI is 24 hours.</li> </ul>		

APPLES: (East of the Rockies)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Primary scab, Black rot (frog-eye), Botrytis blossom-end-rot	5.0	Apply in 20 to 400 gallons of water using ground equipment or in 5 to 20 gallons of water by air.  Apply at 5 to 7 day intervals as needed to maintain control in prebloom, bloom, petal fall, and first cover sprays.
Secondary scab, Brooks fruit spot, Sooty blotch, Fly speck, Black rot, Black pox, Botryosphaeria rot, Bitter rot	2.5 to 5.0	Apply in 20 to 400 gallons of water using ground equipment or in 5 to 20 gallons of water by air.  Apply at 10 to 14 day intervals in second and later cover sprays.
Powdery mildew		If powdery mildew is a problem, add 6 to 12 pounds sulfur per acre to all postbloom sprays until foliage matures. NOTE: Do not use LOVELAND CAPTAN 80 DF in combination with or closely following or in alternation with wettable sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Staymen, Baldwin, King etc. as severe injury and defoliation may occur.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 40 pounds of LOVELAND CAPTAN 80 DF per acre per crop cycle.</li> <li>Do not apply more than 5lbs per acre per application.</li> <li>May be applied up to day of harvest.</li> <li>REI is 24 hours</li> </ul>		



APPLES: (West of the Rockies)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Primary scab	2.5 to 5.0	Apply in 20 to 400 gallons of water per acre using ground equipment or in 5 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, the lower rate may be used in tank mixtures.
Secondary scab		In midsummer cover sprays, the dosage may be reduced to 2.5 pounds per acre.
(Pacific Northwest): Bull's eye rot, Botrytis rot	3.75	Apply in 20 to 400 gallons of water using ground equipment or in 5 to 20 gallons of water by air. Make 1 or 2 applications with late cover sprays and 1 final spray prior to harvest.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 40 pounds of LOVELAND CAPTAN 80 DF per acre per crop cycle.</li> <li>Do not apply more than 5lbs per acre per application.</li> <li>May be applied up to day of harvest.</li> <li>REI is 24 hours.</li> </ul>		

APRICOTS		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Brown rot (twig blight), Jacket rot	1.87 to 3.12	Apply in 20 to 250 gallons of water using ground equipment or in 10 to 20 gallons of water by air.  Apply in red bud, bloom, and 75% petal fall sprays.  To reduce the potential for disease resistance development to other fungicides having a similar spectrum, use the lower rate in tank mixtures.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 15.6 pounds LOVELAND CAPTAN 80 DF per acre per crop cycle.</li> <li>May be applied up to day of harvest.</li> <li>REI is 24 hours.</li> </ul>		

BLUEBERRIES: (East of the Rockies)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Botrytis gray mold or berry rot, Mummy berry	3.12	Apply in sufficient water for thorough coverage or a minimum of 5 gallons of water by air.  Start spray program when buds swell or when buds have loose scales. Repeat at 7-day intervals through blossom period.  Repeat at 7- to 10-day intervals from late bloom.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 43.75 pounds LOVELAND CAPTAN 80 DF per acre per crop cycle.</li> <li>May be applied up to day of harvest.</li> <li>The REI is 48 hours.</li> </ul>		

BLUEBERRIES (West of the Rockies)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Botrytis gray mold or berry rot, Mummy berry	1.25 to 3.12	Apply in 20 to 200 gallons of water by ground or in 5 to 20 gallons of water by air.  Begin at mid-bloom, repeat at 7- to 10-day intervals until maturity.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 43.75 pounds LOVELAND CAPTAN 80 DF per acre per crop cycle.</li> <li>May be applied up to day of harvest.</li> <li>The REI is 48 hours.</li> </ul>		

BLACKBERRIES, RASPBERRIES, DEWBERRIES (Not registered for use in California)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Anthracoese, Botrytis, Spur blight	2.5	Apply 2.5 pounds LOVELAND CAPTAN 80 DF per acre when blossoms are in bud (young canes are 8-10 inches long). Make a second application two weeks later. Apply a fall spray after old canes are removed.
Fruit rot	2.5	Apply at early bloom (5-10% bloom) and again at full bloom. Additional applications can be made at 10-14 day intervals as needed.  Apply LOVELAND CAPTAN 80 DF as indicated above in 45-100 gallons of water per acre. Use the higher volume as foliage increases.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 12.5 pounds of LOVELAND CAPTAN 80 DF per acre per season.</li> <li>Do not apply within 3 days of harvest.</li> <li>The REI is 48 hours.</li> </ul>		

CHERRIES: (East of the Rockies)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Brown rot, Leaf spot, Botrytis rot	2.5	Apply in 20 to 200 gallons of water using ground equipment or in 10 to 20 gallons of water by air.  Apply in pre-bloom, bloom, petal fall, shuck, cover and pre-harvest sprays. Applications at 3- to 4-day intervals may be necessary during bloom to control blossom blight.  Repeat applications at 7- to 20-day intervals as needed to maintain control up to start of harvest.
Powdery mildew		If powdery mildew is a problem, add 6 lbs. sulfur per acre to the petal fall, shuck, and early cover sprays.  If sulfur is added, LOVELAND CAPTAN 80 DF may be reduced to 1.25 pounds per acre in these sprays.
Leaf spot (Post harvest sprays)		Apply in 20 to 200 gallons of water using ground equipment.  Apply immediately after harvest and repeat application in 10 to 14 days.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 17.5 pounds per acre per crop cycle.</li> <li>May be applied up to day of harvest.</li> <li>The REI is 24 hours.</li> </ul>		

CHERRIES: (West of the Rockies)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Brown rot blossom blight, Brown rot (fruit), Leaf spot	1.87 to 2.5	Apply in 20 to 200 gallons of water using ground equipment or in 10 to 20 gallons of water by air.  Apply in pre-bloom, bloom, petal fall, shuck, cover, and pre-harvest sprays.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 17.5 pounds per acre per crop cycle.</li> <li>May be applied up to day of harvest.</li> <li>The REI is 24 hours.</li> </ul>		

GRAPES: (U.S. except CA)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Phomopsis cane and leaf spot, Downy mildew, Suppression of black rot	1.25 to 2.5	Apply in 20 to 200 gallons of water using ground equipment or in 7 to 20 gallons water by air when shoots are 1/2 to 1 1/2 inches long, when shoots are 3 to 5 inches long, and when shoots are 9 to 12 inches long.  Repeat just before bloom, immediately after bloom, and continue at 10- to 14-day intervals as long as disease conditions persist. Use the lower rate when spraying less susceptible grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible grape varieties and during periods of weather highly favorable for disease development.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>• Do not apply more than 15.0 pounds of LOVELAND CAPTAN 80 DF per acre per crop cycle.</li> <li>• May be applied up to day of harvest.</li> <li>• The REI is 48 hours.</li> </ul>		

GRAPES: (California only)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Bunch rot (Botrytis)	2.5	Apply in 20 to 200 gallons of water using ground equipment or in 7 to 20 gallons of water by air.  Make 2 applications before bloom and 1 immediately after bloom. Repeat periodically, making 3 cover applications before the bunches close.
Phomopsis cane and leaf spot (current season infection)	2.0 - 2.5	Apply in 20 to 200 gallons of water using ground equipment or apply 2.5 pounds LOVELAND CAPTAN 80 DF per acre in 7 to 20 gallons of water by air.  Apply first spray when green tissue begins to show but before shoots are 1 inch long and repeat application when shoots are 6 to 8 inches long.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>• Do not apply more than 15.0 pounds of LOVELAND CAPTAN 80 DF per acre per crop cycle.</li> <li>• May be applied up to day of harvest.</li> <li>• The REI is 48 hours.</li> </ul>		

NECTARINES		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Brown rot, Scab	2.5 to 5.0	<p>Apply in 20 to 250 gallons of water using ground equipment or in 10 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, use the lower rates of LOVELAND CAPTAN 80 DF in tank mixes.</p> <p>Apply in full pink, bloom, petal fall, shuck, cover, and preharvest sprays. Applications at 3- to 4-day intervals may be necessary during bloom to control blossom blight.</p> <p>Repeat application at 7- to 14-day intervals as needed to maintain control. Continue applications throughout harvest if conditions favor brown rot.</p>
Coryneum blight (peach blight, shothole)	2.5 to 5.0	<p>Apply in 20 to 250 gallons of water using ground equipment or in 10 to 20 gallons of water by air.</p> <p>Apply in pink bud, full bloom, petal fall, and cover sprays as necessary, and as a postharvest spray (but before leaves drop).</p>
Powdery mildew	2.5 to 5.0	<p>If powdery mildew is a problem, add 7 1/2 lbs. sulfur per acre to the petal fall, shuck, and early cover spray.</p> <p>If sulfur is added, LOVELAND CAPTAN 80 DF may be reduced to 1.6 pounds per acre in these sprays.</p>
<p><b>RESTRICTIONS:</b></p> <ul style="list-style-type: none"> <li>• Do not apply more than 30 pounds per crop cycle (including postharvest sprays).</li> <li>• Pre-harvest sprays may be applied up to day of harvest.</li> <li>• The REI is 24 hours.</li> </ul>		

PEACHES		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Brown rot, Scab	2.5 to 5.0	Apply in 20 to 400 gallons of water using ground equipment or in 10 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, use the lower rates of LOVELAND CAPTAN 80 DF in tank mixes.  Apply in full pink, bloom, petal fall, shuck stages, and in cover and pre-harvest sprays.  When conditions are favorable, make applications at 3- to 4-day intervals during bloom to control blossom blight. Then repeat application at 7- to 14-day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot.
Coryneum blight (peach blight, shothole)	5.0	Apply in 20 to 400 gallons of water using ground equipment or in 10 to 20 gallons of water by air.  Apply in pink bud, full bloom, petal fall stages, and cover sprays as necessary and as a post harvest spray (but before leaves drop).
Powdery mildew	2.5 to 5.0	If powdery mildew is a problem, add 12 pounds sulfur per acre to the petal fall, shuck, and early cover spray.  If sulfur is added, LOVELAND CAPTAN 80 DF may be reduced to 2.5 pounds per acre in these sprays.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 40 pounds per acre per crop cycle (including postharvest sprays).</li> <li>Pre-harvest sprays may be applied up to day of harvest.</li> <li>The REI is 24 hours.</li> </ul>		

PLUMS, FRESH PRUNES (East of the Rockies)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Brown rot	3.75	Apply in 20 to 300 gallons of water using ground equipment or in 10 to 20 gallons of water by air.  Apply in full pink, bloom, and petal fall sprays. Repeat applications at 7- to 14-day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot.  The addition of a neutral spreader has improved coverage.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 33.75 pounds per acre per crop cycle.</li> <li>May be applied up to day of harvest.</li> <li>The REI is 24 hours.</li> </ul>		

PLUMS, FRESH PRUNES (West of the Rockies)		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Brown rot	2.5 to 3.75	Apply in 20 to 300 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Use lower rates when tank mixes with fungicides of similar spectrum of activity are used.  Apply at green bud, popcorn, bloom, and petal fall stages. Repeat in cover sprays as conditions warrant.
Prune russet scab (lacy scab)	2.5 to 3.75	Apply in 20 to 300 gallons of water using ground equipment.  Apply at full bloom.
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 33.75 pounds per acre per crop cycle.</li> <li>May be applied up to day of harvest.</li> <li>The REI is 24 hours.</li> </ul>		

STRAWBERRIES		
DISEASE	APPLICATION RATE (lbs Product/Acre)	DIRECTIONS
Botrytis (gray mold), Leaf spot	1.87 to 3.75	Apply by broadcast spray sufficient water for thorough coverage by ground equipment or in 10 to 20 gallons of water by air.  Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at 7- to 14-day intervals.  Under conditions favorable to fruit rot, continue applications through harvest period treating immediately after each picking.
If applying as a directed/banded spray; use band rate of LOVELAND CAPTAN 80 DF according to the following formula:  $\frac{\text{Plant Bed Width (inches)}}{\text{Row Spacing (inches)}} \times \text{Broadcast rate per acre} = \text{Banded rate of LOVELAND CAPTAN 80 DF per acre}$		
<b>RESTRICTIONS:</b> <ul style="list-style-type: none"> <li>Do not apply more than 30 lbs. per acre per year.</li> <li>May be applied up to day of harvest.</li> <li>The REI is 24 hours.</li> </ul>		

PEACH PREPLANT ROOT DIP (California)		
DISEASE	APPLICATION RATE	DIRECTIONS
Preventative pre-plant dip treatment for crown gall	2.5 lbs product	Use LOVELAND CAPTAN 80 DF plus 3.2 pints sodium hypochlorite (5.25% household bleach) per 100 gallons of water.  Wash nursery trees to remove soil from roots. Cut off all dormant buds and suckers in crown area and prune root system if necessary. Submerge the entire dormant tree for 5 minutes.  Recharge dip during operation at a rate of 3.2 pints diluted sodium hypochlorite per 100 gallons of water.

POSTHARVEST FRUIT APPLICATION

(For use in mechanical fruit dip operations only. Hand dipping of fruit is prohibited.)

For control of various molds and storage rots (Botrytis, Gleosporium, Rhizopus).

Use as a post harvest dip or spray wash on the following fruits:

Apples, Cherries, Pears; Use 1.6 pounds LOVELAND CAPTAN 80 DF per 100 gallons of water. Apply as a spray or in a dip tank. When used as a dip, recharge wash solution periodically when tank volume is reduced by 25%. Bring water back to volume and add 1.6 pounds LOVELAND CAPTAN 80 DF for each 100 gallons added. At end of every 8-hour shift, empty tank, flush, and charge with fresh dilution. Do not allow tank solution to stand overnight. Maintain continuous agitation during dipping operation.

For post-application fruit dips:

Do not contact or allow others to contact the treated fruit until sprays have dried.

DISPOSAL OF LEFTOVER POSTHARVEST TREATMENT MIXTURE:

Leftover dip or spray mixtures containing Captan may be used as a foliar spray for the same crop in case of apples and cherries (but not pears) as treated by the dip or spray mixture, observing all restrictions such as maximum pounds applied per application and season.

When calculating application rates, if analytical services are not available to determine the exact quantity of Captan remaining in the mixture, assume that the tank still contains 1 3/5 pounds of LOVELAND CAPTAN 80 DF per 100 gallons of water. If the dip or spray mixture contains other pesticides in addition to LOVELAND CAPTAN 80 DF, refer to the product label(s) for information regarding disposal. LOVELAND CAPTAN 80 DF wastes are acutely hazardous to the eyes. Improper disposal of spray or dip tank mixtures is a violation of Federal Law. If the leftover dip or spray mixture cannot be disposed of in the manner prescribed above, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance concerning the disposal of spent or excess dip tank mixtures.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Do not store near oxidizing agents.

**PESTICIDE STORAGE:** Store only in original container in locked storage areas. If a spill occurs, visible puddles or drops should be covered with a dry absorbent such as clay, sawdust, lime or soda ash. Sweep up and dispose of in a permitted disposal facility. Avoid cross-contamination with other pesticides and fertilizers during handling and storage. Do not store near heat or open flame. Protect from temperatures below 0 degrees F. Not for use or storage in or around the home.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL: Nonrefillable container.** Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at [www.acrecycle.org](http://www.acrecycle.org). If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**For packages up to 5 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into



application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 56 gallons:** To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**For refillable containers:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container.

Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Offer for recycling or reconditioning, if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**BEFORE BUYING OR USING THIS PRODUCT,** read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. To the extent consistent with applicable law the buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

FORMULATED FOR:  
LOVELAND PRODUCTS, INC.  
P.O.BOX 1286  
GREELEY, CO 80632