

34704-1086

2/7/2014

1/19

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460



OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

FEB - 7 2014

Premjit Halarnkar  
Loveland Products, Inc.  
P.O. Box 1286  
Greeley, CO 80632-1286

Subject: Amended label to reinstate inadvertently deleted directions for tobacco  
Product Name: Match-Up Insecticide  
EPA Registration No. 34704-1086  
EPA Decision No. 487571  
Your submission dated February 6, 2014

Dear Dr. Halarnkar:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable. A stamped copy of the label is enclosed for your records. Please submit one copy of your final printed labeling before you release the product for shipment. Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions, please contact Julie Chao at (703) 308-8735 or [chao.julie@epa.gov](mailto:chao.julie@epa.gov).

Regards,

A handwritten signature in black ink that reads "Venus Eagle".

Venus Eagle, Product Manager 01  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

Enclosure

2/19



**RESTRICTED USE PESTICIDE**  
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**GROUP 1B 3 INSECTICIDE**

# Match-Up<sup>TM</sup>

## INSECTICIDE

For control of listed insects infesting certain field and vegetable crops.

<b>ACTIVE INGREDIENT:</b>	
Chlorpyrifos: O,O-diethyl-O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate .....	28.6%
Bifenthrin* .....	9.0%
<b>OTHER INGREDIENTS</b> .....	62.4%
<b>TOTAL</b>	<b>100.0%</b>

Contains 2.47 pounds of chlorpyrifos per gallon and 0.78 pound of bifenthrin per gallon.  
\*Cis isomers 97% minimum, trans isomers 3% maximum.  
This product contains petroleum distillates.

### KEEP OUT OF REACH OF CHILDREN

### CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

<b>FIRST AID</b> Organophosphate	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.  
**FOR MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL: 1-866-944-8565**  
**NOTE TO PHYSICIAN:** Bifenthrin is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.  
**NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration.  
**NOTE TO PHYSICIAN:** Contains petroleum distillate - vomiting may cause aspiration pneumonia.

ACCEPTED  
FEB - 7 2014

EPA REG. NO. 34704-1086  
EPA EST. NO. 34704-MS-001  
NET CONTENTS 2.5 GAL (9.46 L)

Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended, for the  
pesticide registered under:

EXP 02/14 Tobacco

EPA. Reg. No: 34704-1086

8/19

**MATCH-UP™ INSECTICIDE**  
**EPA REG. NO. 34704-1086**

**PRECAUTIONARY STATEMENTS**  
**HAZARD TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

**CAUTION.** Harmful if swallowed. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Causes moderate eye irritation.

**Personal Protective Equipment (PPE)**

Some materials that are chemical-resistant to this product are barrier laminate and viton.

**Mixers and loaders using a mechanical transfer loading system and applicators using aerial application equipment must wear:**

- Long-sleeved shirt and long pants,
- Shoes and socks.

**In addition to the above, mixers and loaders using a mechanical transfer loading system must wear:**

- Chemical-resistant gloves,
- Chemical-resistant apron,
- A NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any R, P, or HE filter.

See Engineering Controls for additional requirements.

**All other mixers, loaders, applicators and handlers must wear:**

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves,
- Chemical-resistant apron when mixing or loading or exposed to the concentrate,
- Chemical-resistant footwear plus socks,
- Chemical-resistant headgear for overhead exposure,
- A NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any R, P, or HE filter.

**USER SAFETY REQUIREMENTS**

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls**

Mixers and loaders supporting aerial applications must use a mechanical transfer system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)] for dermal protection, and must: Wear the personal protective equipment required above for mixers/loaders, wear protective eyewear if the system operates under pressure, and be provided and have immediately available for use in an emergency, such as broken package, spill, or equipment breakdown: coveralls, chemical-resistant footwear and chemical-resistant headgear if overhead exposure.

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

Use of human flaggers is prohibited. Mechanical flagging equipment must be used.

When handlers use closed cab motorized ground application equipment in a manner that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

4/19

**MATCH-UP™ INSECTICIDE  
EPA REG. NO. 34704-1086**

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish, aquatic invertebrates, small mammals and birds. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply this product or allow it to drift to blooming crops or weeds if bees are actively visiting the treatment area.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging in or adjacent to the treatment area.

The use of bifenthrin is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county, contact the local extension service for procedures and precautions to use to protect endangered species.

**PHYSICAL OR CHEMICAL HAZARDS**

Combustible - Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

**Restricted Use Pesticide**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

This product cannot be reformulated or repackaged into other end-use products.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Match-Up™ Insecticide is an emulsifiable concentrate for use in listed crops. Target pests and application rates are provided in the accompanying tables.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the directions for use associated with each crop.**

**Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Certified crop advisors or persons entering under their direct supervision under certain circumstances may be exempt from the early reentry requirements pursuant to 40 CFR Part 170. PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants,
- Chemical-resistant gloves made out of any waterproof material,
- Chemical-resistant footwear plus socks,
- Chemical-resistant headgear for overhead exposure.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

**USE PRECAUTIONS AND RESTRICTIONS**

Insect control may be reduced at low spray volumes under high temperature and wind conditions.

Some reduction in insect control may occur under unusually cool conditions.

**Flood irrigation:** To avoid contamination of irrigation tail waters, do not flood irrigate within 24 hours following a soil surface or foliar application of Match-Up Insecticide.

**Do not aerially apply this product in Mississippi.**

**Insecticide Resistance Management (IRM)**

Match-Up Insecticide contains a Group 1B insecticide and a Group 3 insecticide. Insect/mite biotypes with acquired resistance to Group 1B and Group 3 may eventually dominate the insect/mite population if Group 1B and Group 3 insecticides are used repeatedly in the same

5/19

**MATCH-UP™ INSECTICIDE  
EPA REG. NO. 34704-1086**

field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Match-Up Insecticide or other Group 1B and Group 3 insecticides.

To delay development of insecticide resistance, the following practices are recommended:

- Avoid consecutive use of insecticides with the same mode of action (same insecticide group) on the same insect species.
- Use tank mixtures or premix products containing insecticides with different modes of action (different insecticide groups) provided the products are registered for the intended use.
- Base insecticide use on comprehensive Integrated Pest Management (IPM) programs.
- Monitor treated insect populations in the field for loss of effectiveness.
- Contact your local extension specialist, certified crop advisor, and/or manufacturer for insecticide resistance management and/or IPM recommendations for the specific site and resistant pest problems.

**SPRAY DRIFT MANAGEMENT**

**Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, non-target crops, aquatic and wetland sites, woodlands, pastures, rangelands, or animals.**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making decision to apply this product.

Observe the following precautions when spraying Match-Up Insecticide adjacent to permanent bodies of water such as rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds.

The following treatment setbacks or buffer zones must be utilized for applications around the above listed aquatic areas with the following application equipment:

<b>Application Method</b>	<b>Required Setback (Buffer Zone) (Feet)</b>
Ground boom	25
Chemigation	25
Orchard airblast	50
Aerial (fixed wing or helicopter)	150

Making applications when wind is blowing away from sensitive areas is the most effective way to reduce the potential for adverse effects.

The following spray drift **best management practices** are recommended to avoid off-target drift movement from applications.

**Aerial Application**

1. The boom width must not exceed 75% of the wingspan or 90% of the rotor blade.
2. Nozzles must always point backward, parallel with the air stream, and never be pointed downward more than 45 degrees.
3. Nozzles must produce a medium or coarser droplet size (255-340 microns volume median diameter) per ASE Standard 572 under application conditions. Airspeed, pressure, and nozzle angle can all effect droplet size. See manufacturer's catalog or USDA/NAAA Applicator's Guide for spray size quality ratings.
4. Applications must not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.
5. Use upwind swath displacement and apply only when wind speed is 3 to 10 mph as measured by an anemometer. Do not apply product when wind speed exceeds 10 mph.
6. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above the ground or crop canopy.

Where states have more stringent regulations, they must be observed.

The applicator must be familiar with and take into account the information covered in the Aerial Drift Reduction Instructions.

**Aerial Drift Reduction Instructions**

**Information on Droplet Size:** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent adverse effects from drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

6/19

**MATCH-UP™ INSECTICIDE**  
**EPA REG. NO. 34704-1086**

**Controlling Droplet Size:**

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

**Boom Length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

**Application Height:** Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**Swath Adjustment:** When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

**Wind:** Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 1.5 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect spray drift.

**Temperature and Humidity:** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions:** Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Sensitive Areas:** The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

**Ground Boom Application**

The following mandatory spray drift best management practices are required to reduce the likelihood of off-target drift movement from ground applications.

1. Choose only nozzles and pressures that produce a medium or coarse droplet size (255 to 400 microns volume median diameter), per ASAE Standard 572. See manufacturer's catalog or USDA/NAAA Applicator's Guide for spray size quality ratings.
2. Apply with nozzle height no more than 4 feet above the ground or crop canopy.
3. Do not apply product when wind speed exceeds 10 mph as measured by an anemometer.

**Orchard Airblast Application**

The following mandatory spray drift best management practices are required to reduce the likelihood of off-target drift movement from airblast applications.

1. Nozzles must be directed so spray is not projected above the canopies.
2. Apply only when wind speed is 3 to 10 mph at the application site as measured by an anemometer outside of the orchard/vineyard on the upwind side.
3. Outward pointing nozzles must be shut off when turning corners at row ends.

*The applicator must take into account the following best management practices to reduce off-site spray drift. This section is advisory and does not supersede mandatory label requirements.*

7/19

**MATCH-UP™ INSECTICIDE  
EPA REG. NO. 34704-1086**

1. Number of nozzles, nozzle orientation and spray volume, air speed and wind direction are key factors in adjusting airblast spray delivery to match the height and density of the crop canopy. Airblast equipment should be adjusted to provide uniform coverage while minimizing the amount of spray movement over-the-top or completely through the crop canopy.
  - High air volumes deliver spray more efficiently than air at high speed. Reducing forward travel speed decreases the air speed necessary to deliver the spray to the top of the crop canopy.
  - Use air guides along with the number and orientation of spray nozzles to achieve the desired spray coverage and directional control.
2. The following steps must be taken to minimize drift and the amount of non-target spray:
  - Orient nozzles and adjust air speed/volume/direction to force the spray through the crop canopy but not allow drift past the canopy.
  - Shut off spray delivery when passing gaps in crop canopy within rows.
  - Spray the outside rows of orchards from outside in, directing the spray into the orchard and shutting off nozzles on the side of the sprayer away from the orchard.
  - When treating smaller trees, vines or bushes, shut off top nozzles to minimize over-the-top spray movement.

**BUFFER ZONES**

**Vegetative Buffer Strip**

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing Bifenthrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers:

*Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21 pp.*

<http://www.in.csusda/v/technical/agronom/newconbuf.pdf>

**Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast)**

Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

**Buffer Zone for ULV Aerial Application**

Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

**Buffer Zone for Non-ULV Aerial Application**

Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

**SPRAY DRIFT REQUIREMENTS**

**Wind Direction and Speed**

Only apply this product if the wind direction favors on-target deposition.

Do not apply when the wind velocity exceeds 15 mph.

**Temperature Inversion**

Do not make aerial or ground applications into temperature inversions.

Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

**Droplet Size**

Use only medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

**Additional Requirements for Ground Applications**

Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

**Additional Requirements for Aerial Applications**

The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

8/19

**MATCH-UP™ INSECTICIDE  
EPA REG. NO. 34704-1086**

**Buffer Zones**

The buffer distances specified in the below table are the distances in feet that must exist to separate sensitive sites from the targeted application site. Buffers are measured from the edge of the sensitive site to the edge of the application site. Sensitive sites are areas frequented by non-occupational bystanders (especially children). These include residential lawns, pedestrian sidewalks, outdoor recreational areas such as school grounds, athletic fields, parks and all property associated with buildings occupied by humans for residential or commercial purposes. Sensitive sites include homes, farm worker housing, or other residential buildings, schools, daycare centers, nursing homes, and hospitals. Non-residential agricultural buildings, including barns, livestock facilities, sheds, and outhouses are not included in this prohibition.

Application Rate (Lb AI/A)	Nozzle Droplet Type	Required Setback (Buffer Zones) (Feet)		
		Aerial	Airblast	Ground
>0.5 to 1.0	coarse or very coarse	10	10	10
>0.5 to 1.0	medium	25	10	10
>1.0 to 2.0	coarse or very coarse	50	10	10
>1.0 to 2.0	medium	80	10	10
>2.0 to 3.0	coarse or very coarse	80 <sup>1</sup>	10	10
>2.0 to 3.0	medium	100 <sup>1</sup>	10	10
>3.0 to 4.0	medium or coarse	NA <sup>2</sup>	25	10
>4.0	medium or coarse	NA	50	10

<sup>1</sup> Aerial application of greater than 2.0 pounds active ingredient per acre is only permitted for Asian Citrus Psylla control, up to 2.3 pounds active ingredient per acre.

<sup>2</sup> NA is not allowed.

Only pesticide handlers are permitted in the setback area during application of this product. Do not apply this product if anyone other than a mixer, loader, or applicator, is in the setback area. Exception: Vehicles and persons riding bicycles that are passing through the setback area on public or private roadways are permitted.

**APPLICATION INSTRUCTIONS**

**Broadcast Foliar Application**

Apply with conventional power-operated spray equipment using nozzles and spray pressures specified for insecticides. Apply Match-Up Insecticide in a spray volume of not less than 2.0 gallons per acre for aerial application equipment (fixed wing or helicopter) or not less than 10.0 gallons per acre for ground equipment, unless otherwise specified. Increase spray volume to ensure adequate coverage with increased density and height of crop canopy. See Spray Drift Management section for specifications on droplet size.

**Ground Application:** Orient the boom and nozzles so that uniform coverage is obtained. The swath width should not be wider than the boom. Follow nozzle manufacturer's specifications for insecticide nozzles with respect to nozzle type, pressure, and spacing.

**Broadcast Soil Application**

Apply with conventional power-operated spray equipment that will apply the product uniformly to the soil surface. Use nozzles that produce medium or coarse droplets (235 to 400 microns). Unless otherwise indicated, a spray volume of 10.0 gallons or more per acre is recommended. For band application, use proportionally less spray volume.

**Aerial Application**

Use a minimum spray volume of 2.0 gallons per acre and follow the **best management practices** for aerial application, above. Marking of swaths by flagging, permanent markers or use of GPS equipment is recommended.

**Chemigation (Sprinkler Irrigation)**

Match-Up Insecticide may be applied to the following crops through sprinkler irrigation equipment: citrus (orchard floors only), corn (field and sweet), cotton and soybeans. Do not apply this product by chemigation unless specified in crop-specific directions in this label or state-specific 24(c) supplemental labeling. Do not apply to labeled crops through any other type of irrigation system.

**Note:** Unless otherwise indicated in specific use directions, the application rates for chemigation are the same as those specified for broadcast application.

**Use Directions for Sprinkler Irrigation**

The following use directions must be followed when Match-Up Insecticide is applied through sprinkler irrigation systems. Thoroughly clean the injection system and tank of any fertilizer or chemical residues, and dispose of the residues according to state and federal laws. Flush the injector with soap and water. Determine the amount of Match-Up Insecticide needed to cover the desired acreage. Mix according to instructions in the Mixing Directions section and bring mixture to desired volume. Do not add crop oil when Match-Up Insecticide is applied by chemigation. Maintain continuous agitation during mixing and throughout the application period. Set the sprinkler system to deliver the desired inches of water per acre. Start the water pump and sprinkler, and let the system achieve the desired pressure and



9/19

**MATCH-UP™ INSECTICIDE  
EPA REG. NO. 34704-1086**

speed before starting the injector. Start the injector and calibrate the injector system according to Calibration Instructions in the following Special Use Precautions section. The mixture containing Match-Up Insecticide must be injected continuously and uniformly into the irrigation water line as the sprinkler is moving to ensure uniform application at the correct rate. When the application is finished, flush and clean the entire irrigation and injector system prior to shutting down the system.

**Use Precautions for Sprinkler Irrigation**

The following use precautions will result in a safe and successful application of mixtures containing Match-Up Insecticide:

1. Apply this product only through the following sprinkler irrigation systems: center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, micro sprinkler, or hand move. Do not apply this product through any other type of irrigation system. Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units.
2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non uniform distribution of treated water.
3. If you have questions about calibration, you should contact state extension service specialists, equipment manufacturers, or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
6. The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Refer to the American Society of Agricultural Engineer's Engineering Practice 409 for more information.
7. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
10. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. The metering pump must provide a greater pressure than that of the irrigation system at the point of injection. The pump must meet Section 675 for "Electrically Driven or Controlled Irrigation Machines" NEC 70 and must contain viton or teflon seals.
12. To insure uniform mixing of the insecticide into the water line, inject the mixture through a nozzle placed in the fertilizer injection port or just ahead of an elbow or tee in the irrigation line so that the turbulence will assist in mixing. It is suggested that the injection point be higher than the insecticide tank to prevent siphoning.
13. The tank holding the insecticide mixture should be large enough to allow the system to complete the application with 1 filling. It must be free of rust, fertilizer, sediment, and foreign material, and equipped with an in-line strainer situated between the tank and the injector pump.
14. **Calibration:** In order to calibrate the irrigation system and injector to apply the mixture of Match-Up Insecticide, determine the following: 1) Calculate the number of acres irrigated by the system; 2) Set the irrigation rate and determine the number of minutes for the system to cover the intended treatment area; 3) Calculate the total gallons of insecticide mixture needed to cover the desired acreage. Divide the total gallons of insecticide mixture needed by the number of minutes to cover the treatment area. This value equals the gallons per minute output that the injector must deliver. Convert the gallons per minute to milliliters or ounces per minute. Calibrate the injector pump with the system in operation at the desired irrigation rate. It is suggested that the timed output of the injector pump be checked at least twice before operation, and the system monitored during operation.
15. Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application if they irrigate non-target areas.
16. Do not allow irrigation water to collect or run off and pose a hazard to livestock, wells, or adjoining crops.
17. Reentry: Follow requirements in the Agricultural Use Requirements section or crop-specific sections of this label.
18. Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units.

**Mixing Directions**

To prepare the spray, add a portion of the required amount of water to the spray tank and with the spray tank agitator operating add Match-Up Insecticide. Complete filling the tank with the balance of water needed. Maintain sufficient agitation during both mixing and application to ensure uniformity of the spray mixture.

Match-Up Insecticide is compatible with insecticides, miticides, and fungicides and non-ressure fertilizer solutions commonly recommended except for alkaline materials such as Bordeaux mixture and lime. It is always recommended that a small jar compatibility test be run prior to tank mixing. Prepare tank mixtures in the same manner as recommended above for use of Match-Up Insecticide alone. When tank mixing Match-Up Insecticide with herbicides, add wettable powders first, flowables second, and emulsifiable concentrates last. When a fertilizer solution is involved, it is strongly recommended that a fertilizer pesticide compatibility agent such as E-Z Mix be used.

10/19

**MATCH-UP™ INSECTICIDE**  
**EPA REG. NO. 34704-1086**

Maintain constant agitation during both mixing and application to ensure uniformity of the spray mixture. Do not allow spray mixtures to stand overnight.

**Tank Mix Compatibility Test:** Test compatibility of the intended tank mixture before adding Match-Up Insecticide to the spray or mix tank. Add proportional amounts of each tank mix ingredient to a pint or quart jar, cap, shake, and let set for 15 minutes. Formation of precipitates that do not readily redisperse indicates an incompatible mixture that should not be used.

**CITRUS ORCHARD FLOORS**  
**(NOT FOR USE IN MISSISSIPPI)**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 5 days unless PPE required for early entry is worn.

Apply Match-Up Insecticide by ground equipment to bare soil beneath citrus trees. Remove weed growth or other obstructions that might prevent the spray from reaching the soil surface.

Match-Up Insecticide must be uniformly applied from the trunk to the drip line of tree. Apply the dosage specified in the table below in a minimum of 40.0 gallons of dilute spray per acre, using equipment that will apply the spray uniformly to the soil surface. Greater spray volume should insure greater uniformity of coverage. A pre- and post-application irrigation may aid in the uniformity of coverage as well.

Pest	Rate/A	Use Instructions
Blue-green citrus root weevil <i>(Pachnaeus opalus)</i> Brown leaf notcher <i>(Epicarus mexicanus)</i> Diaprepes root weevil <i>(Diaprepes abbreviatus)</i> Little leaf notcher <i>(Artipus floridanus)</i> Southern blue-green root weevil <i>(Pachnaeus litus)</i>	40.9 to 51.8 fl oz	<p>The use of this product protects citrus tree roots from <i>Diaprepes</i> and other Citrus root weevil feeding by creating a barrier. As Citrus root weevil eggs hatch in new foliage, neonates fall to the soil surface beneath the tree and come in contact with this product as they attempt to burrow into the root zone. Disturbance of the soil beneath trees should be minimized.</p> <p>Timing of this product's applications is critical. Peak emergence of <i>Diaprepes</i> adults varies by citrus growing region and environmental factors such as soil moisture can affect emergence. Usually, 2 peaks are observed for <i>Diaprepes</i>, first in the spring then late summer or early fall. Southern blue-green and Blue-green citrus weevils and Fuller rose beetle usually have a single emergence peak in the spring. Brown and Little leaf notches usually have 3 emergence peaks, spring, summer, and fall. Since emergence varies by region and season, the best way to time application is to observe the adults. Adults are most active early morning and late afternoon; numbers can be estimated by trapping throughout spring and summer (emergence periods). Eggs are laid 8 to 10 weeks following the adult emergence from the soil; larval invasion into the soil will begin 2 to 3 weeks following adult emergence. It is critical to have this product soil barrier in place prior to drop of the neonates.</p> <p>This product is one of several effective tools in an integrated pest management program for Citrus root weevils. Application of this product should be used in conjunction with good cultural practices, biological control of larvae and foliar control of adults. Consult local university extension personnel for current information to protect citrus trees from Citrus root weevils and other pests.</p> <p>Apply to individual citrus resets, when not in solid planted rows, using hand-gun or shielded sprayer. Peak emergence of <i>Diaprepes</i> root weevil generally occurs in the spring. Depending on weather conditions, a minor emergence of <i>Diaprepes</i> root weevil may also occur in the fall. If the citrus grove to be treated is in an area where weather conditions are conducive to primary emergence occurring in the spring, 53.3 fl oz of this product should be used to obtain the longest residual management of <i>Diaprepes</i> root weevil. If the citrus grove to be treated is in an area where weather conditions will promote more than 1 peak of pest emergence, 42.1 fl oz of this product can be applied early season and 42.1 fl oz of this product can be applied later in the season.</p>

Cont'd. next page

11/19

**MATCH-UP™ INSECTICIDE  
EPA REG. NO. 34704-1086**

**Citrus Orchard Floors cont'd.:**

**Use Restrictions:**

- **DO NOT** apply in tank mixture with Evik® herbicide.
- **DO NOT** apply through irrigation systems.
- **DO NOT** allow any application of this product to contact fruit or foliage.
- **DO NOT** allow meat or dairy animals to graze in treated areas.
- Maximum single application rate, based upon maximum single application limit of 1.0 pound active ingredient chlorpyrifos per acre, is 51.8 fluid ounces of Match-Up Insecticide.
- **DO NOT** apply more than a total of 82.1 fluid ounces of Match-Up Insecticide per acre per year, based upon a maximum limit of 0.5 pound active ingredient bifenthrin per acre per year.
- **DO NOT** make more than 3 applications of Match-Up Insecticide or other products containing chlorpyrifos per year (does not include foliar applications to citrus trees).
- **DO NOT** make a second application of Match-Up Insecticide or other product containing chlorpyrifos within 10 days of the first application.
- Foliar applications of other products containing chlorpyrifos may be made in addition to the orchard floor treatments but must comply with the 10 day re-treatment interval.
- Ground application only. Do not apply by air.
- **DO NOT** apply within 28 days of harvest.

**FIELD CORN and SWEET CORN  
(Grain, Silage and Corn Grown for Seed)**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**PREEMERGENCE INCLUDING CONSERVATION TILLAGE**

Apply as a broadcast spray to surface trash and exposed soil using power-operated ground spray equipment. Use a total spray volume of 20.0 gallons per acre or more.

Pest	Rate/A	Use Instructions
Armyworms Cutworms	6.6 fl oz	This product may also be applied in tank mixes with preemergence herbicides such as paraquat or glyphosate and/or liquid fertilizer solutions.

**Use Restrictions:**

- **For field corn - DO NOT** apply more than 0.3 pound bifenthrin active ingredient equivalent (1.5 quarts Match-Up Insecticide) per acre total per season, including preemergence, preplant incorporated, at-plant and foliar applications of other bifenthrin products.
- **For sweet corn- DO NOT** apply more than 0.2 pound bifenthrin active ingredient equivalent (1.0 quart Match-Up Insecticide) per acre total per season, including preemergence, preplant incorporated, at-plant and foliar applications of other bifenthrin products.

**AT-PLANT T-BAND APPLICATION**

Pest	Rate/A	Use Instructions
Corn rootworm larvae (northern, southern, western)	0.8 fl oz /1000 lin ft of row	Apply as a 5- to 7-inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel centered over the row. Use the table below to determine this product's needs/A. Apply in a minimum of 3.0 gal of finished spray/A. (3.0 gal/A is approximately 0.2 gal/1000 lin ft of row at 36.0 inch spacing). Incorporate into the top 1 inch of soil using press wheel, tines, chains or other suitable equipment. Mix this product with water or fertilizer in the following manner. Fill the spray tank approximately 1/2 full with water or liquid fertilizer, add the proper amount of this product, then add the rest of the water or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture. Applications of this product alone or in recommended tank mixtures, in conjunction with in furrow pop-up fertilizers may be used. A jar compatibility test should be performed with appropriate ratio of this product and fertilizer to ensure mixture will stay in solution. Constant agitation should be maintained during mixing and application.
Army cutworm Grubs Other cutworm spp. Seed corn beetle Seed corn maggot True armyworm or Armyworm spp. Wireworm	0.4 to 0.8 fl oz/1000 lin ft of row	

12/19

**MATCH-UP™ INSECTICIDE**  
**EPA REG. NO. 34704-1086**

**Field Corn and Sweet Corn cont'd..Use Restrictions:**

- **DO NOT** apply to soil where there is greater than 30% cover of crop residue remaining.
- **DO NOT** apply within 30 days of harvest of grain or ears.
- **DO NOT** apply in tank mixes with Steadfast® or Lightning® herbicides.
- **DO NOT** graze livestock in treated area or cut treated crops for feed within 30 days of treatment.
- **DO NOT** apply more than 0.1 pound active ingredient bifenthrin (16.41 fluid ounces Match-Up Insecticide) per acre per season as an at-plant application.

To calculate the amount of Match-Up Insecticide to use per acre based on row spacing, follow the conversion chart below.

<b>Row spacing (inches)</b>	<b>40</b>	<b>38</b>	<b>36</b>	<b>30</b>
Match-Up Insecticide (lb AI chlorpyrifos/bifenthrin per acre)	0.196/0.06	0.207/0.065	0.218/0.069	0.262/0.08
Match-Up Insecticide (fl oz/A)	10.2	10.7	11.3	13.6

**POSTEMERGENCE**

Apply as a postemergence broadcast spray using sufficient spray volume to ensure thorough coverage of treated plants, but no less than 15.0 gallons per acre for ground spray equipment or 2.0 to 5.0 gallons per acre for aircraft equipment. Control may be reduced at low spray volumes under high temperature and wind conditions.

To improve control by aircraft, use 5.0 gallons of finished spray per acre particularly when initial populations are heavier than normal. Thorough coverage is essential to achieve control.

**Chemigation:** This product may be applied through sprinkler irrigation systems at instructed broadcast application rates to control listed foliar pests. See Chemigation section for application instructions.

This product may be tank mixed with glyphosate products when application is to be made to glyphosate-tolerant corn.

Refer to table below for rates of application and for spray volumes for control of specific pests.

**Note:** Do not aerially apply in Mississippi.

*Cont'd. next page*

13/19

**MATCH-UP™ INSECTICIDE**  
**EPA REG. NO. 34704-1086**

**Field Corn and Sweet Corn cont'd.:**

Pest	Rate/A	Use Instructions
Aphids Army cutworm Beet armyworm Billbugs (1) Cereal leaf beetle Chinch bug Common stalk borer (6) Corn earworm Corn rootworm adults Cucumber beetle adults Cutworm species (2) European corn borer (4) Fall armyworm Flea beetle Grasshoppers Greenbug Japanese beetle adult Lesser corn stalk borer Sap beetle Southern armyworm Southern corn leaf beetle Southwestern corn borer (5) Stinkbugs Tarnished plant bug True armyworm or armyworm spp Webworms (3) Western bean cutworm Yellowstriped armyworm	5.5 to 16.4 fl oz	<b>Pest-Specific Instructions:</b> <b>1. For best Billbug control</b> , ground apply in a minimum spray volume of 20.0 to 40.0 gal/A at 40 psi. If corn is less than 6 inches tall, apply in a 9- to 12-inch wide band over the row. For corn greater than 6 inches tall, apply using drop nozzles directed to the base of the plant. Do not reduce the application rate for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone. <b>2. For Cutworms</b> , it is preferable to apply this product when soil is moist and worms are active on or near the soil surface. If ground is dry, cloddy, or crusted at time of treatment, worms may be protected from the spray and effectiveness will be reduced. Shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment may improve control. A second application may be required if damage or density levels exceed economic thresholds established for your area. <b>3. For Webworm control</b> , shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment is necessary. <b>4.</b> University research indicates that achieving greater than 50% control of first-generation <b>European borer</b> with a single liquid insecticide treatment is highly dependent upon timing, insecticide placement and weather conditions. Make application for <b>Corn borer</b> control with initial application at or shortly before egg hatch. <b>5. For Southwestern corn borer</b> , make application for Corn borer control with initial application at or shortly before egg hatch. A second application may be applied 21 days later if needed due to reinfestation. <b>6. Do not</b> use this product in combination with a burndown herbicide for control of Common stalk borer. For Common stalk borer control, treat approximately 11 days after application of glyphosate or after burndown with paraquat herbicide is complete (3 to 5 days). <b>For control of other ear-attacking pests:</b> Apply this product just before silking and repeat as necessary to maintain control. <b>For control of other insect pests:</b> Apply when pests first appear and repeat as necessary.
Banks grass mite (1) Carmine mite (2) Twospotted spider mite (2)	13.2 to 16.4 fl oz	<b>Pest-Specific Instructions:</b> <b>1. Apply for Banks grass mite control</b> when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant. <b>2. For Twospotted spider mite and Carmine mite control:</b> Apply when colonies first form prior to leaf damage or discoloration and before wide-spread Mite dispersal throughout the canopy.  Higher rates within the listed rate range will be necessary for heavier initial populations and corn under heat or drought stress. Field experience with dimethoate at 0.5 lb A/A in tank mixture has demonstrated good control under these conditions. For Mite control in Texas, New Mexico, Oklahoma, and Arizona, apply in a minimum of 5.0 gal of finished spray/A by aircraft or in a minimum of 10.0 gal/A with ground equipment.

Numbers in parentheses (-) refer to Pest-Specific Instructions.

**Use Restrictions:**

- For field corn, do not apply more than 49.2 fluid ounces of Match-Up Insecticide per acre per season, based upon a maximum application limit of 0.3 pound active ingredient bifenthrin per acre per season.
- For sweet corn, do not apply more than 32.8 fluid ounces of Match-Up Insecticide per acre per season, based upon a maximum application limit of 0.2 pound active ingredient bifenthrin per acre per season.
- **DO NOT** make more than 3 applications of any product containing chlorpyrifos per season, including the maximum allowed of 2 granular applications, at the 1.0 pound active ingredient chlorpyrifos rate.

Cont'd. next page

09/19

**MATCH-UP™ INSECTICIDE  
EPA REG. NO. 34704-1086**

**Field Corn and Sweet Corn cont'd.:**

- **DO NOT** make a second application of Match-Up Insecticide or other product containing chlorpyrifos within 10 days of the first application.
- **DO NOT** graze livestock in treated areas or cut treated crops for feed within 30 days of the last application.
- Use of ultra low volume (ULV) application on corn is prohibited.
- **DO NOT** make aerial or ground applications to corn if heavy rainfall is imminent.
- Use of this product on corn is prohibited in all coastal counties.
- **DO NOT** apply in tank mixes with Steadfast or Lightning herbicides.
- **DO NOT** apply within 30 days of harvest of field corn or 21 days of harvest of sweet corn.

**COTTON  
(NOT FOR USE IN MISSISSIPPI)**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Apply as a broadcast foliar spray using aircraft or ground spray equipment. Use a higher rate in the rate range when there is increased pest pressure. Use sufficient spray volume to ensure thorough coverage of treated plants, but no less than 10.0 gallons per acre for ground spray equipment or 2.0 gallons per acre for aircraft equipment. Increase spray volume when foliage is dense and/or pest population is high and/or under high temperature and wind conditions. Treat when field counts indicate damaging insect populations are developing or present.

**Chemigation:** This product may be applied through sprinkler irrigation systems at instructed broadcast application rates to control listed foliar pests. See Chemigation section for application instructions.

Proper application methods are necessary to ensure thorough spray coverage and correct rate, and minimize off-target drift. Follow Application Directions for ground and aerial application and Spray Drift Management instructions of this label.

<b>Pest</b>	<b>Rate/A</b>	<b>Use Instructions</b>
European corn borer Soybean (banded) thrips Tobacco thrips	3.4 to 16.4 fl oz	<b>Pest Specific Instructions:</b> <b>To control Boll weevil:</b> Apply this product at an interval of 3 to 4 days until pest numbers are reduced to acceptable levels. <b>To control Mites and Aphids:</b> Apply when pests first appear. Repeat as necessary to maintain control. Higher rates within the listed rate range will be required once a damaging threshold is established.
Boll weevil Cabbage looper Cotton aphid Cotton fleahopper Cotton leaf perforator Cutworms Fall armyworm Grasshoppers Leafhopper Plant bugs Saltmarsh caterpillar Southern garden leafhopper Stink bugs Tobacco budworm Whitefly Yellow striped armyworm	6.7 to 16.4 fl oz	
Beet armyworm Carmine spider mite Cotton bollworm Kudzu bug <i>Lygus</i> spp. Pink bollworm Twospotted spider mite	9.75 to 16.4 fl oz	

**Use Restrictions:**

- **DO NOT** make more than 3 applications of Match-Up Insecticide or other products containing chlorpyrifos per crop season.
- **DO NOT** apply more than 82.1 fluid ounces of this product per acre per season, based upon the maximum limit of 0.5 pound active ingredient bifenthrin per acre per season.
- **DO NOT** make a second application of this product or other product containing chlorpyrifos within 10 days of the first application.
- **DO NOT** allow meat or dairy animals to graze in treated areas.
- **DO NOT** feed gin trash or treated forage to meat or dairy animals.

Cont'd. next page

15/19

**MATCH-UP™ INSECTICIDE**  
**EPA REG. NO. 34704-1086**

**Cotton cont'd.:**

- **DO NOT** make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season.
- **DO NOT** apply within 14 days of harvest.

**LEGUME VEGETABLES† (SUCCULENT OR DRIED EXCEPT SOYBEANS)**  
**(NOT FOR USE IN MISSISSIPPI)**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

† Adzuki bean, Bean, Blackeyed Pea, Broad bean (dry and succulent), Catjang, Chickpea, Cowpea, Crowder pea, English pea, Field bean, Field pea, Garden pea, Grain Lupin, Green pea, Guar, Lima bean (dry and green), Kidney Bean, Lablab bean, Lentil, Moth bean, Navy bean, Mung bean, Pea, Pigeon pea, Pinto bean, Rice bean, Southern pea, Sweet lupin, Tepary bean, Urd bean, White lupin, White sweet lupin

<b>Pest</b>	<b>Rate/A</b>	<b>Use Instructions</b>
Aster leafhopper Flea beetle Leafhoppers	2.05 to 16.4 fl oz	Apply in a minimum of 2.0 gal of finished spray/A by air or in a minimum of 10.0 gal/A with ground equipment. When applying by air, 1.0 to 2.0 qt of emulsified oil may be substituted for 1.0 to 2.0 qt of water in the finished spray. Thorough coverage is essential to achieve control.
Alfalfa caterpillar Aphids Bean leaf beetle Beet armyworm Cloverworm Corn earworm Corn rootworm (adults) Cucumber beetles Cutworms European corn borer Fall armyworm Grasshoppers Imported cabbageworm Japanese beetle (adults) Leaf miner Loopers Mexican bean beetle Pea leaf weevil Pea weevil Plant bug Saltmarsh caterpillar Sap beetle Southern armyworm Stink bugs Tarnished plant bug Thrips Tobacco budworm Webworms Western bean cutworm Whitefly Yellowstriped armyworm	2.7 to 16.4 fl oz	
Banks grass mite Carmine mite <i>Lygus</i> spp. Twospotted spider mite	6.6 to 16.4 fl oz	

**Use Restrictions:**

- **DO NOT** make more than 3 applications per year of Match-Up Insecticide or other products containing chlorpyrifos.
- **DO NOT** apply more than 32.8 fluid ounces (based on 0.2 pound active ingredient bifenthrin) to peas, or 49.3 fluid ounces (based on 0.3 pound active ingredient bifenthrin) to beans per acre per season.
- **DO NOT** make applications less than 14 days apart.
- **DO NOT** apply within 14 days of harvest.

16/19

**MATCH-UP™ INSECTICIDE**  
**EPA REG. NO. 34704-1086**

**PEANUT**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

<b>Pest</b>	<b>Rate/A</b>	<b>Use Instructions</b>
Beet armyworm Corn earworm Cutworm spp. Fall armyworm Grasshoppers Green cloverworm Leafhoppers Lesser cornstalk borer Loopers Rednecked peanut worm Southern armyworm Southern corn rootworm Stink bugs Threecornered alfalfa hopper Velvetbean caterpillar Yellowstriped armyworm	2.7 to 16.4 fl oz	Apply in a minimum of 10.0 gal/A with ground equipment or 2.0 gal/A by aircraft.
Aphids Spider mites Thrips Whitefly	6.6 to 16.4 fl oz	

**Use Restrictions:**

- **DO NOT** apply more than 82.1 fluid ounces (based on 0.5 pound active ingredient bifenthrin) per acre per season.
- **DO NOT** apply within 21 days of harvest.
- **DO NOT** feed green immature plants and peanut hay to livestock.
- To maintain a proper spray interval, do not make applications less than 14 days apart.
- Aerial application to peanuts is prohibited in Mississippi.

**SOYBEAN**

**(NOT FOR USE IN MISSISSIPPI)**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Apply as a postemergence broadcast spray using sufficient spray volume to ensure thorough coverage of treated plants, but no less than 10.0 gallons per acre at a rate of up to 16.4 fluid ounces of Match-Up Insecticide for ground spray equipment or 2.0 to 5.0 gallons per acre for aircraft equipment.

Apply when field counts indicate damaging pest populations are developing or present and then at a minimum of 30-day intervals.

This product may be tank mixed with glyphosate products when application is to be made to glyphosate-tolerant soybeans. Use a higher rate in the rate range when there is increased pest pressure.

**Chemigation:** This product may be applied through sprinkler irrigation systems at instructed broadcast application rates to control listed foliar pests. See Chemigation section for application instructions.

*Cont'd. next page*



17/19

**MATCH-UP™ INSECTICIDE**  
**EPA REG. NO. 34704-1086**

***Soybean cont'd.:***

<b>Pest</b>	<b>Rate/A</b>	<b>Use Instructions</b>
Alfalfa caterpillar	5.5 to 16.4 fl oz	On determinate soybeans, do not make more than 1 application after pod set.
Aphids		
Aster leafhopper		
Bean leaf beetle		
Beet armyworm		
Cloverworm		
Corn earworm		
Corn rootworm (adults)		
Cowpea curculio		
Cucumber beetles		
Cutworms		
Dectes stem borer		
European corn borer		
Fall armyworm		
Flea beetle		
Grasshoppers		
Green cloverworm		
Imported cabbageworm		
Japanese beetle (adults)		
Leafhoppers		
Leaf miner		
Lesser cornstalk borer		
Loopers		
Mexican bean beetle (adults)		
Pea leaf weevil		
Pea weevil		
Plant bug		
Saltmarsh caterpillar		
Sap beetle		
Southern armyworm		
Stink bugs		
Tarnished plant bug		
Thrips		
Tobacco budworm		
Velvetbean caterpillar		
Webworms		
Western bean cutworm		
Whitefly		
Woollybear caterpillar		
Yellowstriped armyworm		
Kudzu bug	13.2 to 16.4 fl oz	
<i>Lygus</i> spp.		
Twospotted spider mite		
Whitefly		
Velvetbean caterpillar		

**Use Restrictions:**

- **DO NOT** make more than 3 applications per year of Match-Up Insecticide or other product containing chlorpyrifos.
- **DO NOT** apply more than 49.2 fluid ounces of this product per acre per season, based upon a maximum limit of 0.3 pound active ingredient bifenthrin per acre per season.
- Maximum single application rate is 49.2 fluid ounces of Match-Up Insecticide.
- **DO NOT** make a second application of this product or other product containing chlorpyrifos within 14 days of the first application.
- **DO NOT** allow meat or dairy animals to graze in treated areas or otherwise feed treated soybean forage, hay, and straw to meat or dairy animals.
- **DO NOT** apply within 28 days of harvest.

12/19

**MATCH-UP™ INSECTICIDE**  
**EPA REG. NO. 34704-1086**

**TOBACCO**  
**(Preplant Soil Application)**

**Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Apply as a preplant broadcast spray to reduce the feeding damage caused by listed pests. Apply 24 to 48 hours before bedding and transplanting using a spray volume of 10.0 GPA or more. Incorporate immediately after application to a depth of 4 inches using suitable incorporation equipment.

Pest	Rate/A	Use Instructions
Armyworm spp. Cutworm spp. Mole crickets Stalk borers Tobacco flea beetle (larvae) Wireworms White grubs	10.3 to 16.4 fl oz	Before broadcast application of this product onto existing beds, knock down beds to final shape for transplanting. Use PTO-driven implements that will incorporate this product to a depth of 4 inches.
Aphid spp. Armyworm spp. Chinch bugs Cucumber beetle Cutworm spp. Flea beetle (adults) Grasshoppers Green bugs Japanese beetles Saltwater caterpillar Stink bugs Tarnished plant bugs Thrips Tobacco budworm Tobacco hornworm Whiteflies	6.6 to 16.4 fl oz	
<i>Lygus</i> spp. Spider mites	16.4 fl oz	

**Use Restriction:**

• **DO NOT** make more than 1 application of Match-Up Insecticide or other product containing chlorpyrifos per season.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL: Nonrefillable container.** Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at [www.acrecycle.org](http://www.acrecycle.org). If not recycled, then puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**For packages up to 5 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix

19/19

**MATCH-UP™ INSECTICIDE  
EPA REG. NO. 34704-1086**

**Storage & Disposal cont'd.:**

tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For packages greater than 56 gallons:** To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**For refillable containers:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC - 1-800-424-9300.**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Evik is a registered trademark of Syngenta Group Company  
Lightning is a registered trademark of BASF  
Match-Up is a trademark of Loveland Products, Inc.  
Steadfast is a registered trademark of E.I. duPont de Nemours & Company

**FORMULATED FOR  
LOVELAND PRODUCTS, INC.  
P.O. BOX 1286, GREELEY, COLORADO 80632-1286**